

Pycnodysostose Associated with Stridor: Case Report

Nadia Mebrouk ^{a*}, Rachid Abilkassem ^b and Aomar Agadr ^b

^a Nadia Mebrouk, Physician Resident in Pediatrics, Children's Hospital Rabat, Morocco.

^b Department of Pediatrics, Faculty of Medicine and Pharmacy, Military Teaching Hospital Mohammed V, University Mohammed V, Rabat, Morocco.

Authors' contributions

This work was carried out in collaboration among all authors. NM, RA and AA designed the study, performed the statistical analysis, wrote the protocol, wrote the first draft of the manuscript, and managed the analyses of the study. NM, RA and AA managed the literature searches. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJPR/2022/v8i430249

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/86050>

Received 12 February 2022

Accepted 20 April 2022

Published 27 April 2022

Case Study

ABSTRACT

Pycnodysostosis is a rare genetic disease, caused by a mutation of the cathepsin K gene which is involved in bone renewal. It is associated with dwarfism and bone fragility, but the association with a stridor is exceptional. We report the case of a child treated for stridor with sleep apnea syndrome confirmed by sleep recording. His clinical examination showed stridor with laryngomalacia, the characteristic dysmorphism of pycnodysostosis and failure to thrive. The patient was put under non-invasive ventilation during evenings, and underwent ablation of adenoids, which resulted in a stable blood oxygen saturation. A surgery to lengthen the patient's mandibular is being considered, depending on her evolution. Practitioners should overlook this rare disease when faced with atypical stridor associated with high obstruction, failure to thrive and dysmorphism.

Keywords: Pycnodysostosis; stridor; sleep apnea syndrome.

1. INTRODUCTION

Pycnodysostosis or Toulouse-Lautrec disease is a rare autosomal recessive disease of unknown

exact prevalence (between 1/100,000 and 1/1 million) [1], classified as an osteochondrodysplasia. It affects both sexes equally and is caused by a 1q21 mutation in the

*Corresponding author: Email: nadia.mebrouk@hotmail.com;

cathepsin K gene [2]. The absence of this protein, secreted by the osteoclasts, leads to a defect in the cleavage of the proteins of the bone matrix and the bone becomes abnormally fragile [3]. The arguments allowing the diagnosis to be made are: clinical, with growth retardation predominant in the limbs which are short and massive, fontanelle open or closing late, characteristic facial dysmorphism (macrocephaly, protrusion of frontal and occipital bumps, micrognathia, hypomaxillia with prominent nose, koilonychia, dental implantation defect with double row) [4]. The facial abnormalities of pycnodysostosis, such as micromandibuli or maxillary hypoplasia, are anatomical factors that can reduce the upper airway pathway. However, upper respiratory obstruction, leading to obstructive sleep apnea syndrome (OSAS), is rarely the main symptom of the disease [5–6]. We report a case of pycnodysostosis associated with stridor with OSAS.

2. CASE REPORT

The patient is a child aged two years and 4 months who was referred to our unit at two months of age for stridor. He is of first degree consanguineous parents. The interrogation revealed that the stridor dates back to birth. The examination found a dysmorphic syndrome made up of frontal hump, persistent anterior fontanelle, micrognathia, short fingers, with curved nails (Fig. 1), glossophtosis and elongation of the soft palate. Asymmetrical thorax, scoliotic attitude of the dorsal spine, with significant stature delay (-3

standard deviations), cranial circumference 41 cm (+1 standard deviation). Chest CT angiography did not show any vascular abnormalities.

Otorhinolaryngological examination in favor of mild laryngomalacia. The skeletal radiographs showed densification of the bones of the base of the skull, persistence of the anterior fontanelle, diffuse condensation of the rib cage and pelvis with a dense appearance of the spine (Fig. 2), diaphyseal densification and metaphyseal long bones and hypoplastic phalanges (Fig. 3). A computed tomography of the skull and the facial bone showed mandibular hypoplasia (Fig. 4). In view of the clinical signs and radiological manifestations, the diagnosis retained is pycnodysostosis. Consultation with a geneticist subsequently confirmed the diagnosis of pycnodysostosis. Genetic counseling has been proposed for the family and the risk of recurrence, in the patient's couple, during subsequent pregnancies is 25%. The child is regularly taken care of in otolaryngology and orthopedics. Faced with chronic snoring with installation of obstructive sleep apnea syndrome (OSAS), a ventilatory polygraphy was performed and showed that the patient presents with obstructive sleep apnea hypopnea syndrome with a severe snoring index and desaturation index figured at 54.4/h. Our patient then presented the disease of pycnodysostosis with obstruction of the upper airways secondary to mandibular hypoplasia, the patient was then put on non-invasive ventilation during sleep, which made it possible to improve her saturation and reduce sleep apnea pending surgical treatment.



Fig. 1. short fingers, with curved nails



Fig. 2. diffuse condensation of the rib cage and pelvis with a dense appearance of the spine



Fig. 3. diaphyseal densification and metaphyseal long bones and hypoplastic phalanges



Fig. 4. Mandibular hypoplasia

3. DISCUSSION

Pycnodysostosis is a rare autosomal recessive lysosomal genetic disease [7]. The most common clinical manifestations are dwarfism, acroosteolysis of the distal phalanges, bone fragility with spontaneous fractures and dysplasia of the clavicles. The characteristic cranial malformations are a bulky skull with the presence of wormian bones and persistence of the anterior fontanel, a small mandible, decayed, poorly implanted or abnormally shaped teeth. The nails are sometimes irregular and cracked. Respiratory distress and sleep apnea are sometimes observed, at any age of life, but stridor is very rare. Diagnosis is clinical and should be confirmed by full skeletal radiographic examination and skull radiographs. A CTSK gene analysis may also be requested. The few respiratory obstructions associated with pycnodysostosis reported in the literature are located at different levels of the upper airways, at any age of life. Thus, in children, narrowing of the pharynx associated with glossoptosis has been described in two boys aged 4 and 5 years with pycnodysostosis [8]. Another case of narrow and obstructive cavum was reported in a 4-year-old boy [5]. Finally, mandibular hypoplasia was also implicated in the genesis of the obstructive syndrome in a 3-year-old boy [6]. In adults, a

cephalometric study in two sisters aged 70 and 67 localized the narrowing to the pharynx by the association of maxillary and mandibular hypoplasia, glossoptosis and an excessively long soft palate [9]. The management of respiratory obstructive symptoms in pycnodysostosis depends on the level and severity of the obstruction. Thus, maxillofacial surgery can enlarge the pharyngeal canal in cases of marked bone hypoplasia. Lengthening of the mandibular ramus by bilateral rib graft to propel the base of the tongue forward was performed in a 3-year-old boy [6]. A Lefort I osteotomy with maxillary distraction for 6 weeks was performed on a 15-year-old girl [10]. Non-invasive ventilation is an interesting therapeutic possibility in children with OSAS of multiple etiology, daytime hypercapnia is the most often accepted criterion for initiating non-invasive ventilation (NIV). The occurrence of acute respiratory failure, particularly during superinfection, for example, is also a recognized indication for NIV.

Pycnodysostosis is caused by mutations in the gene for cathepsin K (located in 1q21), a lysosomal enzyme secreted by osteoclasts which allows the cleavage of proteins from the bone matrix (collagen type I, osteonectin or osteopontin).

4. CONCLUSION

Faced with atypical stridor associated with high obstruction, failure to thrive and dysmorphism, it is important not to overlook a rare disease. Pycnodysostosis is one of them and its early diagnosis makes it possible to detect and prevent its complications, which are bone fractures, head trauma and the consequences on the development of the child of a sleep apnea syndrome. The level of respiratory obstruction is variable and its precise identification allows appropriate treatment, often combining surgery and non-invasive ventilation. As this genetic is autosomal recessive, genetic counseling is important.

CONSENT

As per international standard, parental written consent has been collected and preserved by the author(s).

ETHICAL APPROVAL

As per international standard or university standard written ethical approval has been collected and preserved by the author(s).

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Hodder A, Huntley C, Aronson KJ, Ramachandran M. Pycnodysostosis and the making of an artist. *Gene*. 2015;555(1):59-62. PubMed | Google Scholar
2. Puri R, Saxena A, Mittal A, Arshad Z, Dwivedi Y, Chand T, et al. Pycnodysostosis: an anaesthetic approach to this rare genetic disorder. *Anesthesiology*. 2013;2013:716756
3. Troen BR. The role of cathepsin K in normal bone resorption. *Drug News Perspect*. 2004;17:19–28.
4. Sudarshan R, Vijayabala GS. Pycnodysostosis'a review. *SEAJCRR*. 2012;1(2):42-5
5. Abourazzak S, Chaouki S, Akodad Z. Apnées de sommeil et retard statural révélant une pycnodysostose. *Arch Pediatr*. 2013;20:307–10.
6. Teissier N, Jacquemont ML, Blancal JP, et al. Severe snoring in a child with pycnodysostosis treated with a bilateral rib graft. *Cleft Palate Craniofac J*. 2009;46:93–6.
7. Mujawar Q, Naganoor R, Patil H, Thobbi AN, Ukkali S, Malagi N. Pycnodysostosis with unusual findings: a case report. *Case J*. 2009;2:6544
8. Yousefzadeh DK, Agha AS, Reinertson J. Radiographic studies of upper airway obstruction with cor pulmonale in a patient with pycnodysostosis. *Pediatr Radiol*. 1979;8:45–7.
9. Muto T, Yamazaki A, Takeda S, et al. Pharyngeal narrowing as a common feature in pycnodysostosis – a cephalometric study. *Int J Oral Maxillofac Surg*. 2005;34:680–5.
10. Nørholt SE, Bjerregaard J, Mosekilde L. Maxillary distraction osteogenesis in a patient with pycnodysostosis: A case report. *J Oral Maxillofac Surg*. 2004;62:1037–40.

© 2022 Mebrouk et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/86050>