



## Multi-Drug Resistant Bacterial Status of Borehole Water in Kogi State University, Anyigba, Nigeria

S. A. Adegoke <sup>a\*</sup>

<sup>a</sup> Department of Microbiology, Kogi State University, Anyigba, Nigeria.

### Author's contribution

The sole author designed, analysed, interpreted and prepared the manuscript.

### Article Information

DOI: 10.9734/AJOB/2022/v16i1293

### Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/91512>

Original Research Article

Received 07 July 2022  
Accepted 13 September 2022  
Published 29 September 2022

### ABSTRACT

Transmission of pathogens through water is of grave public health concern. Bacteria are of major concern because of the pathogenicity and etiologic agents of life threatening infections. The multidrug resistant (MDR) bacterial status of the borehole water samples from Kogi State University, Anyigba was studied. The samples ten each from point A and B were collected from two main borehole water sources and analyzed for MDR bacteria. A total of seven isolates (*Escherichia coli*, *Bacillus* sp, *Klebsiella* sp, *Pseudomonas* sp, *Staphylococcus* sp, *Streptococcus* sp and *Salmonella* sp) were identified by standard microbiological methods. Phenotypic identification of antibiotic resistance profile using the disk diffusion method was carried out. *Pseudomonas* sp, *Streptococcus* sp, *Staphylococcus* sp, *Escherichia coli* and *Klebsiella* sp. were found to be 100% resistant to all the imported antibiotics while 55.6% and 66.7% resistance were recorded for *Salmonella* sp. and *Bacillus* sp respectively. Percentage resistance to all indigenous antibiotics recorded were 40% for *Pseudomonas* sp., 30% for *Salmonella* sp, 40% for *Escherichia coli*, 30% for *Klebsiella* sp, 20% for *Staphylococcus* sp, 20% for *Streptococcus* sp, and 10% for *Bacillus* sp. The results showed that all the isolates were multi drug resistant (MDR) and the presence of these organisms poses great risk to the university community as well as individuals that consume the water and use for other domestic purposes.

\*Corresponding author: E-mail: detunjioutreach68@gmail.com;

*Keywords: Antibiotics; bacteria; borehole; multi-drug resistant; pollution and waste.*

## 1. INTRODUCTION

Potable water is an essential ingredient for good health and the socio-economic development of man [1] but it is lacking in many societies. Clean water is priceless and a limited resource that man has begun to treasure only recently after decades of pollution and waste [2]. World population cannot be sustained without access to safe water [3]. It is therefore important to conjunctly consider both water quality and quantity in water resources management [4]. Borehole water become unsuitable for domestic use as a resource due to contamination that makes it unfit for many purposes [5]. Standards and guidelines in water quality stem from the need to protect human health. Borehole water serves as the major source of drinking water in the local population of Nigeria. Since only few can afford and rely on purified and treated bottled water for consumption. Chowdhury [6] stressed the importance of groundwater as a source of potable water in Africa and constitutes about two thirds of the freshwater resources of the world. Ground water provides a reasonably constant supply for domestic use, livestock and irrigation. (Carlow et al., 2011) stated this source can buffer the effects of rainfall variability across seasons. In many arid and semi-arid areas of Africa, boreholes water is a means of coping with water deficiencies in areas where rainfall is scarce or highly seasonal and surface water is extremely limited [5].

Underground water naturally is water from the atmosphere and earth surface that has percolated through the coarse particulate network of the soil, which serves as a natural filter against microbial contaminants, into the water table and thus is expected to be of better quality than the surface water. Underground water is gradually becoming over exploited in Nigeria due to the inability of the government to provide adequate potable water supply to all communities in the country [7]. There are several rules guiding the digging of wells, but because of poverty some cannot afford the sinking of deeper wells called boreholes and thus employ the services of local well diggers who most times hand-dig wells irrationally and even sometimes closer to locations like soak away pit, latrine, and sewage septic tanks that usually encourages microbial contamination of such wells [7]. Some of these hand-dug wells are not hygienically

protected and are never treated from time to time [7]. Most are made in a way that provides easy access to reptiles and insects that defecate or sometimes get drowned in the wells. The manner in which some even draw water from the wells encourages contaminations through the rope and containers being used. The world health organization (WHO) has defined portable water as water in which the physical, chemical and microbiological quality is within acceptable limit [8]. However, the truth is, over one billion people worldwide have no access to potable water [9]. This has resulted to increased cases and spread of waterborne diseases throughout the world (Obeta, 2013). WHO reported an estimation of over two million deaths as a result of waterborne diseases and over four billion diarrhea cases worldwide annually. In Africa, the WHO has estimated that a child has five episodes of diarrhea in a year with about 800, 000 deaths of children per year from diarrhea and dehydration [9]. These have been attributed to the presence of bacterial pathogens in the drinking water which resulted in various waterborne diseases such as cholera, typhoid fever, bacillary dysenteries and many gastrointestinal diseases [10]. Drinking of water contaminated with human and animal faeces exposes individuals to high risk of microbial infections, especially faeces from infected or carriers of waterborne disease-causing agents.

Contamination of water bodies has increasingly become an issue of serious environmental concern. In the case of underground waters like bore holes, this may arise from the construction process of a borehole, drilling fluids, chemical casings and other materials which may find their way into the well thereby polluting the water [11]. An open well during the construction stage can also be a direct route for contaminants from the surface to the aquifer thereby providing an ideal opportunity for chemical casing and bacteriological pollution to occur [3]. Even if no source of anthropogenic contamination may exist, there is potential for natural levels of metals and other chemicals to be harmful to human health.

The occurrence and spread of antibiotic-resistant bacteria (ARB) are pressing public health challenges worldwide, and aquatic ecosystems are recognized reservoirs for ARB as well as antibiotic resistance genes (ARGs). The

emergence of antibiotic-resistant bacteria limits the clinical use of antibiotics. So, there is increasing concern that existing antibiotics would not be potent any longer against these pathogens. It is in view of this that the borehole water sources from Kogi State University Ayingba were assessed to ascertain the multi-drug resistant bacteria status of the water.

## 2. MATERIALS AND METHODS

**Sample collection:** The borehole water samples were collected randomly from the university two main borehole water sources in sterile screw cap bottle and held in iceboxes until delivery to the laboratory.

**Isolation and enumeration:** Serial dilution was carried out for each of the borehole water samples before inoculation on Nutrient agar and MacConkey agar. One ml of the borehole water was diluted in 9ml to give tenfold serial dilution. With the aid of a sterile syringe, 1ml of the sample was aseptically transferred from the various dilution factors into the sterile nutrient agar Petri dishes. Enumeration was done using the pour plate method.

**Characterization and identification of bacteria:** Identification and characterization of bacteria was done by isolation of bacteria in a pure form by streaking on nutrient agar to obtain a pure isolate and Gram staining was carried out as well as other biochemical tests contained in the standard diagnostic protocol (ISO 6885-1; 1999). The biochemical tests that were carried out include.

**Catalase test:** This test demonstrates the presence of catalase, an enzyme that catalyses the release of oxygen from hydrogen peroxide. The enzyme catalase mediates the breakdown of hydrogen peroxide into water and oxygen. The presence of the enzyme in a bacterial isolate is evident when a small inoculum is introduced into hydrogen peroxide, and the rapid elaboration of oxygen bubbles. The lack of catalase is evident by lack of or weak bubble production.

The test was carried out by transferring a small amount of colony growth into a slide; a drop of 3% hydrogen peroxide was placed on the organism and observed for immediate bubbling.

**Citrate test:** This test detects the ability of an organism to use citrate as the sole source of carbon and energy. Use of citrate involves the enzyme citrase, which breaks down citrate to oxaloacetate and acetate. Oxaloacetate is further broken down to pyruvate and carbondioxide.

Production of sodium bicarbonate as well as ammonia from the use of sodium citrate and ammonium salts results in alkaline pH. This results in a colour change. Bacterial colonies were picked and inoculated into a slope of Simmons citrate agar and incubated overnight at 37 degree Celsius. The organism that has the ability to use citrate, changes the color of the medium from green to blue.

**Urease test:** Urea is a di-amide of carbonic acid. It is hydrolyzed with the release of ammonia and carbon dioxide. Many organisms, especially those that infect the urinary tract, have a urease enzyme which was able to split urea in the presence of water to release ammonia and carbon dioxide. The ammonia combines with carbon dioxide and water to form ammonium carbonate which turns the medium alkaline, turning the indicator phenol red from its original yellow color to bright pink.

**Indole test:** Indole was performed on bacteria species to determine the ability of the organism to convert tryptophan into indole. The pure bacteria culture was grown in peptone broth for 24 hours. After incubation, 5 drops of kovac's reagent was added to the culture broth. Positive result shows the presence of a red or red violet color in the surface alcohol layer of the broth, negative result appears yellow.

**Methyl Red:** It was used to identify and characterize enteric bacteria based on their pattern of glucose metabolism. An isolate was inoculated into a tube with a sterile transfer loop; the tube was incubated at 35 degrees Celsius for 2 days. After incubation, 2.5ml of the medium was transferred to another tube. Five drops of pH indicator methyl red is added to this tube, the tube was gently rolled between the palms of the hand to disperse the methyl red. A red color represents a positive test that is; they subsequently metabolize pyruvic acid to other acids. A yellow color represents a negative result.

**Voges-Proskauer test:** Alpha-naphthol and potassium hydroxide was added to the incubated bacteria (VP broth). A cherry red color indicates a positive result while a yellow-brown color indicates a negative result.

**Multiple tube fermentation technique:** Coliforms are detected in 3 stages: presumptive, confirmed and completed. In the presumptive test, dilutions from the water sample are added to tubes of lactose broth medium with a pH indicator and incubated for 24 to 48 hours. After

the period of time, the development of gas and how much produced was examined. In the confirmed test, samples are streaked from the positive presumptive tubes at the highest dilutions onto plates of differential media, eosin methylene blue (EMB) agar that contains lactose. Since coliforms produce acid from lactose, and the eosin methylene blue dyes are absorbed under acid conditions, the coliforms form dark-centered colonies with or without metallic sheen; these colonies indicate a positive confirmed test. In the completed test, separate colonies are selected from the confirmed test and inoculated in lactose broth and on a nutrient agar slant for 24 hours at 35°C. If gas and acid are produced in lactose broth, and the isolated microorganism is a Gram negative non-endospore-forming rod, it indicates a positive completed test.

**Anti bacteria susceptibility testing:** The Kirby-Bauer method was adopted for the antibacterial susceptibility. If the bacteria are using the Mueller-Hinton agar. The isolates in broth form were poured on the molten agar and allowed to solidify before application of the antibiotics on the plates. Sensitivity to the antibiotic by the bacteria was indicated with a clear ring or a zone of inhibition seen around the disc. A ruler was used to measure the diameter of the disk plus the surrounding clear area in millimeters (mm).

### 3. RESULTS

The phenotypic identification revealed seven bacteria genera (Table 1). The colonial edges of the organisms were mostly entire and most with yellow colour. All were subjected to biochemical characterization.

Table 2 shows the biochemical characterization of the seven bacterial genera from the borehole water. Gram negative species were present more than the Gram positive. The organisms were predominantly rod shaped.

*Pseudomonas* sp. and *Klebsiella* sp. showed 40% resistance to gentamicin and nalidixic acid (Table 3) while *Salmonella* sp. and *E. coli* had 30%. Both had similar resistance pattern except for augmentin, ampicillin, aminoglycosides (gentamicin and streptomycin) (Table 3).

The isolates were largely susceptible to the indigenous antibiotic disk Table 4, with% susceptibility range of 50 – 70%. The resistance percentage range was 10% - 20% as shown in Table 4.

The resistance was more pronounced with the imported antibiotic disk as shown in Table 5. The % resistance range was 55.6% - 100%. Only *Streptococcus* had an intermediate resistance.

**Table 1. Colonial morphology of isolates**

Samples	Shape	Colour	Elevation	Edges	Consistency	Probable organism	
A	1	Round	Blue green	Convex	Entire	Dry	<i>Pseudomonas</i> sp
	2	Round	Milky	Flat	Entire	Moist	<i>Streptococcus</i> sp
	3	Round	Yellow	Raised	Entire	Moist	<i>Salmonella</i> sp
	4	Round	Milky	Convex	Entire	Moist	<i>Staphylococcus</i> sp
B	1	Circular	Yellow	Convex	Entire	Moist	<i>Escherichia coli</i>
	2	Round	Pink	Flat	Entire	Mucoid	<i>Klebsiella</i> sp
	3	Round	Yellow	Raised	Undulate	Mucoid	<i>Bacillus</i> sp

**Table 2. Gram Reaction and Biochemical Tests of Isolates**

Sample	Gram reaction shape		Biochemical tests						Probable microorganisms
			IND	MR	VP	CAT	CIT	URE	
A1	-	ROD	+ve	-ve	+ve	+ve	+ve	+ve	<i>Pseudomonas sp</i>
A2	+	COCCI	-ve	+ve	-ve	-ve	+ve	-ve	<i>Streptococcus sp</i>
A3	-	ROD	-ve	-ve	-ve	+ve	-ve	-ve	<i>Salmonella sp</i>
A4	+	COCCI	-ve	+ve	-ve	+ve	+ve	-ve	<i>Staphylococcus sp</i>
B1	-	ROD	+ve	+ve	-ve	+ve	-ve	+ve	<i>Escherichia coli</i>
B2	-	ROD	-ve	-ve	+ve	+ve	+ve	+ve	<i>Klebsiella sp</i>
B3	+	ROD	-ve	+ve	-ve	-ve	+ve	-ve	<i>Bacillus sp</i>

Key: IND= indole MR= Methyl red VP- Vogesproskaner, CAT- catalase, CIT= citrate and URE= urease

**Table 3. Antibiotic susceptibility test on set 1 (gram negative)**

<b>Chemical class of antibiotics</b>	<b>Antibiotics</b>	<b><i>Pseudomonas</i> sp.</b>	<b><i>Salmonella</i> sp.</b>	<b><i>E. coli</i></b>	<b><i>Klebsiella</i> sp.</b>
Beta Lactam	PEF (10µg)	S	S	S	S
	AU(20µg)	R	R	I	I
	PN(20µg)	I	I	S	R
Aminoglycosides	CN(10µg)	R	I	R	R
	S(30µg)	R	I	R	S
Fluoroquinolones	OFX(10µg)	S	S	S	S
	CPX(10µg)	S	S	S	S
	NA(30µg)	R	R	R	R
Cephalosporin	CEP(10µg)	S	R	R	I
Sulfonamides	SXT(30µg)	S	S	S	I
	Susceptible	50	40	50	40
Percentage (%)	Intermediate	10	30	10	30
	Resistant	40	30	40	30

Key: R=resistance, I= intermediate, S= susceptible, OFX= tarivid, CPX = ciproflox, AU= augmentin, CN=gentamycin, S = streptomycin, CEP= Ceporex , NA= Nalidixic acid, SXT= septrin, PN, ampicillin

**Table 4. Antibiotic susceptibility test on set 2 (gram positive)**

<b>Chemical class of antibiotics</b>	<b>Antibiotics</b>	<b><i>Staphylococcus</i> sp</b>	<b><i>Streptococcus</i> sp</b>	<b><i>Bacillus</i> sp</b>
Fluoroquinolones	CPX(10µg)	S	S	S
	NB(10µg)	R	I	R
	LEV(20µg)	S	S	S
Aminoglycosides	CN(10µg)	S	S	I
	S(30µg)	S	S	S
Macrolide	E(30µg)	R	S	S
Beta Lactam	AML(20µg)	I	R	S
	APX(20µg)	I	R	I
Antitubercular	RD(20µg)	S	I	S
Chloramphenicol	CH(30µg)	S	I	S
Percentage (%)	Susceptible	60	50	70
	Intermediate	20	30	20
	Resistant	20	20	10

Key: R= resistant, S= susceptible, I= intermediate, CPX= Ciproflox, NB= Norfloxacin, CN= gentamycin, AMX= amoxyl S= streptomycin, RD= rifampicin, E= erythromycin, CH=chloramphenicol, APX= ampiclox, LEV=levofloxacin.

**Table 5. Antibiotic susceptibility test on the imported DISC**

<b>Chemical class of antibiotics</b>	<b>Antibiotics</b>	<b><i>Pseudomonas</i> sp</b>	<b><i>Streptococcus</i> sp</b>	<b><i>Salmonella</i> sp</b>	<b><i>Staphylococcus</i> sp</b>	<b><i>E. coli</i></b>	<b><i>Klebsiella</i> sp</b>	<b><i>Bacillus</i> sp</b>
B-Lactam	Amp (10µg)	R	R	R	R	R	R	R
	OX(1µg)	R	R	R	R	R	R	R
	AUG(30µg)	R	R	R	R	R	R	R
Macrolide	VAM(30µg)	R	R	I	R	R	R	S
Cephalosporin	CAZ(30µg)	R	R	R	R	R	R	R
	CTX(30µg)	R	R	R	R	R	R	R
Tetracycline	TE(30µg)	R	R	I	R	R	R	R
	DO(30µg)	R	R	I	R	R	R	S
chloramphenicol	C(30µg)	R	R	I	R	R	R	S
Percentage (%)	Susceptible	0	0	0	0	0	0	33.3
	Intermediate	0	0	44.4	0	0	0	0
	Resistant	100	100	55.6	100	100	100	66.7

Key: CTX= cefotaxime, CAZ= ceftazidime, AMP= ampicillin, OX= oxacillin, VAN= vancomycin, TE= tetracycline, DO= doxycycline, C= chloramphenicol. Aug= amoxicillin



Table 6. MPN result of borehole water

Sample	Number of tubes giving a positive result			MPN (per 100ml)	95% confidence limit	
	10ml	1ml	0.1ml		Low	high
A	5	3	0	79	25	190
B	3	2	1	140	37	340

Table 6 shows the results of the most probable number (MPN) in the enumeration of the microorganisms. It shows high number of bacterial counts per millilitres of the water sample.

#### 4. DISCUSSION

This study was conducted to assess the multidrug resistant bacteria in the borehole water of Kogi State University Anyigba. A total of seven isolates were enumerated and characterized from the borehole water sampled. Bacteria isolated were *Escherichia coli*, *Klebsiella* sp, *Staphylococcus aureus*, *Salmonella* sp, *Bacillus* sp, *Streptococcus* sp and *Pseudomonas* sp. These organisms pose several health risks to immune compromised individuals in particular and consumers in general [12]. The presence of *Klebsiella* sp in the borehole water is unacceptable from the public health point of view and it agrees with the findings of Ogu et al. [13].

Microorganisms isolated were identified with Gram staining; three isolates (*Streptococcus* sp, *Staphylococcus* sp, and *Bacillus* sp) were Gram positive while four isolates (*Pseudomonas* sp, *Salmonella* sp, *Escherichia coli*, *Klebsiella* sp) were Gram negative.

Further identification with biochemical tests were carried out on the isolates. *Pseudomonas* sp was positive for catalase, citrate, urease, vogesproskauer, indole test but negative for methyl red. *Streptococcus* sp was positive for methyl red and citrate test but negative for indole, vogesproskauer, catalase and urease test. *Salmonella* sp was positive for catalase and urease test but negative for indole, methyl red, vogesproskauer and citrate test. *Staphylococcus* sp was positive for methyl red, catalase, citrate but negative for urease, voges-proskauer and indole test. *Escherichia coli* were negative for voges-proskauer and citrate but positive for indole, catalase, methyl red, and urease. *Klebsiella* sp was negative for indole and methyl

red but positive to catalase, citrate, urease, voges-proskauer. *Bacillus* sp was positive to methyl red and citrate but negative to catalase, urease, voges-proskauer and indole.

Antibiotic susceptibility test was carried out for each isolate on an indigenous and imported disk. The percentage of resistance of the Gram negative isolates on the indigenous disk ranges between 30-40%, intermediate ranges between 10-30% and the susceptibility ranges between 40-50%. The percentage of resistance of the Gram positive isolates on the indigenous disc ranges between 10-20%, intermediate between 20-30% while susceptibility ranges from 50-70%. Most of the isolates had 100% resistance on the foreign disk.

The multiple antibiotic resistances of *E. coli* established in this study agreed with other findings [14]. Strains of *E. coli* and *Salmonella* spp. accounted for several outbreaks worldwide, partly due to resistance to chloramphenicol, ampicillin, and trimethoprim [15].

The frequency of resistance to penicillin in the current study was high among the isolates as compared with resistance to chloramphenicol and ampicillin observed in the isolates obtained from the various water sources. *E. coli* resistance against ampicillin was observed by Çelebi et al. [16] and Olowe et al., [17]. Lastly, periodic monitoring of antibiotic sensitivity of the water sources is of importance to detect any changing patterns that may arise in future in order to keep pace with such changing patterns for better curative measures or policies formulation and implementation.

For the MPN (most probable number) method, the maximum acceptable concentration for drinking water is none detectable per 100ml and the samples exceeded this, which means that the borehole water is unsafe for drinking without further treatment and this work agrees with the

work of Okafor, [18] on borehole water in Nsukka [19,20].

## 5. CONCLUSION

It can be deduced that most bacteria found in the borehole water of Kogi State University Ayingba are multidrug resistant because they showed resistance that cut across different classes of antibiotics. On this note, water treatments and regular washing of overhead tanks if practice would help in reducing the menace of spread of multidrug resistance bacteria through the use of borehole water. There is need to pay attention to surface aquifers to check pollution as this could be the source of the multidrug resistance organisms.

## COMPETING INTERESTS

Author has declared that no competing interests exist.

## REFERENCES

1. Udom GJ, Ushie FA, Esu EO, Ofojekwu PC, Ezenwaka IS, Alegbeleye WO, et al. A geochemical survey of groundwater in Khana and Gokana local government area of Rivers State, Nigeria. *J Appl Sci Environ Manag.* 2002;6(1):53-9. DOI: 10.4314/jasem.v6i1.17196
2. Sinderberg MS. Higher education chemistry. In: *The molecular nature of matter and change.* McGraw-Hill; 2003.
3. Brainstein J. Trading the Rain, Should the World's fresh water resources be an international traded commodity? In: *integrated water resources management and security in the middle east.* NATO Science for and security series Springer. Dordrecht; 2007. Available: <https://doi.org/10.1007/978-1-4020-5986-12>.
4. Xinghui X, Zhifeng Y, Yuxiang W. Incorporating eco-environmental water requirements in integrated evaluation of water quality and quantity- A case study for the Yellow River. *Water Resour Manag.* 2009;23:1067-79.
5. Agbaire PO, Oyibi PI. Seasonal variation of some Physico-chemical properties of borehole water in Abraka, Nigeria. *Afr J Pure Appl Chem.* 2009;3(6):116-8.
6. Chowdhury S, Mazumder MAJ, Al-Attas O, Husain T. Heavy metals in drinking water: occurrences, implications, and future needs in developing countries. *Sci Total Environ.* 2016;569-570:476-88. DOI: 10.1016/j.scitotenv.2016.06.166, PMID 27355520.
7. Ayantobo OO, Oluwasanya GO, Idowu OA, Eruola AO. Water quality evaluation of hand-dug wells in Ibadan, Oyo State, Nigeria. *Glob J Sci.* 2013;13(10):21-7.
8. WHO. Guidelines for drinking-water quality, health criteria and other supporting information. 2nd ed. Vol. 2004. Geneva: World Health Organization. 2004;2.
9. WHO. Water Supply and Sanitation Council, Global water supply and sanitation assessment report 2000. New York; 2000.
10. Adeyinka SY, Wasiru J, Akintayo CO. Review on prevalence of waterborne diseases in Nigeria. *J Adv Life Sci.* 2014; 1(2):1-3.
11. Angulo FJ, Tippen S, Sharp DJ, Payne BJ, Collier C, Hill JE, et al. A community waterborne outbreak of salmonellosis and the effectiveness of a boil water order. *Am J Public Health.* 1997;87(4):580-4. DOI: 10.2105/ajph.87.4.580, PMID 9146435.
12. Iroha C, Iroha I, Nwakeze E, Ajah M, Ejike C. Evaluation of the metal content and bacteriological parameters of selected borehole water sources in Abakaliki, Nigeria. *Int J Waste Resour.* 2018;8:338.
13. Ogu GI, Madar IH, Olueh AA, Tayubi IA. Antibiotic susceptibility profile of bacteria isolated from drinking water sources in Amai Kingdom, Delta State, Nigeria. *Annu Res Rev Biol.* 2017;14(1):1-9. DOI: 10.9734/ARRB/2017/34326
14. Park J, Kim JS, Kim S, Shin E, Oh K, Kim Y, et al. A waterborne outbreak of multiple diarrhoeagenic *Escherichia coli* infections associated with drinking water at a school camp. *Int J Infect Dis.* 2018;66:45-50. DOI: 10.1016/j.ijid.2017.09.021
15. Uma B, Prabhakar K, Rajendran S, Kavitha K, Sarayu YL. Antibiotic sensitivity and plasmid profiles of *Escherichia coli* isolated from pediatric diarrhea. *Journal of Global Infectious Diseases.* 2009;1(2): 107.
16. Çelebi A, Duran N, Öztürk F, Açıık L, Aslan G, Aslantaş O. Identification of clinic uropathogen *Escherichia coli* isolates by antibiotic susceptibility, plasmid and whole cell protein profiles. *Advanced Molecular Biology.* 2007;1:31–40.

17. Olowe OA, Okanlawon BM, Olowe RA, Olayemi AB. Antimicrobial resistant pattern of Escherichia coli from human clinical samples in Osogbo, south western Nigeria. African Journal of Microbiology Research. 2008;2(1):8-11.
18. Okafor N. Aquatic and waste Microbiology (1st ed.). Fourth Dimension Publishing Coy.P. Enugu. 1985;52.
19. Roger CC, McDonald AM, Nicol AL, Robins NS. Groundwater security and drought in Africa. Linking Availability, Access and Demand. 2011. DOI: 10.1111/j.1745-6584.2009.00558.x
20. Obeta MC, Ocheje FJ. Assessment of groundwater quality in Ankpa Urban, Kogi State, Nigeria. Environ Res J. 2003; 7(3):37-47.

---

© 2022 Adegoke; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/91512>