



Resident Perceptions on the Health Risks Associated with Solid Waste Disposal at Kalundu Dumpsite, Kitui County – Kenya on the Surrounding Human Settlement

Valilian Kavutha Matheka ^{a*}, Bonface O. Manono ^a and Felista Muriu-Ng'ang'a ^a

^a South Eastern Kenya University, P.O.Box 170 – 90200, Kitui, Kenya.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/AJEE/2022/v19i130335

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/89210>

Original Research Article

Received 15 May 2022
Accepted 20 July 2022
Published 01 August 2022

ABSTRACT

Aims: Urban centers of developing countries are growing at very rapid rates. This has been accompanied by increased use of resources and enormous waste generation. The generated waste is dumped in the outskirts of the urban where they can have negative impacts on the environment and surrounding human settlements. This study therefore aims to assess the perceived health risks of Kalundu dumpsite in Kitui Town, Kitui County of Kenya.

Study Design: A survey design was used to collect data from people living within 500 meter radius of Kalundu dumpsite.

Place and Duration of Study: This study was carried out between January 2020 and December 2020 in Kitui Town of Kitui County, Kenya.

Methodology: A Semi - structured questionnaire was administered to 78 respondents to collect data on perceived health risks they associate with the presence of Kalundu dumpsite in their locality. The collected data was coded and then analyzed. Pearson Correlation Coefficient was used to test the link between respondent characteristics and perceived health risks.

Results: 83.3% of respondents perceived the presence of the dumpsite in their vicinity as a health hazard that they attributed to cause diseases (Malaria, chest pain; diarrhea, and cholera) and other negative environmental effects (smoke, odour, and habitat for disease vectors and pathogens).

Children under the age of 5 years were the most vulnerable.

Conclusion: There is a perceived link between waste deposited onto Kalundu dumpsite and human and environmental health of the surrounding community.

Keywords: Dumpsite; health risks; solid waste; Kalundu; waste management; developing countries; Kitui-Kenya.

1. INTRODUCTION

In recent years, the world population is rapidly increasing with a majority of the increase experienced in urban areas [1]. In Africa, urban population is projected to increase from 395 million in 2009 to 1.23 billion persons in 2050 [2]. This has been characterised by an exponential increase in resource consumption and tremendous increase in waste generation [3]. This process has accelerated in Kenya since the inception of the devolved administration system in after the promulgation of the 2010 Constitution.

In many cases, the generated wastes are thrown into poorly managed waste disposal sites [4], located either within the town centres or on their outskirts [5,6]. To prevent detrimental effects to the environment and human health [7], these produced wastes must be well managed [8]. Unfortunately, these dumpsites are ineffective and poorly managed and pose greater risks to public health and environment especially to the residents of nearby settlements [9,10].

This link has been demonstrated in a number of studies (Aluko & Sridhar 2005) [11-15]. For example, direct handling of solid organic waste can result in various types of infectious and chronic diseases [16,17] while chemicals, broken glass, hypodermic needles and other hospital wastes risk injuring or poisoning waste workers or playing children [18]. Similarly, the waste can cause smells, dust, filth and dirt in residential areas [4] which will adversely affect the air people breath [19-21]; respiratory complications, irritation, and allergies [22] and the public and environmental health [23]. Furthermore, these waste dumpsites can also provide fertile ground for the proliferation of flies, mosquitoes, rodents and other disease vectors and pathogens.

Open dumping is the preferred method of solid waste disposal [24]. This situation creates suitable conditions for increased health risks to people living closer to dumpsites [25-27]. This scenario also applies to Kitui Town, which is the

County headquarters and largest town of Kitui County in Kenya. In Kitui, the Kitui Town Administration unit is responsible for managing all the generated solid waste in Kitui County. The collected waste is dumped at Kalundu dumpsite which derives the name from the adjacent Kalundu slum, a residential area that is surrounded by several businesses, a garage and a cattle traders market. At the edge of the dumpsite, there is the Kalundu River.

Because of its location and the fact that it is not protected or fenced, the dumpsite is a risk to the neighbouring community since it can adversely affect the immediate human health of those living in close proximity. It is therefore necessary to assess the health effects of the dumpsite location to the local community. It is on this background that this study assesses the health effects of solid waste disposal on Kalundu dumpsite in Kitui Town on the surrounding human settlements.

2. METHODOLOGY

This study was carried out in Kitui Town the headquarter of Kitui County of Kenya. The town lies between latitudes 0°10 and 3°0 South and longitudes 37°50 and 39°0 East. Kitui County is mostly dry and hot with temperatures ranging between 14°C in July-August and 34°C in January-March with rainfall ranging between 500mm and 1050mm annually. Kalundu open dumpsite is currently the only disposal site for all the solid waste generated and collected in the town. The dumpsite lies in Kitui Central Sub County, Kitui Township Ward in Kalundu village.

For this study, a survey was used to collect data from people living close to the dumpsite vicinity in a radius of 500 meters. In order to assess the perceptions of people living close to the dumpsite on the health risks, the respondents were stratified into two categories (i) those very close to the dumpsite (less than <250 meters), and those beyond 250 meters but within the 500 meters range. The number of respondents was determined using the formula by Kothari [28].

$$n = \frac{z^2 pqN}{e^2 (N - 1) + z^2 pq} \quad (1)$$

Where:

n: is the sample size for a finite population (smaller than 50,000)

N: size of the population = 800 households (Kalundu village)

p: population reliability (or frequency estimated for a sample of size n), where p is 0.5 which is taken for all developing countries population and p + q= 1

e: margin of error = 10%

Z- normal reduced variable at 5% level of significance (which is 1.96)

From the formula, 78 respondents were selected representing 39 from each stratum. A structured questionnaire was then administered to the respondents to collect data on their perceived health risks they associate with the presence of the dumpsite in their locality. The collected data was then coded and inputted into the Statistical Package for Social Sciences (SPSS) for analysis.




Fig. 1. Location of Kalundu Dumpsite in kalundu locality of Kitui Town, Kitui County – Kenya
Source; GIS Field Survey

Pearson Correlation Coefficient was used to test the links between respondent characteristics, household location from the dumpsite and the health perceptions. Participation was voluntary, and the respondent's privacy and anonymity was guaranteed.

3. RESULTS AND DISCUSSION

3.1 Demographic and Socio-economic Characteristics of the Respondents

Out of the 78 respondents, 50 were male while 28 were female. The respondents' ages ranged from 29 to 35 years. Majority of them were primary school leavers with those who had attained secondary school level representing 32% and only 10.3% had proceeded beyond secondary education. 56.4% of the respondents within 250 m from the dumpsite (near the dumpsite) were self-employed. This category was represented by 51.3% for those further than 250 m but within 500 m of the dumpsite (far from dumpsite). The average monthly earnings were below Ksh.10, 000, basically translating to below 3 dollars in a day. This low education and employment status of residents contributed to them resolving to make a living through solid waste scavenging. Here, they collected waste valuables for sell thereby exposing themselves to increased risk of health hazard exposure, cases that have been reported in other studies [29-31].

3.2 Perceived Health Risks Associated with Waste Dumping at Kalundu Dumpsite

83.3% of respondents were in agreement that Kalundu dumpsite posed a health hazard to the surrounding community and environment (Fig. 2).

Just like in other studies [32-34,15], the perceived health risks reported by the respondents in this study included; smoke emanating from burning waste, odour from rotting organic waste, providing an habitat for breeding mosquitoes, and harboring other disease vectors (Fig. 3). These together with high incidence of snake presence in the dumpsite vicinity exacerbate the health risks associated with living in close proximity to Kalundu dumpsite. Further, the respondents noted that the dumpsite together with the overlying human settlement areas, adjacent mechanic garages and livestock market acted as major sources of contamination to their only water source – the Kalundu River. Thus, they were exposed to health risks associated with various forms of water contamination as reported in other studies [35-37,7]. There was an indication of water contamination evidenced through waste spilled from dumpsite lying along the bank of Kalundu River (Fig. 4).




Fig. 2. Respondent view of the dumpsite being a health hazard to the people and surrounding environment




Fig. 3. Perceived health risks associated with presence of dumpsite in the Kalundu neighbourhood



Fig. 4. Waste leakage and spillage along the banks of Kalundu River

According to the respondents, Malaria was the most common disease associated with the presence of the dumpsite within their vicinity (Fig. 5). This was attributed to mosquito bites, an indication that the dumpsite provided a conducive habitat for mosquitoes to thrive. This was followed by chest pain; diarrhea and cholera respectively (Fig. 5). However, cholera was perceived to be a seasonal disease with high incidences during the rainy season. Skin infections, eye irritation and nose irritation were cited as minor occurrences. This finding corroborate other studies that demonstrate a direct link between health effects

and location of dumpsites, thus dumpsites pose significant health risks to the environment and people living close by Yongsu, [38-40]. Kalundu dumpsite is not fenced, waste dumping is indiscriminate and there is no control of who or what entered into it or came out of it. This creates a conducive environment that poses major health risks as reported by the respondents and corroborated by the findings of another study [41]. Furthermore, it has been observed and reported that solid wastes that are not well managed offer serious health risks to the surrounding human settlements [42].




Fig. 5. Perceived disease incidences associated with the presence of Kalundu dumpsite




Fig. 6. Relationship between distance from dumpsite and reported frequency of ill-health cases specified over time

Although both groups of respondents as per the study demarcation indicated that the dumpsite posed environmental and human health challenges to the community, this study found that those living within 250 m of the dumpsite (near the dumpsite) were more affected (Fig. 6; Table 1). It indicates that increasing the distance

between settlements and the dumpsite reduces the frequency of illnesses and vice versa (Fig. 6). This observation corroborates other studies linking living close to dumpsite and perception of environmental health risks [43,44,45,46]. The study also brought into attention the environmental implications of poor solid waste

Table 1. Respondents' perceptions on environmental health risks associated with solid waste dumped at Kalundu dumpsite

Environmental implications	Intensity	Near dumpsite	Far from dumpsite	Total
Land degradation	Very intense	52.5%	54.5%	53.5%
	Slightly intense	19.8%	17.8%	18.8%
	Intense	4.0%	10.9%	7.4%
	Not intense	23.8%	16.8%	20.3%
	X ²	4.60		
	P- value	0.203		
Air pollution	Very intense	62.4%	48.5%	55.4%
	Slightly intense	7.9%	25.7%	16.8%
	Intense	7.9%	12.9%	10.4%
	Not intense	21.8%	12.9%	17.3%
	X ²	14.78		
	P- value	0.00***		
Water pollution	Very intense	15.8%	13.9%	14.9%
	Slightly intense	61.4%	11.9%	36.6%
	Intense	8.9%	60.4%	34.7%
	Not intense	13.9%	13.8	13.8
	X ²	72.546		
	P- value	0.00***		
Unhygienic environment	Very intense	80.2%	7.9%	44.1%
	Slightly intense	13.9%	44.6%	29.2%
	Intense	3.0%	31.7%	17.3%
	Not intense	3.0%	15.8%	9.4%
	X ²	109.09		
	P- value	0.00***		

Note *** indicates significance at 1%




Fig. 7. Age group vulnerability to health risks associated with living in the dumpsite vicinity

management at Kalundu dumpsite on land degradation where 53.5% perceived the disposed waste to have a land degrading effect; 55.4% perceived it to be degrading the air while

44.1% perceived the waste as creating unhygienic conditions (Table 1). These effects were more severe in the near the dumpsite settlements compared with the far the dumpsite

ones. In contrast, water pollution was perceived to be very intense or intense by over 75% of respondents residing far from the dumpsite compared to only slightly over 34% of those living near the dumpsite (Table 1).

Further, the result indicates a statistically significant relationship between intensity of air pollution, water pollution and unhygienic environments ($p < 0.01$) whilst the association between intensity of land degradation and the study areas was not statistically significant ($p = 0.203$) (Table 1).

According to the respondents' views, children under the age of 5 years were the most vulnerable to contract disease compared to other age groups. Children from the group living near the dumpsite were more susceptible with 70% of them affected compared to 55% in the group living further away from the dumpsite (Fig. 7). Further, respondents highlighted the presence of dangerous items that risked injuring or poisoning waste workers and/or playing children, a finding that was also reported by Davis and Cornwell, 2008.

4. CONCLUSION

This study concludes that there is a link between waste deposited onto dumpsites and human and environmental health of the surrounding community with those living closer to the dumpsite more vulnerable.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Liu X, Huang Y, Xu X, Li X, Li X, Ciais P, Zeng Z. High-spatiotemporal-resolution mapping of global urban change from 1985 to 2015. *Nature Sustainability*. 2020;3(7):564-570.
- Habitat UN. State of the world's cities 2010/2011: bridging the urban divide. Earthscan, London; 2010.
- Ameen RFM, Mourshed M. Urban environmental challenges in developing countries—A stakeholder perspective. *Habitat International*. 2017;64:1-10.
- Rachel OA, Komine H, Yauhara K, Murakami S. Municipal Waste Management in Developed and Developing Countries: Japan and Nigeria as case Studies. *Solid Waste Audit Report, Federal Capital Territory Abuja Nigeria*, 2004; 2009.
- Karak T, Bhagat RM, Bhattacharyya P. Municipal solid waste generation, composition, and management: the world scenario. *Critical Reviews in Environmental Science and Technology*. 2012;42(15):1509-1630.
- Smit W. Urban governance in Africa: an overview. *African Cities and the Development Conundrum*. 2018;55-77.
- Aluko OO, Obafemi TH, Obiajunwa PO, Obiajunwa CJ, Obisanya OA, Odanye OH, Odeleye AO. Solid waste management and health hazards associated with residence around open dumpsites in heterogeneous urban settlements in Southwest Nigeria. *International Journal of Environmental Health Research*. 2021;1-16.
- Ferronato N, Torretta V. Waste mismanagement in developing countries: A review of global issues. *International Journal of Environmental Research and Public Health*. 2019;16(6):1060.
- Wilson DC, Araba AO, Chinwah K, Cheeseman CR. Building recycling rates through the informal sector. *Waste Management*. 2009;29(2):629-635.
- McGill R. Making towns work: Habitat III—what relevance?. *Planning Theory & Practice*. 2018; 19(1):140-148.
- Nwanta JA, Onunkwo J, Ezenduka E. Analysis of Nsukka metropolitan abattoir solid waste and its bacterial contents in south eastern Nigeria: Public health implication. *Archives of Environmental & Occupational Health*. 2010;65(1):21-26.
- Norsa'adah B, Salinah O, Naing NN, Sarimah A. Community health survey of residents living near a solid waste open dumpsite in Sabak, Kelantan, Malaysia. *International Journal of Environmental Research and Public Health*. 2020;17(1):311.
- Olu J, Iyere A. Perception of residents and workers toward the environmental health effects of the dumpsite in Solous Igando, Lagos, Nigeria. *Journal of Advances in Environmental Health Research*. 2020; 8(4):242-249.
- Amugsi DA, Haregu TN, Mberu BU. Levels and determinants of perceived health risk from solid wastes among communities living near to dumpsites in Kenya.

- International Journal of Environmental Health Research. 2020;30(4):409-420.
15. Etea T, Girma E, Mamo K. Risk perceptions and experiences of residents living nearby municipal solid waste open dumpsite in Ginchi Town, Ethiopia: A Qualitative Study. *Risk Management and Healthcare Policy*. 2021;14:2035.
 16. Vimercati L, Baldassarre A, Gatti MF, De Maria L, Caputi A, Dirodi AA, Bellino RM. Respiratory health in waste collection and disposal workers. *International Journal of Environmental Research and Public Health*. 2016;13(7):631.
 17. Heldal KK, Halstensen AS, Thorn J, Eduard W, Halstensen TS. Airway inflammation in waste handlers exposed to bioaerosols assessed by induced sputum. *European Respiratory Journal*. 2003; 21(4):641-645.
 18. Davis ML, Cornwell DA. *Introduction to environmental engineering*. McGraw-Hill; 2008.
 19. De Feo G, De Gisi, S, Williams ID. Public perception of odour and environmental pollution attributed to MSW treatment and disposal facilities: A case study. *Waste Management*. 2013; 33(4):974-987.
 20. Sakawi Z, Sharifah SA, Jaafar O, Mahmud M. Community perception of odor pollution from the landfill. *Research Journal of Environmental and Earth Sciences*. 2011;3(2):142-145.
 21. Kumar S, Gaikwad SA, Shekdar AV, Kshirsagar PS, Singh RN. Estimation method for national methane emission from solid waste landfills. *Atmos. Environ*. 2004;38:3481–3487. DOI: 10.1016/j.atmosenv.2004.02.057
 22. Vinti G, Bauza V, Clasen T, Medlicott K, Tudor T, Zurbrugg C, Vaccari M. Municipal Solid Waste Management and Adverse Health Outcomes: A Systematic Review. *International journal of Environmental Research and Public Health*. 2021;18(8):4331.
 23. Boningari T, Smirniotis PG. Impact of nitrogen oxides on the environment and human health: Mn-based materials for the NO_x abatement. *Current Opinion in Chemical Engineering*. 2016; 13:133-141.
 24. Muiruri J, Wahome R, Karatu K. Assessment of methods practiced in the disposal of solid waste in Eastleigh Nairobi County, Kenya. *AIMS Environmental Science*. 2020;7(5):434-448.
 25. Sankoh FP, Yan X, Tran Q. Environmental and health impact of solid waste disposal in developing cities: a case study of granville brook dumpsite, Freetown, Sierra Leone. *Journal of Environmental Protection*; 2013.
 26. Brender JD, Maantay JA, Chakraborty J. Residential proximity to environmental hazards and adverse health outcomes. *American Journal of Public Health*. 2011;101(S1):S37-S52.
 27. Vrijheid M. Health effects of residence near hazardous waste landfill sites: a review of epidemiologic literature. *Environmental Health Perspectives*. 2000;108(suppl 1):101-112.
 28. Kothari CR. *Research methodology: Methods and techniques*. New Age International; 2004.
 29. Bradney L, Wijesekara H, Palansooriya KN, Obadamudalige N, Bolan NS, Ok YS, Kirkham MB. Particulate plastics as a vector for toxic trace-element uptake by aquatic and terrestrial organisms and human health risk. *Environment International*. 2019;131:104937.
 30. Naidoo T, Rajkaran A. Impacts of plastic debris on biota and implications for human health: A South African perspective. *South African Journal of Science*. 2020;116(5-6):1-8.
 31. Silva ALP, Prata JC, Duarte AC, Barcelò D, Rocha-Santos T. An urgent call to think globally and act locally on landfill disposable plastics under and after covid-19 pandemic: Pollution prevention and technological (Bio) remediation solutions. *Chemical Engineering Journal*, 2021;131201.
 32. Milosevic LT, Mihajlovic ER, Djordjevic AV, Protic MZ, Ristic DP. Identification of fire hazards due to landfill gas generation and emission. *Polish Journal of Environmental Studies*. 2018; 27(1).
 33. Nair AT, Senthilnathan J, Nagendra SS. Emerging perspectives on VOC emissions from landfill sites: Impact on tropospheric chemistry and local air quality. *Process Safety and Environmental Protection*. 2019;121:143-154.
 34. Vaverková MD. Landfill impacts on the environment. *Geosciences*. 2019;9(10):431.
 35. Sibanda LK, Obange N, Awuor FO. Challenges of solid waste management in Kisumu, Kenya. In *Urban Forum*. Springer

- Netherlands. 2017, December;28(4):387-402.
36. Waturu G, Mutai W, Kiiru J, Musyoki M, Ochieng L. Antimicrobial resistance encountered in garbage collection areas and dumpsites in Nairobi, Kenya using *Escherichia coli* and *Klebsiella* as indicator species. *Advances in Microbiology*. 2017; 7(9):653-665.
 37. Awino FB. Assessment of Metal Contaminants and Human Health Risks in Food Crops at the Uncontrolled Mbale Dumpsite, Uganda (Doctoral dissertation, University of Canberra); 2020.
 38. Yongsi HB. Pathogenic microorganisms associated with childhood diarrhea in Low-and-middle income countries: Case study of Yaoundé–Cameroon. *International Journal of Environmental Research and Public Health*. 2008;5(4):213-229.
 39. Giusti L. A review of waste management practices and their impact on human health. *Waste Management*. 2009;29(8): 2227-2239.
 40. Gouveia N, Prado RRD. Health risks in areas close to urban solid waste landfill sites. *Revista de Saúde Pública*. 2010; 44(5):859-866.
 41. Sood D. Solid waste management study for freetown (Component Design for World Bank, Draft Report Project No. P078389). Great Falls, Virginia; 2004.
 42. Nwogwugwu N, Ishola AI. Solid waste management and public health challenges: Appraisal of local government capacity to achieve effective environmental governance. *Asian Social Science*. 2019; 15(5):1-9.
 43. Okot-Okumu J. Solid Waste Management in Uganda: issues challenges and opportunities. In POVIDE programme Workshop. The Netherlands; 2008.
 44. Gakungu NK, Gitau AN, Njoroge BNK, Kimani MW. Solid waste management in Kenya: A case study of public technical training institutions. *ICASTOR Journal of Engineering*. 2012;5(3):127-138.
 45. Abul S. Environmental and health impact of solid waste disposal at Mangwaneni dumpsite in Manzini: Swaziland. *Journal of Sustainable Development in Africa*. 2010;12(7):64-78.
 46. UN-Habitat. *The State of African Cities*;2010. Available:<http://www.unhabitat.org/documents/SOAC10?SOAC-PRI-en.pdf>; retrieved on August, 31, 2019.

© 2022 Matheka et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/89210>