



Intercropping of Pulses and Oilseeds with Sisal (*Agave sisalana*) in North Coastal Zone of Andhra Pradesh

Malleswari Sathineni ^{a*}, T. Sreelatha ^a and S. Mitra ^b

^a AINP on Jute and Allied Fibres, Agricultural Research Station, Amadalavalasa, Acharya NG Ranga Agricultural University, Andhra Pradesh-532185, India.

^b ICAR-Central Research Institute for Jute and Allied Fibres, Barrackpore, West Bengal-700120, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2022/v12i1211551

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/95158>

Original Research Article

Received: 18/10/2022

Accepted: 25/12/2022

Published: 28/12/2022

ABSTRACT

A field experiment was conducted during *Kharif, Rabi, 2020 and 2021* under AINP on Jute and Allied Fibres at Agricultural Research Station farm, Amadalavalasa – ANGRAU, Andhra Pradesh. The experiment was conducted to study the suitability and economic feasibility of short duration pulses and oilseeds as intercrops in between paired row spaces of sisal plantation. Results revealed that, short duration pulses are more feasible than oilseeds as intercrops with increased productivity per unit area. Sisal fibre yield and intercrops yield was higher with pulse intercropping than oilseeds. Sisal fibre equivalent yield of intercrops varied between 1.87 q ha⁻¹ in case of sisal + mustard to 5.28 q ha⁻¹ in sisal + cowpea. The net returns per rupee invested was highest in sisal + cowpea (3.42) followed by sisal + greengram (3.34), sisal + rajma (3.05) and lowest among pulses was sisal + blackgram (2.83). While, among oilseeds, highest was in sisal + niger (2.90) followed by sisal + mustard (2.80) and sisal + sesame (2.65) compared to lowest in sole sisal (2.49). However, the total system productivity in terms of sisal fibre equivalent yield of cropping system varied between 14.05 q ha⁻¹ in case of sole sisal to 21.64 q ha⁻¹ in sisal + cowpea. The results inferred

*Corresponding author: E-mail: sn.malleswari@angrau.ac.in;

that, growing of short duration pulses in between paired row spaces of sisal could be efficiently used by growing short duration pulses like cowpea, greengram, rajma and blackgram for increased cropping system under sisal plantation.

Keywords: *Sisal based intercropping; short duration pulses; oilseeds; economics; yield.*

1. INTRODUCTION

Sisal (*Agave sisalana*), is a leaf fibre producing plant belongs to Asparagaceae family, commonly known as Agave in India. It is semi perennial, xerophytic and mainly grown in arid and semi-arid regions of Andhra Pradesh, Orissa, Bihar, Karnataka, Maharashtra, and West Bengal [1]. In the context of climate change and land degradation neutrality, sisal is an excellent CDM (clean development mechanism) crop for biochemical as well as for afforestation in degraded lands besides improvement in soil carbon sequestration [2]. Generally, sisal plant resistant to many pest and diseases consequently reduces pesticide load in the environment, reduces soil erosion through its extensive root system, contributes positively to watershed management. Therefore, designated as "specialty crop" for conservation agriculture [2-4].

Several species of sisal are indigenous to India largely grown as live fence as well as in wild conditions. Different species are *Agave sisalana*, *A. cantala*, *A. veracruz*, *A. amanuensis*, *A. angustifolia*, and *A. fourcroyodes*. Among these, *A. sisalana* contributes nearly 85% of total sisal fibre production in the world. Healthy sisal plant produces 500 -750 leaves in about 10-12 years of its life span. Thus a normal sisal leaf with 600-gram weight produces about 1000 fibres for each leaf [1,4]. Globally, sisal fibre can be used for multiple purposes and commonly used in shipping industry and several other domestic to industrial uses including high strength requiring long lasting geo-textile and specialty composites.

In this scenario, renewed interests in recent times has been noticed to grow sisal primarily for its quality fibre having great demand in the country [4]. Further, some pockets of north coastal region of Andhra Pradesh are having undulated terrain, poor soil fertility, low water retention capacity, where no other arable crop could be grown by the resource poor farmers. Sisal appears to be a promising alternative in this

region, if profitability of this crop could be sustained by introduction of suitable inter crops in sisal for increasing the system productivity.

However, improved production technology of paired row system is usually practiced in north coastal zone for sisal production. But in this system a large amount of land area unutilized in between rows. Therefore, the inter-row space may be utilized with suitable intercrops to improve profitability of the system. But, sufficient and detailed research work on suitable inter crops for sisal plantation in north coastal zone of A.P is truly lacking. Therefore, a field experiment was conducted to identify suitable intercrops for increasing the system productivity as well as profitability of sisal-based intercropping system.

2. MATERIALS AND METHODS

The field experiment was conducted during *kharif* and *Rabi*, 2020 -21 and 2021-22 in sandy clay loam soil at Agricultural Research Station farm, Amadalavalasa, ANGRAU, Andhra Pradesh under All India Network Project on Jute and Allied Fibres (AINPJAF). Geographically, the experimental field is situated at latitude / longitude of 18.4°N, 83.89°E with an altitude of 35 m above mean sea level.

The experiment was laid out in randomized block design with 9 treatments and three replications. The treatment details are T₁: Sisal + Black gram (GBG-1); T₂: Sisal + Greengram (IPM 2-14); T₃: Sisal + Cowpea (DC-15); T₄: Sisal + Horsegram (CRHG -22); T₅: Sisal + Sesame (YLM-66); T₆: Sisal + Niger (KGN-2); T₇: Sisal + Mustard (Varuna); T₈: Sisal + Rajma (Amber) and T₉: Sole Sisal. The agro-techniques adopted for the short duration pulses and oilseeds as intercrops in sisal plantation is given in Table 1.

The sole sisal crop was sown during *kharif* season and in the *Rabi* season sisal along with short duration pulses and oilseeds crops were grown. Sisal crop was sown in 4.0 m × 4.0 m paired row and inter crops were sown in line with

Table 1. Agro-techniques adopted for pulses and oilseeds as intercrops in between paired rows of sisal

| Crop | Variety | Duration (Days) | Spacing (cm) | RDF N:P:K:S (kg ha ⁻¹) |
|--|----------|-----------------|--------------|------------------------------------|
| Pulses | | | | |
| <i>Vigna mungo</i> (L.) (Blackgram) | GBG-1 | 70-75 | 30 × 10 | 20:50:0 |
| <i>Vigna radiata</i> (L.) (Greengram) | IPM 2-14 | 60-65 | 30 × 10 | 20:50:0 |
| <i>Vigna unguiculata</i> (L.) (Cowpea) | DC-15 | 75-80 | 30 × 15 | 15:50:50 |
| <i>Macrotyloma uniflorum</i> (L.) (Horsegram) | CRHG -22 | 85-90 | 30 × 10 | 10:25:4 |
| <i>Phaseolus vulgaris</i> (Rajmash) | Amber | 70-75 | 30 × 10 | 90:60:0 |
| Oilseeds | | | | |
| <i>Sesamum indicum</i> (L.) (Sesame) | YLM-66 | 80-90 | 30 × 15 | 50:20:20 |
| <i>Guizotia abyssinica</i> (L.) (Niger) | KGN-2 | 95-100 | 30 × 10 | 10:0:0 |
| <i>Brassica nigra</i> (Mustard) | Varuna | 135-140 | 30 × 15 | 60:30:15:20 |

4.0 m × 2.4 m spacing in between the lines of sisal. The main crop sisal and inter crops were grown completely under rainfed conditions. Agronomic management for growing of inter crops were followed as per the package of practices as prescribed by ANGRAU. Weeding and thinning operations in intercrops was done after 30 days after sowing and harvested at maturity. Biometrical observations of all the component crops were recorded at regular intervals. The yield and yield attributes were recorded at harvest. The recorded observations for different parameters were statistically analysed using analysis of variance (ANOVA) technique and results are presented at 5 % level of significance (P=0.05).

3. RESULTS AND DISCUSSION

3.1 Effect of Intercrops on Growth Parameters of Sisal

Effect of pulses and oilseeds as intercrops on growth parameters of sisal did not vary significantly during the two years of study (Table 2). The pooled data of the two-year study revealed that, highest number of leaves per sisal plant was recorded in pulses as intercrops than oilseeds. Highest number of leaves (31.9) recorded in sisal + horsegram followed by sisal + cowpea (31.8), sisal + rajma (31.7), sisal + greengram (31.3) and lowest was in sisal + blackgram (30.1). While oilseeds as intercrops, highest number of sisal leaves was recorded in

sisal + sesame and sisal + niger (29.7) and lowest was in (29.4) as compared to sole sisal (29.2). Whereas, highest leaf length was recorded in Sisal + Greengram (101.0 cm) followed by (sisal + horsegram) (97.9 cm), which was on par with sole sisal. Among oilseeds as intercrops highest leaf length was recorded in sisal + niger (99.5 cm) followed by sisal + sesame (98.7 cm) and lowest was recorded in sisal + mustard (98.4 cm) as compared to sole sisal. Similar findings were also reported by Sarkar et al. [4] in intercropping system with pulses followed in double rowed sisal plantation.

However, similar trend was recorded with respect to leaf width of sisal wherein, highest was in sisal + cowpea (9.6 cm) among pulses, sisal + sesame and sisal + niger (9.6 cm) among oilseeds than sole sisal (9.5 cm). While the green biomass of sisal was obtained with sisal + cowpea (47.25 t/ha) followed by sisal + greengram (47.18 t/ha) in both pulses and oilseeds as intercrops compared to sole sisal (45.97 t/ha). The results are in conformity with Sarkar et al. [4]. All intercrops in the present study successfully established in sisal interspaces as pulses and oilseeds fit well in intercropping system. However, pulses influenced yield attributes of sisal than oilseeds might be due to plant stature of the pulses and provided enough space for the sisal to expand. Besides, intercrops positively influenced the soil fertility.

Table 2. Effect of pulses and oilseeds grown as intercrops on growth parameters of Sisal

| Treatments | No. of Leaves / Plant | | | Leaf length (cm) | | | Leaf width (cm) | | | Green biomass (t/ha) | | |
|-------------------|-----------------------|------|--------|------------------|-------|--------|-----------------|------|--------|----------------------|-------|--------|
| | 2020 | 2021 | pooled | 2020 | 2021 | pooled | 2020 | 2021 | pooled | 2020 | 2021 | pooled |
| Sisal + Blackgram | 25.4 | 34.9 | 30.1 | 103.5 | 90.7 | 97.1 | 10.6 | 8.3 | 9.4 | 43.34 | 41.39 | 42.37 |
| Sisal+ Greengram | 26.5 | 36.2 | 31.3 | 102.5 | 101.0 | 101.7 | 10.5 | 8.5 | 9.5 | 45.12 | 49.24 | 47.18 |
| Sisal + Cowpea | 27.2 | 36.5 | 31.8 | 100.7 | 95.0 | 97.8 | 10.6 | 8.5 | 9.6 | 46.38 | 48.13 | 47.25 |
| Sisal + Horsegram | 26.9 | 36.9 | 31.9 | 103.4 | 92.4 | 97.9 | 10.7 | 8.2 | 9.5 | 45.89 | 45.29 | 45.59 |
| Sisal + Sesame | 26.5 | 32.9 | 29.7 | 102.6 | 94.8 | 98.7 | 10.8 | 8.3 | 9.6 | 45.22 | 38.39 | 41.80 |
| Sisal + Niger | 27.2 | 32.1 | 29.7 | 101.2 | 97.8 | 99.5 | 10.7 | 8.4 | 9.6 | 46.46 | 43.26 | 44.86 |
| Sisal + Mustard | 26.6 | 32.3 | 29.4 | 102.3 | 94.5 | 98.4 | 10.6 | 8.3 | 9.4 | 45.40 | 37.33 | 41.36 |
| Sisal + Rajma | 26.4 | 37.0 | 31.7 | 102.5 | 90.6 | 96.5 | 10.7 | 7.8 | 9.3 | 45.10 | 46.75 | 45.93 |
| Sole Sisal | 25.5 | 33.0 | 29.2 | 102.9 | 92.8 | 97.9 | 10.8 | 8.1 | 9.5 | 43.49 | 48.46 | 45.97 |
| CD (P=0.05) | NS | NS | NS | NS | NS | NS | NS | NS | NS | NS | NS | NS |
| SEm ± | 0.97 | 2.71 | 1.44 | 0.88 | 3.98 | 2.04 | 0.21 | 0.29 | 0.24 | 1.66 | 3.53 | 1.95 |
| CV (%) | 6.4 | 13.5 | 11.5 | 1.5 | 7.3 | 5.1 | 3.3 | 5.9 | 4.5 | 6.4 | 13.8 | 10.7 |

3.2 Effect of Intercrops on Yield Parameters of Sisal

The fibre yield of sisal during two years of study did not vary significantly (Table 3). However, pooled data of fibre yield revealed that, effect of intercrops on yield of sisal was higher with pulses than oilseeds. Among pulses, except sisal + black gram which recorded lower yield (14.69 q/ha) and remaining treatments were on par with highest yield in sisal + cowpea (16.35 q/ha). Among, oilseeds highest yield was recorded in sisal + niger (15.57 q/ha) followed by sisal + sesame (14.54 q/ha) and sisal + mustard (14.40 q/ha). However, sole sisal was comparatively higher than oilseeds and recorded 15.88 q/ha of sisal fibre yield. The similar findings were also reported by Rajesh kumar et al. [5] in maize intercropped with pulses.

Pooled data of intercrops yield was revealed that, yields among intercrops was on par and highest yield was recorded in sisal + cowpea (11.47 q/ha) and lowest was in sisal + rajma (3.73 q/ha). However, among oilseeds as intercrops highest yield was recorded in sisal + mustard (3.07 q/ha) followed by sisal + sesame (2.47 q/ha) and sisal + niger (2.43 q/ha). Similar trend of results was obtained when intercrops yield was converted to sisal fibre equivalent yield. The results are in conformity with the findings of Sarkar et al. [6] who reported similar trend of results in pigeonpea and sisal intercropping system.

In the present study, fibre yield of sisal, yield of short duration pulses and oilseeds was expressed as sisal fibre equivalent yield of cropping system (q/ha). Pooled results of cropping system as a whole reveal that, effect of short duration pulses as intercrops was significantly higher as compared to oilseeds. The results are in line with the findings of Sujatha et al. [7] and Behera et al. [8].

Similarly, both short duration pulses and oilseeds as intercrops significantly influenced the fibre yield of sisal. Highest sisal fibre equivalent yield of cropping system was obtained in sisal + cowpea (21.64 q/ha) followed by sisal + greengram (21.14 q/ha), sisal + rajma (19.81 q/ha), sisal + blackgram (18.27 q/ha) and lowest among pulses was with sisal + horsegram (17.29 q/ha). Among oilseeds as intercrops highest was obtained in sisal + mustard (17.81 q/ha) followed

by sisal + niger (17.64 q/ha) and lowest was in sisal + sesame (16.72 q/ha). However, sole sisal crop recorded lowest equivalent yield (14.05 q/ha) as compared to both short duration pulses and oilseeds as intercrops in the cropping system. The results are in line with the findings of Sarkar et al. [4] who reported that growing of legume intercrops like pigeonpea, cowpea and blackgram with sisal improved the system productivity. In the present study, pulse crops impacted the Sisal fibre equivalent yield of cropping system (q/ha) than oilseeds. The pulse crops are having positive impact on symbiosis for nitrogen fixation and increase in soil fertility. They have high soil ameliorating properties and ability to thrive better under unfavourable conditions makes the pulses an integral component of agriculture perse as intercrops. This has increased the fibre yield and yield attributes of sisal and increased the system productivity.

3.3 Economics of Sisal Based Intercropping System

The economics of sole sisal crop and intercropping with short duration pulses and oilseeds was worked out by considering the variable cost of fertilizers, labours, seeds and price of each crop yield. Calculated the gross returns, net returns and benefit cost ratio of the sole sisal, intercropping with pulses and oilseeds and sisal based cropping system (Table 4). The results revealed that, inter cropping with short duration pulses positively contributed to economic feasibility and sustainability of the system by ensuring higher economic income to farmers in sisal based cropping system.

Sisal intercropped with cowpea treatment among different treatments ensures highest profit to resource poor farmers as it was reflected in gross, net profit and highest B:C ratio (3.42) compared to other treatments and lowest B:C ratio was recorded in sole sisal (2.49). Intercropping with sisal + cowpea ensured highest B:C ratio followed by sisal + greengram (3.34), sisal + rajma (3.05), sisal + niger (2.90), sisal + horsegram (2.90), sisal + blackgram (2.83), sisal + mustard (2.80) and sisal + sesame (2.65). In the present study, intercropping with short duration pulses contributed higher than oilseeds for cropping system productivity. The results are in conformity with the findings of Padhi et al. [9] and Sarkar et al. [4].

Table 3. Yield and fiber equivalent of Sisal in the intercropping system

| Treatments | Sisal Fibre yield (q/ha) | | | Intercrops yield (q/ha) | | | Sisal fibre equivalent yield of intercropping system (q/ha) | | | Sisal fibre equivalent yield of cropping system (q/ha) | | |
|-------------------|--------------------------|-------|--------|-------------------------|------|--------|---|------|--------|--|-------|--------|
| | 2020 | 2021 | pooled | 2020 | 2021 | pooled | 2020 | 2021 | pooled | 2020 | 2021 | pooled |
| Sisal + Blackgram | 15.79 | 13.60 | 14.69 | 4.55 | 4.74 | 4.64 | 3.41 | 3.73 | 3.57 | 19.20 | 17.33 | 18.27 |
| Sisal+ Greengram | 16.43 | 16.18 | 16.31 | 5.07 | 5.61 | 5.34 | 4.56 | 5.10 | 4.83 | 20.99 | 21.28 | 21.14 |
| Sisal + Cowpea | 16.89 | 15.82 | 16.35 | 14.50 | 8.45 | 11.47 | 6.34 | 4.22 | 5.28 | 23.24 | 20.04 | 21.64 |
| Sisal + Horsegram | 16.71 | 14.88 | 15.80 | 3.98 | 3.96 | 3.97 | 1.49 | 1.48 | 1.49 | 18.21 | 16.36 | 17.29 |
| Sisal + Sesame | 16.47 | 12.62 | 14.54 | 2.64 | 2.30 | 2.47 | 2.27 | 2.10 | 2.18 | 18.73 | 14.72 | 16.72 |
| Sisal + Niger | 16.92 | 14.21 | 15.57 | 2.34 | 2.53 | 2.43 | 1.96 | 2.19 | 2.07 | 18.88 | 16.41 | 17.64 |
| Sisal + Mustard | 16.53 | 15.36 | 14.40 | 2.95 | 3.19 | 3.07 | 1.71 | 2.02 | 1.87 | 18.25 | 17.38 | 17.81 |
| Sisal + Rajma | 16.43 | 15.92 | 15.89 | 4.58 | 2.88 | 3.73 | 4.58 | 2.70 | 3.64 | 21.00 | 18.63 | 19.81 |
| Sole Sisal | 15.84 | 12.27 | 15.88 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 0.00 | 15.84 | 12.27 | 14.05 |
| CD (P=0.05) | NS | NS | NS | - | - | - | - | - | - | 1.69 | 3.41 | 1.83 |
| SEm± | 0.61 | 1.16 | 0.65 | - | - | - | - | - | - | 0.56 | 1.14 | 0.64 |
| CV (%) | 6.4 | 13.8 | 10.3 | - | - | - | - | - | - | 5.1 | 11.5 | 8.5 |

Table 4. Effect of pulses and oilseeds grown as intercrop on economics of sisal (pooled data of 2020 and 2021)

| Treatment | Gross Returns of cropping system (Rs./ha) | Cost of Cultivation of cropping system (Rs./ha) | Net Returns of cropping system (Rs./ha) | B:C Ratio |
|-------------------|---|---|---|-----------|
| Sisal + Blackgram | 146160 | 51600 | 94560 | 2.83 |
| Sisal+ Greengram | 169120 | 50640 | 118480 | 3.34 |
| Sisal + Cowpea | 173120 | 50580 | 122540 | 3.42 |
| Sisal + Horsegram | 138320 | 47634 | 90686 | 2.90 |
| Sisal + Sesame | 133760 | 50436 | 83324 | 2.65 |
| Sisal + Niger | 141120 | 48740 | 92380 | 2.90 |
| Sisal + Mustard | 139760 | 49980 | 89780 | 2.80 |
| Sisal + Rajma | 156160 | 51130 | 105030 | 3.05 |
| Sole Sisal | 112400 | 45184 | 67216 | 2.49 |

Even though sole crop of sisal was grown as live fence and also as an erosion control measure in degraded soils. However, in the present study, inclusion of short duration pulses as intercrops in the system enhanced the productivity of the cropping system. This is due to positive impact of pulses and oilseeds on sisal fibre yield and in improving the system productivity. Similar findings were reported by Sarkar et al. 2015 wherein, they reported that, sisal equivalent yield from the system was highest with pigeonpea followed by cowpea and blackgram as intercrops along with sisal. In addition, the cropping system including pulses ensures food and nutritional security of farm family [10].

4. CONCLUSION

Sole sisal is generally grown in poor fertile soils, sloppy lands or in degraded soils as live fence, as erosion control measures and for its fibre in various diversified uses. The vacant space in between the paired rows of sisal plantation may be effectively utilized by growing short duration pulses, as they are relatively hardy, being adopted and fitted into various cropping systems compared to oilseeds. Thus, cultivation of short duration pulses in paired row sisal plantation effectively utilizes space available, protects soil and makes cropping system more remunerative by ensuring nutritional support and higher farm income to resource poor farm family than traditional sole sisal.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Nayak L, Nag D, Das S, Ray DP, Ammayappan L. Utilisation of Sisal Fibre (*Agave sisalana* L.) – A review. *Agricultural Reviews*. 2011;32(2):150-152.
2. Sarkar S, Jha AK. Research for sisal (*Agave* sp.) fibre production in India. *International Journal of Current Research*. 2017;9(11): 61136-61146.
3. Sarkar S, Saha AR, Majumdar B, Abdullah SK. Sisal: Speciality crop of conservation agriculture for peninsular India. In: Book of papers. XIX National Symposium on Resource Management Approaches Towards Livelihood Security. 2-4 December, 2010, Bengaluru; 2010.
4. Sarkar S, Kundu DK, Saha AR, Majumdar B, Jha AK. Intercropping system in double rowed sisal (*Agave sisalana*) plantation. *Indian Journal of Natural Fibres*. 2015;1(2):221-224.
5. Rajeshkumar A, Venkataraman NS, Ramadass S, Ashokkumar N, Thirumeninathan S. Effect of weed control and cropping system on weed population and productivity of maize grown sole or intercropped with pulses. *Journal of Crop and Weed*. 2017;13(3):150 -155.
6. Sarkar S, Kundu DK, Saha AR, Majumdar B. Feasibility of growing annual intercrops in sisal plantation in plateau region of India, In: Book of Abstracts National Seminar on Jute and Allied fibres in changing times: Issues and Strategies held at NIRJAFT, Kolkata, during 3-5, January. 2013;45.
7. Sujatha S, Bhat R, Kannan C, Balasimha D. Impact of intercropping of medicinal and aromatic plants with organic farming approach on resource use efficiency in arecanut (*Areca catechu* L.) plantation in India. *Industrial Crops and Products*. 2011;33(1):78-83.
8. Behera MS, Kundu DK, Satpathy S, Jha AK, Singh A, Nayak RK. Scope and opportunity of intercropping medicinal and aromatic plants with sisal cultivation. *International Journal of Tropical Agriculture*. 2015;33(2):1771-1773.
9. Padhi AK, Panigrahi RK, Jena BK. Effect of planting geometry and duration of intercrops on performance of pigeonpea and finger millet intercropping system. *Indian Journal of Agricultural Research*. 2010;44(1):43 -47.
10. FAO. Pulses: nutritious seeds for a sustainable future. Rome; 2016. Available: <https://doi.org/10.4060/i5528e>