



Impact of Crude Oil Contamination on Soil Physicochemical Properties around the Sildubi Oil Spill, Borhola, Assam

Tarun Saikia ^{a*}, Jhorna Bora ^b and Cinu Gogoi ^c

^a Department of Chemistry, D.C.B. Girls' College, Jorhat, Assam, India.

^b Department of Chemistry, Dibrugarh University, Dibrugarh, Assam, India.

^c CSIR – NEIST, Jorhat, Assam, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2023/v35i224130

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/108464>

Original Research Article

Received: 08/09/2023

Accepted: 11/11/2023

Published: 20/11/2023

ABSTRACT

Crude oil contamination can affect the soil physicochemical properties and it is a serious constraint on the rice ecosystem near the oil drilling sites. The objective of this study was to identify the effects of crude oil contamination through oil exploration on soil health and the horizontal distribution of soil nutrients in rice field soils in the Sildubi area, Jorhat district, Assam. Soil samples were collected at a depth of 0–15 cm from the drilling site and from the four directions (East, West, South and North) in a circular mode with a radius of 500 and 1000m away from the oil spill. Each sample was mounded with four replications. Samples were analyzed for physicochemical properties by using standard methods that reflect soil nutrient status.

The percentage of organic carbon, organic matter and soil pH are found to be higher at the drilling site compared to the results obtained in the rice field soils, apart by 500 and 1000m from the central zone. Similarly, bulk density, moisture percentage and electrical conductivity were found to

*Corresponding author: E-mail: saikia.tarun55@gmail.com;

be lowest in the soils of the drilling site. In our study, considerable variation in the horizontal distribution of soil nutrients was also observed. The highest P, Ca and nitrogen contents were recorded in the oil-contaminated soils compared to soils collected from 500 and 1000m radius. The total soil Fe was significantly higher in the samples collected at 1000m. Spilled crude oil is certainly responsible for alterations in soil physicochemical properties and the horizontal distribution of some plant-specific nutrients. High concentrations of Fe and P observed due to crude oil contamination have been developing Fe stress situations reducing crop production in nearby agricultural lands.

Keywords: Crude oil; physicochemical; bulk density; horizontal; Fe stress.

ABBREVIATIONS

TOC : Total organic Carbon,
TOM : Total organic Matters,
THC : Total petroleum Hydrocarbon,
EC : Electrical Conductivity,
DL Site : Drilling Site

1. INTRODUCTION

Soil is one of the most important resources in nature, an important reservoir of nutrients and moisture for the plant's production. Healthy soil is indeed alive and dynamic, consisting of microorganisms. The top-most layer of soil is comparatively richer in nutrients and supports maximum crop production. Normally, soil physicochemical properties vary distinctly from place to place, particularly with respect to depth in the soil profile, climate, water content of the soil, and soil organic carbons.

Crude oil contamination is one of the most severe problems for soil health. Oil Contamination reduces the strength, permeability, maximum dryness and optimum moisture content [1]. Crude oil contamination of the soil environment can undoubtedly limit its protective function, disturb metabolic activity, adversely affect its physicochemical characteristics, reduce fertility and negatively influence crop production. The oil deposited on the leaves of the plants penetrates into the leaves and reduces transpiration and photosynthesis rate [2]. For upland oil spills, the effects become more serious in nearby low-land crop fields. Crude oil pollution is a serious constraint on the rice ecosystem near the oil drilling sites.

Oil leakage is inadvertent due to oil exploitation, transportation pipeline damage, coastal facility discharge and oil tank accident [3]. Such unintended crude oil contamination comprises contaminants like heavy metals [4] and aromatic hydrocarbons. For crude oil contamination, pore

spaces might be blocked, which could reduce soil aeration and water infiltration and increase bulk density. Such a dramatic modification in crude oil-contaminated soil directly affects the growth and productivity of nearby rice fields. Moreover, oils that are denser than water might reduce and restrict soil permeability [5], Crude oil-contaminated soils are hydrophobic compared to immaculate sites [6]. Hydrocarbon contamination can increase soil total organic carbon [7] and change soil chemical properties like pH, total organic matter and soil mineral nutrients such as sodium, potassium, sulphate, phosphate and nitrate, thus indirectly affecting the growth and development of plants and microorganisms [8,9]. Crude oil contamination may increase the soil pH up to 8.0 [10] and in alkaline soils crude oil adversely affects soil physicochemical properties, particularly soil available phosphorous concentration and causes deterioration of wetland soil health.

The effects of crude oil contamination on soil health may be explained by two probable mechanisms. It may penetrate the soil interlayer, where it directly affects the plant root system, microbial population and thus depletes soil oxygen content. Crude oil may also develop a pseudo nutrient deficiency through a reduction in the level of available plant nutrients or by elevating the concentration of certain elements to toxic levels, such as iron. Oil usually develops an anaerobic environment in soil by limiting soil particles and blocking air respiration in the soil pores. Such anoxic situations affect basic soil chemical and physical indicators and thus, the soil microbial communities [11]. Heavy crude oil contamination can cause the complete mortality of wetland vegetation [12].

North-eastern India, particularly Assam is well-known in the production of petroleum oil. Crude oil pollution is a usual phenomenon in the oil drilling sites as well as the periphery areas through which oil transportation pipelines carry the crude oil to the oil collecting stations. Large,

well-established reports are available regarding the effects of crude oil pollution on soil physicochemical properties and different stabilization methods have been proposed as a remedy for such soil to regain its properties at the regional level [13]. However, the literature regarding the geographical position of oil drilling sites and the effects of crude oil on the horizontal distribution of soil chemical constituents is still inadequate. Hence, an effort has been made to investigate some realistic information regarding the effects of crude oil pollution on soil physicochemical properties and to compare the horizontal variation of different physicochemical parameters in paddy soil around the Sildubi oil spill in Borhola, Assam.

2. MATERIALS AND METHODS

2.1 Study Location

The study location of this investigation was the Sildubi oil field area in the Jorhat district of upper Assam, where oil exploration activities are conducted by Oil & Natural Gas Co-operation Limited (ONGCL). Geographically, the south-east side of this oil spill was hilly and the other sides were low-land rice fields.

2.2 Collection of Soil Samples

Soil samples were collected at 0–15 cm depth from the top surface using a soil auger. Soil collected from the drilling site is represented as DL site. Four samples were collected from North (N), South (S), East (E) and West (W) zones at radius of 500 m from drilling site in circular mode and another set of four samples were collected at 1000 m radius in similar way. Each sample was mounted in replicate quadrants, from which samples were collected. The collected samples were put into polythene bags and taken to the laboratory for analysis after labelling.

2.3 Preparation of Soil Samples

In the laboratory, the samples were spread on brown paper and dried in open air and then ground with a mortar pestle, taking the precaution not to break down the ultimate particles of soil.

2.4 Analysis of Soil Physicochemical Properties

In this study, the analysis of soil samples was carried out according to standard methods using

analytical-grade reagents. The analysis process of different parameters of the soil samples is discussed below:

The colour determination was made by visual observation. The pH was tested in deionised water at a 1:3 soil - water solution ratio by using a calibrated pH meter and moisture content was determined gravimetrically by weighing samples before and after oven-drying at 105°C for 24 h [14]. The bulk density (ρ_d) of the soil was determined from the core samples [15]. Total organic carbon (TOC) and total organic matter (TOM) were also determined in the laboratory by the titrimetric method of Walkey and Black [16]. Available nitrogen was determined by the distillation method [17]. The K, Na and Ca were measured with a flame photometer [18]. Soil iron and phosphorous was determined by the spectrophotometric method as outlined by Jackson [18] at 540 and 660 nm, respectively.

Statistical analyses of experimental data were carried out using SPSS software. An analysis of variance was carried out to test the significance of the treatment effect. The F-test, coefficient of variance and critical difference were calculated by the standard method [19].

3. RESULTS AND DISCUSSION

3.1 Soil Bulk Density

Soil bulk density is a function of soil textures, structures, and organic matter content, but in a given soil type, it can be used to monitor the degree of soil compaction and peddling [20]. Changes in soil bulk density may affect water and oxygen supply to the host. In this investigation, significant horizontal variations were observed in soil bulk density. Lower soil bulk density was recorded in the sample collected from the drilling site compared to the samples of 500 and 100m radius (Table 1 and Fig. 1A). Soil samples collected at 1000m radius showed high compactness ($\rho_b = 1.243 \text{ g/cm}^3$) compared to soils at 500m radius (Table 1). In contrast to the results of previous studies [21,20], a lower bulk density in crude oil-contaminated soil was recorded in our study. The effect of crude oil contamination on soil physicochemical properties may also fluctuate with sampling season [10] and location of the oil spills. In the case of upland oil spills, due to heavy rainfall in the summer season, the effect of crude oil pollution may be more pronounced in nearby lowland rice field soils. In this investigation, the

Table 1. Effect of crude oil contamination on soil physical parameters in the samples collected from different sites

Sample collection sites	Physical Properties					
	Colour		Bulk Density (mg.cm ⁻³)		% Moisture	
	500 m	1000 m	500 m	1000 m	500 m	1000 m
East	Light Brown	Pale Brown	1.168±0.005	1.396±0.002	6.213±0.028	17.255±0.03
West	Light Brown	Brown	1.287±0.003	1.372±0.002	9.135±0.027	16.66±0.03
North	Light Brown	Light Brown	1.308±0.002	1.472±0.002	23.455±0.06	18.625±0.32
South	Brown	Brown	1.334±0.003	1.228±0.001	21.42±0.27	25.127±0.1
Drilling site	Black		0.763±0.01		3.855±0.02	
Mean			1.172±0.11	1.246±0.13	12.816±5.76	16.303±3.7
CV			1.11%	7.18%	7.73%	2.11%
F value			1790.87***	3217.72***	330.81***	2.23.61***
CD at 0.01%			0.04	0.027	3.024	1.049

*** represent the significance at 0.1% level of probability




Fig. 1. Effect of crude oil contamination on (A) bulk density (mg.cm⁻³), (B) % moisture content and (C) soil pH in the samples collected from different sites. The vertical bars represent the mean standard error. The ±SEM values in the error bars for (A) and (B) are shown as a multiple of 10

*** represent the significance at 0.1% level of probability

geographical position of the Sildubi oil spill might be the major reason for the low bulk density value in the soils on the oil drilling side.

3.2 Moisture

Moisture content is an important physical indicator of fertile soil. Soils with very low moisture content show a stressful condition for plant growth. Particle size, void ratio, clay minerals, organic matter and groundwater conditions determine the soil moisture.

In this study, soil moisture contents at the contaminated site were appreciably lower than the samples collected at 500 and 1000m radii. The samples collected from 1000m, showed comparatively high moisture content (Table 1 and Fig. 1B). It is evident from the results that the crude oil, which penetrates the soil interlayer, prevents top-down movements of water. Similar results were reported by Rowell, [22] and De Jong [23] in their observations who advocated

that the water-holding capacity of crude oil-contaminated soil was directly proportional to porosity. They recorded lower water holding capacity at the highly polluted site with reduced porosity. The formation of hydrophobic crude oil coating above the soil surface might have blocked soil pores and reduced the permeability of water and air [24,10,25,26,27], thus affecting the soil water content.

3.3 Soil pH

Surface soil pH plays a vital role in ensuring soil fertility and the availability of other plant nutrients like Fe, Mn, Zn and Cu, which are more available in acidic than alkaline soils. Hence, pH is one of the most significant physical properties of soil and from a farmer's point of view; it is an important soil parameter for crop production.

The soil pH value at the drilling site was 7.85 and was significantly higher than the samples collected from 500m and 1000m radius (Fig. 1C)

and differs from the results of previous studies that showed an increase in petroleum hydrocarbon concentrations in soil could lower pH values [8]. There was significant horizontal variation on the soil pH values among the sampling sites except the samples collected from north (Table 2). The pH values of the samples collected at the 1000m radius were lower than at the other sampling sites. The results of previous studies on oilfields [10,9] showed that oil pollution raised soil pH which is supported by our results. The higher pH values in the soils of the drilling site in this study might be due to the hydrophobic nature of crude oil, which might induce a potential drought in the surface layers of polluted soil [28,10]. Moreover, crude oil contamination in soil may intensify the accumulation of exchangeable bases like Ca^{2+} or may reduce the exchangeable acidity [10,29,30], which also supports the increase of pH in crude oil-polluted soil.

3.4 Electrical Conductivity

Electrical conductivity is the measure of ions present in a soil solution. Usually, electrical conductivity varies with depth, but horizontally, the range of variation becomes more prominent between upland and nearby lowland profiles. This probably occurred due to the slope of the land surface, which is responsible for leaching out alkali and alkaline bases [31]. In this study, electrical conductivity is quite variable in the soil samples collected at different radii from the central drilling site (Fig. 2D). Significant variations in horizontal electrical conductivity were recorded in the present study. The lowest electrical conductivity was recorded in oil-contaminated soils at the drilling site. Although high EC was recorded in the soils collected from 1000m radius to the west, the results of other samples at this radius were quite lower.

Moreover, the samples collected at 500m radius showed significantly higher EC compared to the soil samples of drilling site and 1000m radius. Exceptionally high EC was observed in the soils at 500m radius to the north (Table 2). This variation in soil EC might be due to the horizontal flow of alkali and alkaline bases from upland oil drilling sites to near-low-land rice fields for heavy rain in the summer season.

3.5 Soil Organic Carbon and Organic Matter

Organic carbon content was found to be higher in the soils of crude oil-contaminated zones, while the soil samples collected at a 1000m radius were found to be the lowest. Significant horizontal variations in organic carbon content and percentage organic matters were recorded in current investigation (Table 2).

Organic carbon decreases with an increase in horizontal distance. Organic carbon in the soil is generally derived from biota, such as peat formation with time, plant fine root turnover, microbial biomass, etc. However, crude oil contamination in oilfield marshes might also contribute to the TOC in the soil. In the present investigation, the observed increase in the total organic carbon in the soils of the drilling site may be attributed to the accumulation of high hydrocarbon content during oil exploration (Fig. 2E and 2F). Wang et al. [9], in their study, also reported that oil contamination significantly increased the TOC contents, most probably because of the much higher TPH concentration [10]. Similar findings have been reported by Agbogidi et al. [30] who concluded that high organic carbon in crude oil-contaminated soil might be due to the slow decomposition rate by soil organisms since contamination of soil with crude oil might have poor soil aeration.




Fig. 2. Effect of crude oil contamination on (D) conductivity (mS/cm), (E) % organic carbon and (F) % organic matters in the samples collected from different sites. The vertical bars represent the mean standard errors. The \pm SEM values in the error bars for (D) are shown as a multiple of 10

*** represents the significance at the 0.1% level of probability

Table 2. Effect of crude oil contamination on soil chemical properties in the samples collected from different sites

Sample collection sites	Soil Chemical Properties							
	Soil pH		Conductivity(m S/cm)		% Organic Carbon		% Organic Matters	
	500 m	1000 m	500 m	1000 m	500 m	1000 m	500 m	1000 m
East	5.4±0.12	5.53±0.1	0.562±0.001	0.156±0.002	1.05±0.03	0.916±0.01	1.752±0.04	1.588±0.02
West	5.87±0.1	5.38±0.06	0.214±0.001	0.711±0.001	1.007±0.05	1.03±0.03	1.857±0.03	1.595±0.02
North	7.35±0.16	6.78±0.63	1.11±0.01	0.165±0.002	2.004±0.06	0.676±.004	3.467±0.06	1.37±0.18
South	5.57±0.1	5.1±0.1	0.342±0.001	0.364±0.001	1.255±0.08	1.13±0.01	2.022±0.05	2.085±0.04
Drilling site	7.85±0.14		0.248±0.003		2.383±0.09		3.777±0.02	
Mean	6.41±0.5	6.12±0.52	0.495±0.165	0.327±0.1	1.54±0.27	1.23±0.3	2.575±0.43	2.083±0.44
CV	4.16%	2.53%	1.54%	2.82%	5.81%	6.80%	2.75%	8.03%
F value	69.79***	229.14***	9412***	2500.98***	185.31***	252.33***	718.04***	139.45***
CD at 0.1%	0.814	0.473	0.023	0.027	0.273	0.255	0.216	0.518

*** represent the significance at 0.1% level of probability

Our study also agrees with the results of Ekundayo and Obuekwe [32], who reported an increase in the organic carbon content of oil-polluted soils in southern Nigeria. Udo&Faymei [33] also reported that an increase in organic carbon is directly proportional to an increase in crude oil levels in the soil [24].

3.6 Nitrogen

In the present investigation, the total nitrogen content in the soils of the drilling site was found to be higher than the samples collected from 500 and 1000m radii (Table 3 and Fig. 3P). The increase in total nitrogen in the crude oil-contaminated soil is attributed to the impact of crude oil, which promotes the organic carbon content in the soil. Agmi et al. [34], Akamigbo [35] and Essien and John [36] also found a positive relationship between soil organic carbon and total nitrogen in soils. Udo and Fayemi [33] also reported that available nitrogen increases with increased crude oil contamination in the soil. The present results sustain the results of Agim et al. [34]; however, Obasi et al. [37] and Onuh et al. [38] reported a decrease in total nitrogen with an increase in crude oil pollution in the soil. Significant horizontal variations in total nitrogen were observed in the present study. Soil samples collected at a 1000m radius recorded the lowest nitrogen content compared to those observed at a 500m radius from the drilling site.

3.7 Phosphorous

Phosphorous is one of the key micronutrients essential for plant growth and helps in the transfer of energy. Most of the activities of plants, such as growth, respiration and reproduction, depend on the phosphorous levels of the soil where the plants are grown. Soils with high organic matter content have better supplies of organic phosphate for plant uptake.

The results of the present study have recorded distinct horizontal variations in soil phosphorous concentration (Fig. 3Q). The phosphorous concentrations in the soils of the drilling site (2214.22µg/g) were significantly higher than the mean phosphorous concentration in the samples collected at 500m (2080.24µg/g) and 1000m radius (1578.46µg/g). Samples collected at 1000m radius from the drilling site recorded the lowest phosphorous content (Table 3). Moreover, in this study, the highest phosphorous concentrations were detected in the samples collected from the south direction at 500m

(2488.02µg/g) and at 1000m (2185.09µg/g). This southern sampling site is a low-land area suitable for rice cultivation. The increase in phosphorous content in the soils of the drilling site might be due to the increase in soil organic matter and pH. Organic matter is an important factor in controlling phosphorous availability. With the increase in organic matter, the availability of phosphorous increases, this can be explained in two ways. Firstly, the mineralization of organic matter releases plant-available forms of phosphorous into soils, and secondly, organic molecules will compete with phosphate adsorbed to soil surfaces and reduce phosphorous retention. This process will increase the availability of phosphorous. Soil pH between 6 and 7 will result in maximum phosphorous availability. At low pH, soils have greater amounts of iron, which form very strong bonds with phosphate. At high pH, when calcium is the dominant cation, phosphate tends to precipitate with calcium. Higher Ca concentrations in the drilling site, as observed in this investigation, may precipitate phosphorous in the form of phosphates on the soil surface, which were spilled over by petroleum waste frequently. The poor microbial activities to degrade the crude oil hydrocarbons in such anoxic soil conditions might be another reason for the higher concentration of available phosphorous at the drilling site. Similar findings have been reported by Agbogidi et al. [30] and they concluded that the increase in available phosphorous content of the crude oil-contaminated soil might be due to the increase in soil pH resulting from amendment applications. However, Da-Silva and Fitzsimmons [39] reported that alkaline pH could cause flocculation of Ca, Mg and Fe with phosphorous and reduce their availability to plants.

The result of this investigation agrees with the observation of Siddiqui and Adams [40], who reported an increase in phosphorous concentration with increasing concentrations of oil up to a certain limit, and then it declined. Bielski and Ferguson [41] also noted that increasing pH increases weathering and mineralization rates. This could have increased phosphorous availability.

3.8 Potassium

Potassium is considered an important nutrient for plants and plays a key role in a vast array of physiological processes vital to plant growth, from protein synthesis to the maintenance of plant water balance [42]. In the current study, the

concentration of available potassium was found to be the lowest at the oil drilling site (1.945 $\mu\text{g/g}$). The mean potassium content in the soils at 500m radius was found to be highest (2.694 $\mu\text{g/g}$), while the soils at 1000m radius recorded 2.501 $\mu\text{g/g}$ available potassium. Except for the sample of North direction collected at 500m (3.86 $\mu\text{g/g}$), comparable results of the available potassium content were recorded at other sampling sites (Table 3 and Fig. 3R). The significant variations in the horizontal distribution of available potassium content might be due to excessive rainfall during the summer season, which accelerates the rate of parallel nutrient flow from the upland drilling site to the lowland paddy field. Agbogidi et al. [30] had also recorded lower levels of K^+ and Na^+ in crude oil-contaminated soil and reported that the reduction of K^+ and Na^+ might be due to nutrient immobilization, consequent on the formation of complexes in the soil after degradation and uptake.

3.9 Sodium

Sodium is not considered an essential nutrient for the growth of plants; instead, high sodium concentrations build up salt stress in the soil and adversely affect the plants' growth. However, it acts as a counter-ion in the soil where the potassium concentration is very low. In this study, significant horizontal variations in the Na^+ concentration were recorded at different sampling points Fig. 4X. The sodium concentration at the drilling site was 6.492 $\mu\text{g/g}$. The average value for samples at 500m radius was 13.113 $\mu\text{g/g}$, and at 1000m radius it was 6.738 $\mu\text{g/g}$. The highest Na^+ and K^+ concentrations were recorded in the sample collected from north direction at a radius of 500 m. The higher soil pH (pH = 7.35) and electrical

conductivity (1.1mS/cm) in this sample might be the reason for the high accumulation of exchangeable bases like Na^+ and K^+ ions [30]. The alkaline pH of the soils contaminated with crude oil might be attributed to the accumulation of high sodium content that entered the nearby rice fields during heavy rains. The observed increase in salt concentration at 500m may be due to the leaching of Na^+ and K^+ ions from the drilling site to the neighbouring rice field area during the summer season.

3.10 Calcium

Calcium is present in sufficient amounts in most soils. Calcium is a component of several primary and secondary minerals in soil that are essentially insoluble for agricultural purposes. However, calcium is also present in relatively soluble forms as cations adsorbed to the soil colloidal complex, which is considered to be available to crops. Crude oil contamination may increase the concentration of Ca in the soil [31,30]. In this study, significant horizontal variation in soil Ca concentrations was recorded (Fig. 4Y). The highest Ca concentration (402.71 $\mu\text{g/g}$) was detected in the soils of the drilling site, which were heavily contaminated with crude oil waste. The average values of Ca content in the samples collected at 500 and 1000m radius were 338.344 $\mu\text{g/g}$ and 259.603 $\mu\text{g/g}$, respectively (Table 4). The results of this study support the findings of Agbogidi et al. [30], Konwar and Jha [43], and Wang et al. [10]. In that study, the maximum calcium content was estimated in highly contaminated soil and the minimum in control soil. The result of the current study supports the findings of Onojake and Osuji [44], who recorded lower Ca content in soils contaminated with petroleum waste.

Table 3. Effect of crude oil contamination on soil macronutrient concentrations in the samples collected from different sites

Sample collection sites	Concentration of different elements in soil					
	Soil Nitrogen ($\mu\text{g} / \text{gm}$)		Soil K ($\mu\text{g} / \text{gm}$)		Soil P ($\mu\text{g} / \text{gm}$)	
	500m	1000m	500m	1000m	500m	1000m
East	469.25 \pm 13.8	502.85 \pm 3.85	2.322 \pm .02	2.187 \pm .02	2050.09 \pm 6.53	1068.54 \pm 5.7
West	441.5 \pm 10.18	407.25 \pm 4.34	2.89 \pm .04	2.797 \pm .05	1867.9 \pm 3.37	1160.22 \pm 5.5
North	541.25 \pm 10.5	324.5 \pm 9.07	3.86 \pm .032	2.865 \pm .05	1780.99 \pm 9.66	1264.24 \pm 15.8
South	501.5 \pm 5.2	475.25 \pm 3.47	2.225 \pm .03	2.710 \pm .045	2488.02 \pm 11.38	2185.09 \pm 4.7
Drilling site	994.75 \pm 7.8		1.945 \pm .07		2214.22 \pm 10.4	
Mean	589.65	540.92	2.649	2.501	2080.24	1578.46
CV	2.33%	13.59	3.15%	3.99%	8.85%	2.10%
F value	1112.02***	51.15***	309.82***	67.84***	942.75***	1188.07***
CD at 0.01%	42.02	224.46	0.255	0.305	56.23	101.23

*** represent the significance at 0.1% level of probability




Fig. 3. Effect of crude oil contamination on (P) soil nitrogen (µg/g), (Q) available phosphorus (µg/g) and (R) potassium concentration (µg/g) in the samples collected from different sites. The vertical bars represent the mean standard errors. The ±SEM values in the error bars are shown as a multiple of 10

*** Significant at 0.1% level of probability

Table 4. Effect of crude oil contamination on calcium, sodium and total iron concentrations in the soil samples collected from different sites

Sample collection sites	Concentration of different elements in soil					
	Soil Ca (µg / gm)		Soil Na (µg / gm)		Soil Fe (µg / gm)	
	500m	1000m	500m	1000m	500m	1000m
East	394.778±6.12	210.81±.68	15.56±.52	5.415±.23	3990.67±6.1	7659.46±0.5
West	285.9±6.37	275.848±4.66	14.745±.2	6.703±.168	2690.05±6.53	8176.54±0.95
North	283.595±6.55	189.793±5.78	16.768±.15	6.255±.045	2050.85±5.6	7239.59±23.9
South	324.735±2.86	218.853±4.53	11.997±.06	8.823±.12	2646.82±8.65	8078±24.49
Drilling site	402.71±8.27		6.492±.14		2859.17±16.5	
Mean	338.344	259.603	13.113	6.738	2847.91	6802.55
CV	3.54%	4.55%	5.08%	5.03%	3.81%	2.78%
F value	92.248***	212.20***	269.525***	56.724***	169.9***	18927.2***
CD at 0.01%	36.608	36.096	1.72	1.035	331.55	99.23

*** represent the significance at 0.1% level of probability




Fig. 4. Effect of crude oil contamination on (X) sodium concentration (µg/g), (Y) soil Ca concentration (µg/g) and (Z) total iron (µg/g) in the samples collected from different sites. The vertical bars represent the mean standard errors. The ±SEM values in the error bars are shown as a multiple of 10

*** Significant at 0.1% level of probability

3.11 Soil Iron

Iron is the most abundant element in the earth crust, which plays an important role in the mobilization of toxic elements in the environment and functions as an electron carrier in many complex red-ox reactions in the plant body. Iron is a co-factor of many enzymes involved in plant

defence systems. The study of soil iron is very significant because its deficiency or toxicity in soil may cause many physiological and biochemical disorders in the plant body. Above 6,000 soils Fe may cause iron toxicity in paddy rice [45].

Significant variations in the horizontal distribution of soil Fe contents were recorded in this study

(Fig. 4Z). The samples collected at 1000m away from the drilling site recorded the highest mean Fe concentration (6802.55µg/g). The Fe concentrations in the samples collected from the east, west, north and south directions at a 1000m radius were 7659.46µg/g, 8176.54µg/g, 7239.59µg/g and 8078µg/g, respectively (Table 4). A considerable increase in total Fe concentrations in the samples collected at 1000m radius might be due to artificial flooding and the spontaneous flow of rainwater from upland oil spill during rainy season. The total Fe concentrations (2844.59µg/g) in the soils at a 500m radius were lower than the results of the drilling site. Although the amounts of major elements such as Ca, K, Na, Mg and Fe in petroleum waste-contaminated soils primarily depend on the extent of petroleum waste, the effects of other parameters like soil pH, precipitation of these elements in the form of phosphates, hydroxides, oxy-hydroxides and the geographical position of the oil-contaminated area also play an important role in the availability of these elements. In the current study, significant increase in total Fe in the samples collected from 1000m radius might also be attributed to the horizontal flow of the element either through flooding or by soil osmotic properties. Soil heavily contaminated with crude oil shows threshold levels of major elements [31]. Moreover, there are positive correlations between soil pH, available P content and soil Fe concentrations. Ch'ng et al. [46] reported that in an acidic soil environment, if phosphate ion concentrations are high, they compete with Fe and thus Fe becomes unavailable to plants, while pH greater than 7 may cause the flocculation of Ca and Fe with phosphorus in the form of their phosphates, thus reducing their availability to plants. In this study, the high concentration of available phosphorous and alkaline elements in the soil of the drilling site might be the cause for higher concentration of total Fe in the neighbouring agricultural soils. The high concentration of Fe under acidic pH in the peripheral rice field around the oil drilling site might be the reason for severe oxidative stress in rice leaves called 'leaf bronzing', which in turns deplete crop productivity [45].

4. CONCLUSION

From the current study, it can be concluded that the test results obtained from the soil analysis of the oil impacted sites and compared to the results of the soil samples collected from 500 and 1000 m, indicate that the total hydrocarbon

levels observed at the soils of the oil-spilled locations have provided evidence of severe hydrocarbon contamination. The alteration in soil physical and chemical properties under such conditions lowers crop productivity on neighbouring agricultural lands. Significant variations in the results of different parameters were observed among the soil samples collected from different collection zones. Sol pH, organic carbon and organic matter were significantly higher at the drilling site, while other studied parameters were higher in the samples collected from nearby agricultural lands. The significant horizontal variations in soil physical and chemical properties primarily depend on the extent of petroleum waste. Moreover, the effects of other parameters like soil pH, precipitation of elements in the form of phosphates, hydroxides, oxy-hydroxides and the geographical position also accelerate the nutrient alteration in the oil-contaminated area. Among the different mechanisms available to explain the variability in nutrient concentration upon crude oil contamination, flooding, soil osmotic properties, and the formation of hydrophobic crude oil layers over the root surface also play important roles in nutrient modification and their availability for plants' uptake. Similarly, antagonistic relationships among the nutrients and their competitive absorption and mobilization by the plant's chelators might also be the cause of specific nutrients unavailability under crude oil contamination. In this study, high concentrations of Fe and phosphorous observed due to crude oil contamination have been developing Fe stress situations, either by Fe deficiency in alkaline pH or by toxicity in acidic pH, thus sinking crop production in neighbouring agricultural lands.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Salimnezha A, Soltani-Jigheh H, Soorki AA, Effects of oil contamination and bioremediation on geotechnical properties of highly plastic clayey soil. *Journal of Rock Mechanics and Geotechnical Engineering*. 2021;13:653–670.
2. Okafor UC, Evaluation of the impact of crude oil contamination on soil's physicochemical characteristics, Microflora and Crop Yield. *Tropical Aquatic and Soil Pollution*. 2023;3(1):24-35.

3. Zhang S, Li R, Bai W, Yang Q. Effects of Oil Contamination on the Physical-Mechanical Behavior of Loess and Its Mechanism Analysis. *Geofluids*. 2021(3): 1-12.
4. Borah G, Deka H. Crude oil associated heavy metals (HMs) contamination in agricultural land: Understanding risk factors and changes in soil biological properties. *Chemosphere*. 2023;310: 136890.
5. Abosede EE. Effect of crude oil pollution on some soil physical properties. *Journal of Agriculture and Veterinary Science*. 2013;6(3):14–17.
6. Quyum A, Achari G, Goodman R. Effect of wetting and drying and dilution on moisture migration through oil contaminated hydrophobic soils. *Science of the Total Environment*. 2002;296(1):77–87.
7. Ekundayo E, Obuekwe O. Effects of an oil spill on soil physico-chemical properties of a spill site in a typicudipsamment of the Niger delta basin of Nigeria. *Environmental Monitoring and Assessment*. 2000;60(2): 235–249
8. Kusic I, Mesic S, Basic F, Brkic V, Mesic M, Durn G, Zgorelec Z, Bertovic L. The effect of drilling fluids and crude oil on some chemical characteristics of soil and crops. *Geoderma*. 2009;149(3–4):209–216.
9. Wang XY, Feng J, Zhao JM. Effects of crude oil residuals on soil chemical properties in oil sites, Momoge Wetland, China. *Environmental Monitoring and Assessment*. 2010;161(1):271–280.
10. Wang Y, Feng J, Lin Q, Lyu X, Wang X, Wang G. Effects of crude oil contamination on soil physical and chemical properties in momoge wetland of China. *Chin. Geogra. Sci*. 2013;23(6):708–715.
11. Sutton NB, Maphosa F, Morillo JA, Abu Al-Soud W, Langenhoff AA., Grotenhuis T, Rijnaarts HH, Smidt H. Impact of long-term diesel contamination on soil microbial community structure. *Applied and Environmental Microbiology*. 2013;79(2): 619–630.
12. Lin Q, Mendelssohn IA. Impacts and recovery of the deepwater horizon oil spill on vegetative structure and function of coastal salt marsh in the northern Gulf of Mexico. *Environmental Science & Technology*. 2012;46(7):3737–3743.
13. Ezeokpube GC, Ahaneke IE, Alaneme U, Attah IC, Etim RK., Olaiya BC, Udusoro IM. Assessment of mechanical properties of soil-lime-crude oil-contaminated soil blend using regression model for sustainable pavement foundation construction. *Advances in Materials Science and Engineering*. 2022;(1):1-18.
14. Zhang C, Nie S, Liang J, Zeng G, Wu H, Hua S, Liu J, Yuan Y, Xiao H, Deng L, Xiang H. Effects of heavy metals and soil physicochemical properties on wetland soil microbial biomass and bacterial community structure, *Science of the Total Environment*. 2016;557-558:785–790.
15. Ma YY, Lei TW, Zhang XP, Chen YX. Volume replacement method for direct measurement of soil moisture and bulk density. *Trans Chin Soc Agric Eng (in Chinese)*. 2013;29:86–93
16. Walkley AJ, Black IA. Estimation of soil organic carbon by the chromic acid titration method. *Soil Sci*. 1934;37:29-38.
17. Bremner JM. Nitrogen availability indexes IN Methods of soil analysis. Part 2. *Agronomy 9, Am. Soc. of Agron. Medison, Wisconsin*. 1965;1324-1345.
18. Jackson ML. *Soil chemical analysis*, prentice-hall of India Private limited, New Delhi. 1973;227-255.
19. Cochran WG, Cox GM. *Experimental Design*. John Wiley and Sons Inc., New York; 1963.
20. Oyem ILR, Oyem IL. Effects of crude oil spillage on soil physico-chemical properties in ugborodo community. *International Journal of Modern Engineering Research (IJMER)*. 2013;3(6): 3336-3342
21. Agmi LC, Amos HI, Osi AF, Okoli HN, Egboka NT. Lethality of crude oil contamination on properties of soil and growth parameters of maize (*Zea mays L.*). *Int J Environ Sci Nat Res*. 2021;28(3).
22. Rowell MJ. The effects of crude oil spill on soil - A review. In: *Reclamation of agricultural soils after oil spills, Part I: Eds., Toogood J.A., Edmoter, Deptt. of Soil Science, University of Alberta*. 1977;1-13.
23. De Jong E. The effects of crude oil spills on cereals. *Env. Pollut*. 1980;22:187-196.
24. Baruah D. Physico-chemical properties of soil and quantitative analysis of a herbaceous community after blowout of an

- oil well, Nature Environment and Pollution Technology. 2007;6(2):251-258.
25. Bennett PC, Siegel DE, Baedecker MJ, Hult MF. Crude oil in a shallow sand and gravel aquifer—I. Hydrogeology and inorganic geochemistry. Applied Geochemistry. 1993;8(6):529–549.
 26. Roy JL, McGill WB, Rawluk MD. Petroleum residues as water-repellent substances in weathered nonwettable oil-contaminated soils. Canadian Journal of Soil Science. 1999;79(2):367–380
 27. Khamsehchiyan M, Hossein Charkhabi A, Tajik M. Effects of crude oil contamination on geotechnical properties of clayey and sandy soils. Engineering Geology.2007; 89(3):220–229.
 28. Njoku K, Akinola M, Oboh B. Phytoremediation of crude oil contaminated soil: The effect of growth of Glycine max on the physico-chemistry and crude oil contents of soil. Nature and Science. 2009;7(10):79–87.
 29. Osuji LC, Egbuson EJ, Ojinnaka CM. Assessment and treatment of hydrocarbon inundated soils using inorganic nutrient (NPK) supplements: II. A case study of Eneka oil spillage in Niger Delta, Nigeria. Environmental Monitoring and Assessment. 2006;115(1):265–278.
 30. Agbogidi OM, Eruotor PG, Akparobi SO, Nnaji GU. Evaluation of crude oil contaminated soil on the mineral nutrient elements of maize (*Zea mays* L.). Journal of Agronomy.2007;6(1):188.
 31. Shehla S, Samina S, Irfan UJ. Major elements distribution in the soil profile of petroleum contaminated soils. Journal of Himalayan Earth Sciences. 2018;51(2A): 120-131. 34.
 32. Ekundayo EO, Obuekwe CO. Effects of an oil spill on soil physico-chemical properties of a spill site in a Typic Paleudult of midwestern Nigeria. Environmental Monitoring and Assessment. 1997;45(2): 209–221.
 33. Udo EJ, Fayemi AA. The effects of oil pollution of soils on germination, growth and nutrient uptake of corn. Journal of Environmental Quality.1975;4: 537-540.
 34. Agim LC, Osuji GE, Onweremadu EU, Ndukwu BN, Osuaku S. Seasonal dynamics of soil organic carbon and total nitrogen in soils under different land uses in Owerri Southeastern Nigeria. Agro Science Journal of Tropical Agriculture, Food, Environment, and Extension.2012; 11(1):43-54.
 35. Akamigbo FOR. Influence of land uses on soil physical properties of the humid tropical ecology of south eastern Nigeria. Nigeria Agricultural Journal. 1999;30:59-76.
 36. Essien OE, John IA. Impact of crude oil spillage pollution and chemical remediation of agricultural soil properties and crop growth. Journal of Applied Sciences Management. 2010;14(4):147-154.
 37. Obasi NA, Eberechukwu E, Anyanwu DI, Okorie UC. Effects of organic manures on the physicochemical properties of crude oil polluted soils. African Journal of Biochemistry Research. 2013;7(6):67-75.
 38. Onuh MO, Maduekwe DK, Ohia GU. Effects of poultry manure and cow dung on physical and chemical properties of crude oil polluted soil. Science World Journal. 2008;3(2):45-50.
 39. Da-Silva Cerozi B, Fitzsimmons K. The effect of pH on phosphorus availability and speciation in an aquaponics nutrient solution. Bioresource Technology. 2016; 219:778-781.
 40. Siddiqui S, Adams WA. The fate of diesel hydrocarbons in soils and their effect on the germination of perennial ryegrass. Environmental Toxicology. 2002;17:49 – 62.
 41. Bielski RL, Ferguson IB. Physiology and metabolism of phosphate and its compounds in inorganic plant nutrition. Encyclopedia Plant Physiology. 1983;5: 422–449.
 42. Agbogidi MO, Eshegbeyi OF. Performance of *Dacryodes edulis* seeds and seedling in crude oil-contaminated soil. Journal of Sustainable Forestry. 2006;22(3-4):1-13.
 43. Konwar D, Jha DK. Response of Rice (*Oryza sativa* L.) to Contamination of Soil with Refinery Effluents under Natural Conditions. Assam University Journal of Science & Technology. 2010;5(1):14-22
 44. Onojake MC, Osuji LC. Assessment of the physico-chemical properties of hydrocarbon contaminated soil. Archives of Applied Science Research. 2012; 4(1):48-58.

45. Saikia T. Baruah KK. Iron Toxicity Tolerance in Rice (*Oryza sativa*) and Its Association with anti-oxidative Enzyme Activity. Journal of Crop Science. 2012; 3(3):90-94.
46. Ch'ng HY, Ahmed OH, Majid NMA. Improving phosphorus availability in an acid soil using organic amendments produced from agro industrial wastes; 2014.

© 2023 Saikia et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/108464>