

International Journal of Environment and Climate Change

Volume 14, Issue 2, Page 407-422, 2024; Article no.IJECC.112900 ISSN: 2581-8627 (Past name: British Journal of Environment & Climate Change, Past ISSN: 2231–4784)

Impacts of Climate Change on Teesta River Basin Char Lands: A Study of Livelihood and Ecosystem Dynamics in Bangladesh

Rozina Parvin ^a, Md Shafiqul Bari ^b and Noor Muhammad ^{c,d*}

^a Faculty of Science and Technology, Bangladesh University of Professionals, Dhaka-1216, Bangladesh.

^b Department of Agroforestry and Environment, Hajee Mohammad Danesh Science and Technology University, Dinajpur-5200, Bangladesh.

^c Rural Development Academy, Bogura-5842, Ministry of Local Government, Rural Development and Cooperatives, Bangladesh.

^d Department of Agriculture, King Abdulaziz University, Jeddah-80208, Kingdom of Saudi Arabia.

Authors' contributions

This work was carried out in collaboration among all authors. Author RP designed the research work, developed the protocol, conducted field survey and wrote the first draft of the Manuscript. Author MSB developed the conceptual framework and supervised the research. Author NM did the statistical analyses and edited the manuscript. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJECC/2024/v14i23956

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <u>https://www.sdiarticle5.com/review-history/112900</u>

> Received: 01/12/2023 Accepted: 05/02/2024 Published: 10/02/2024

Original Research Article

ABSTRACT

This study anticipates identifying how does climate change affect the livelihood and ecosystem of char land of Teesta River basin char land. Structured questionnaire and related documents were used for both primary and secondary data collection. Survey was conducted in two districts

*Corresponding author: E-mail: noormd.rda@gmail.com;

Int. J. Environ. Clim. Change, vol. 14, no. 2, pp. 407-422, 2024

(Rangpur and Nilphamari) and four unions (Lakhitari, Topa Madhupur, Shatibari, Gulmund) of four upazilas (Gangachara, Kaunia, Dimla, Jaldhaka). A total of 50 respondents from each selected char union was selected following a multistage random sampling procedure. Thus, there was altogether 200 respondents selected for this study. Data related to age, education level, occupation, knowledge about climate change of char inhabitants and perception of char dwellers regarding climate change was measured. A total number of 93% of the respondents clearly grasp that climate change refers to changes in long-term typical weather. It was observed that ongoing climate change has 78% impact on agriculture followed by health of peoples and other animals that is 20%. In the rainy season they observed increase in lightning from sun as the highest effect of climate change that has a WAI of 1.77. Teesta river basin people perceived dry spell frequency as the second highest extent of climate change that had a WAI of 1.74 followed by timing of rain offset. uneven distribution of rainfall and timing of rain onset. According to their perception it was found that great extent of loss was found in skills of char peoples (48.35%) followed by knowledge of char peoples (44.27%). It was also observed that a third great extent of loss was found in char peoples good health (41.67%) followed by ability to work (39.06%). Out of all the respondents 166 (43.2%) observed that soil is becoming dry day by day due to climate change followed by rivers becoming dry (37.8%) (Table 8). They found that growing trees have become difficult (45.1%), due to drought underground water is decreasing (61.7%) and air became dry due to climate change, and these are small extent of loss. Char inhabitants observed that climate change impact on physical capital and great extent impact was perceived on people's migration to another place (41.4%). Teesta River basin char dwellers opined about 24 adaptation strategies to reduce the impact of climate change on livelihood and ecosystem in that area.

Keywords: Livelihood; ecosystem; climate change; char land; Teesta River Basin.

1. INTRODUCTION

1.1 Background of the Study

Bangladesh has achieved significant progress over the last decade, and the country is currently on its path to transitioning to a developing economy [1]. However, due to the unequal distribution of resources and development efforts, not everyone has benefited [2]. Due to limited access and rights to resources and even basic requirements, some people in the country tend to remain marginalized [3,4]. Furthermore, this situation is deteriorating as the frequency and severity of various climatic problems such as cyclones, floods, river erosion, heavy rain, and drought has increased. People in Bangladesh are increasingly vulnerable due to social, economic, and political reasons, as well as the climate calamity.

Bangladesh is one of the South Asia's most vulnerable countries to climate change due to its physical, social, and economic characteristics [5]. Natural hazards such as floods, erosion, cyclones, heavy rain, drought, and other natural disasters occurrence in Bangladesh is very common, and climatic variability is one of the main reason for these catastrophes [6,7]. Bangladesh is a country with extreme geographical vulnerability, with 70% of the population living in flood-prone areas and 26% in cyclone-prone areas [8]. In Bangladesh, significant natural calamities such as floods, erosion, and cyclones have displaced around 39 million people since 1970 [9]. Experts predict that by 2050, roughly 6 to 8 million more Bangladeshis may be relocated as a result of rising global temperatures and sea levels [10].

Chars are home to over 10% of the world's population [11]. Furthermore, it is estimated that the char's area, which encompasses roughly 7200km2, is home to 4-5% of Bangladesh's population [12]. There are 56 large and 226 little chars in this country. Bangladesh's char land is often vulnerable to multiple disasters due to its complex environment [12] and the people who live there are the ones who are at most risk. Approximately 12 million char dwellers in Bangladesh are affected by annual floods, erosion, and poverty. Char land, on the other hand, is remote and vulnerable land, with nearly 80% of its residents living in extreme poverty. since they don't own land. More than 70% of the char land population comprises of farmers and fishermen [13].

Nearly 14 percent of the total cultivated area of Bangladesh is covered by the Teesta flood plain and it provides livelihood opportunities to approximately 7.3 percent of the population. A total number of 9.15 million people's live in this flood plain areas in five districts of Rangpur Division (Gaibandha, Kurigram, Lalmonirhat, Nilphamari, and Rangpur). This river water is being used in various sectors of human life including livelihood, irrigation, fisheries and household uses. For their livelihood around 70% of these people of northern part of Bangladesh are directly dependent on the river. The Teesta flood plain basin is mainly agrarian. Rice, wheat, maize, pumpkin, jute, potato is grown in this area as sole crop. Rubber and tea are cultivated as supplementary crops. Char areas are often called as the place of multiple vulnerabilities due to access of education, medicine and even daily basic needs are inadequate and insufficient.

1.2 Problem Statement

The inhabitants of Teesta River basin char were heavily dependent on fishing on the Teesta River. The river was very rich in fish and the char land people never returned empty-handed [14]. Every day in the dry season, they caught many kinds of fish and cultivated rice in Char land (a small island on the river). The soil was fertile and suitable for agriculture, and six different seasons seemed to be a blessing to the people. There was seasonal rainfall at the time, it was easy to adjust in winter, and summer temperatures were not as high as they are today. Agriculture and fishing were the main sources of income for the villagers of the Teesta River basin. The Teesta River was small but very deep and had water all vear round. The availability of natural resources. weather, seasonality, and livelihoods changed depending on the resources of the river. Today, many families live in shelters on the banks of the Teesta River, and it has become difficult for them to meet their basic needs [15]. They face problem of food security, and many are not able to get three meals a day.

Climate change is already affecting biodiversity, and it is expected to become a more serious hazard over the next few decades. The melting of Arctic Sea ice poses a threat to biodiversity throughout the ecosystem and beyond. Water scarcity is found in the char regions due to dryness of Teesta during the pre- and postmonsoon seasons each year. It limits the livelihood options available to the char dwellers. River-dependent livelihood options such as fishing, and boating become unavailable due to the dryness. As a result, char dwellers depend on agriculture as a major means of earning a living. climate change floods, droughts, storms, hailstorms, char erosion, erratic rainfall, cold waves, and climate variability have tremendous impact on the livelihood and ecosystem of Teesta River basin char land. Climate change impacts their daily life and livelihood. The char people are mainly involved in agriculture and often they seek different sources of income such as livestock rearing, daily wage labor, handicraft etc. Char peoples are below poverty line. To cope up with the changed climate they are required to live through this hard situation. It is needed to find out some climate smart ways and strategies to confirm their better survival.

Bangladesh is extremely vulnerable to climate change due to hydroelectric dams. Geological and socioeconomic factors, such as: (a) its location in the world; (b) the kind of geology; (c) the type of geology. The majority of the population relies on crop cultivation, which has heavily influenced by climate change. been Climate variability and hazard are two terms that are often used interchangeably. Climate change is already having a negative influence on agriculture, biodiversity, extreme environmental dangers, and socioeconomic situations of Bangladesh. As a result, residents of the lower Teesta basin had to deal with drought and flood in the same year. The Teesta literature does not provide a detailed and uniform depiction of the key time or adaptation ways.

Rangpur, Nilphamari, Kurigram, Ghaibanhha, Mymensingh, Jamalpur. Sirajgonj, Bogura. Raishahi, Chapai Nawabgonj, Pabna, Taingail, Shariatpur, Faridpur, Barishal, Patuakhali, Blola, Manikgonj, Munshigonj, and Noakhali are the major Char populated districts of Bangladesh. A vast number of people live in these Char areas and rely on Char-based farming systems for their livelihood. As a result, an integrated approach of crop and trees cultivation is required to increase productivity, maintain ecological balance, and improve the socioeconomic status of the Char land people [16]. Teesta river basin char areas as highlighted tremendously have been vulnerable areas. Several natural disasters such as floods, drought, river erosion etc. are common in these areas. In the char areas, more than 60 percent of people do not have any cultivated land other than homestead only. Therefore, for improving the livelihood of the char land people, increasing productivity of the homestead farming system is necessary [14]. Moreover, there is huge scope to introduce improved cropland agroforestry production systems in char land area of Teesta River basin which may ensure sustainable environment friendly climate resilience land use system. In char land area climate smart well adapted agricultural option is much needed. No systematic and comprehensive study about the impact of climate change on agriculture, eco-system and livelihood has been conducted . So, considering the fact a study on the impact of climate change on the livelihood and eco-system of the Teesta River char land of Bangladesh was conducted.

2. METHODOLOGY 2.1 Study Area Selection

The study area was specially selected due to location advantage, flood and drought risks, topographical properties and population diversity. The survey was conducted in the Union of Lakhitari in Gangachara upazila and Union of Topa Madhupur in Kaunia Upazila of the Rangpur district and in the Union of Kha Khagibari of the Dimla Upazila and Union of Gulmonda in Jaldhaka Upazila in the Nilphamari district (Fig. 1). Both districts are located in the lower part of the Teesta basin.

2.2 Data Collection Methods

Data was collected with the help of a pre-tested questionnaire. Structured questionnaire was applied for survey to detect how does climate change influence the livelihood and ecosystem of Teesta River basin char land. In this research interview was taken about the biodiversity as well as ecosystem of the selected char land and some climate related data were collected from weather office and from some NGOs.



Fig. 1. Selection of study area



Fig. 2. Map of study area

2.3 Sampling Technique

Four unions of four Upazilas of two districts were surveyed for the study. By using multistage random sampling procedure, a total of 96 respondents from each selected char union were selected. As the population of the study was heterogenous and simple random sampling method was used, large number of samples were selected to ensure representation of the population. The following well-known statistical formula was used to determine the sample size from an known population [17].

$n_0 = \frac{Z^2 p q}{e^2}$	
$\mathbf{n}_0 = ((1.96)2 \text{ x } .5(.5)) / (.05)2$	Where n_0 is the sample size, Z^2 is the abscissa of the normal curve that cuts off an
= (3.8416 x .25) / .0025	area α at the tails. (1 – α) equals the desired confidence level, e.g., 95%).
=.9604 / .0025	e is the desired level of precision, p is the estimated proportion of an attribute that is
=384.16	present in the population, and q is 1-p. The value for Z is found in statistical tables which
=384	contain the area under the normal curve. e.g., $Z = 1.96$ for 95 % level of confidence

Table 1 Distribution of sample size

SI No	Name of Districts	Name of Upazilas	Name of Unions	No. of Household	No. of Population	Household Proportion (%)	Household no./Respo ndents
1	Rangpur	Gangachara	Lakhitari	5262	28165	20.95%	384x20.95 % =81
2		Kaunia	Tepamadhup ur	9179	35633	36.54%	384x36.54 %=140
3	Nilphamari	Dimla	Khaga Kharibari	6100	31200	24.30%	384x24.30 %=93
4		Jaldhaka	Gulmund	4573	36305	18.21%	384x18.21 % =70
		Т	otal	25114	94998	100%	384

2.4 Data Collection Process

Primary data was collected on the spot in the research location. To acquire both quantitative and qualitative data, a questionnaire survey was undertaken in the research region. Both primary and secondary data was used in the research study. Primary data is collected from the household of Lakhitari, Topa Madhupur, Khaga Khagibari and Gulmunda Union during study period. Furthermore, observation and group discussions were done to acquire additional and extra information concerning the influence of climate change on the livelihood and ecosystem of the study region's char land. Primary data for the study was collected through a survey, one-on-one interviews, and focus group discussions. To detect the climate change impact some of the climate related data was collected from different GOs and NGOs office which was treated as secondary sources of data. Secondary data on the study area's physical setting, total household population, institutional framework, policy guidelines, and maps were obtained from a variety of government and non-governmental organizations, including the published Government statistical yearbook, physical map from LGED, Upazilla Agriculture Office, and Land Office, and others. on the other hand, data was gathered from a variety of government agencies, non-governmental organizations, and published works such as reports, books, records, journals, maps, and papers from various libraries.

3. RESULTS AND DISCUSSION

3.1 Demographic Characteristics of the Study Area

Demographic characteristics like age, education level, occupation, annual income, and farm size of the respondents of the Teesta River basin char land is not same. These factors significantly influence the impact of climate change on livelihood and ecosystem of that char unions as well as affect the mitigation strategies.

3.1.1 Age of the respondents

The result of the household survey showed that the maximum number of respondents (n=166) belong to the age category of 30-40 years. The second largest age category was > 60 years and there were 110 respondents in this category. There were 80 respondents in the 50-60 years age category. also 78 respondents were found to be in category of 41-49 years (Table 2).

3.1.2 Education level of the respondents

The respondents' education level was calculated in terms of their achievement of certificate. The result showed that 78.39% of respondents have recognized education and the rest 21.61% respondents have no formal education. Among the literate respondents 103 respondents had primary level education, 93 respondents had secondary level education, 62 respondents achieved higher Secondary School education, and 43 respondents were Graduate (Table 3). Similar results observed by Karim et al., [14]. Huda et al., [15]. and Haque et al., [18].

3.1.3 Occupation of the respondents

Respondents were involved in different occupations in the study villages. Agriculture is the key activity in both the villages. Besides this, fishing, stone collecting, boating and daily labor are major activities for northwestern char land of Bangladesh. Analysis of the respondents' (n=384) occupation showed that 158 engaged respondents were primarily in agriculture as a primary occupation followed by small business (107), day labour (33), private service (23). On the other hand, 21 respondents are housewife and other 21 respondents are involved in other activities like Van/ rickshaw puller, auto/nosimon driver. Among them 5 respondents reported that they do govt service and 9 respondents retired from govt service. Another 5 respondents were engaged in livestock rearing activities and out of all the respondents, only one respondent was involved in fishing activities and another 1 respondent is engaged in handicraft.

Age	_	Study Unions									
Group	Lakh	itari	Tepan	Tepamadhupur I		Khaga Kharibari		Gulmund			
	HH	%	HH	%	HH	%	HH	%	HH	%	
30-40	22	27.2	42	30.0	21	22.6	31	44.3	116	124.1	
41-49	20	24.7	22	15.7	16	17.2	20	28.6	78	86.2	
50-60	20	24.7	26	18.6	17	18.3	17	24.3	80	85.9	
>60	19	23.5	50	35.7	39	41.9	2	2.9	110	104	
Total	81	100	140	100	93	100	70	100	384	400	

Table 3. Education level of the respondents

Table 2. Distribution of respondents by age

Education Level	_	Study Unions									
	Lakh	khitari Tepamadhupur		Khaga Kharibari		Gulmund		Total			
	HH	%	HH	%	HH	%	HH	%	HH	%	
Illiterate and No Formal education	29	35.8	11	7.9	28	30.1	15	21.4	83	95.2	
Primary	23	28.4	16	11.4	42	45.2	22	31.4	103	116.4	
Secondary	21	25.9	42	30.0	15	16.1	15	21.4	93	93.4	
Higher Secondary	5	6.2	43	30.7	5	5.4	9	12.9	62	55.2	
Graduate	3	3.7	28	20.0	3	3.2	9	12.9	43	39.8	
Total	81	100	140	100	93	100	70	100	384	400	

Occupation	Study Unions									
	Lakhitari		Tepar	Tepamadhupur		Khaga Kharibari		und	Total	
	нн	%	НН	%	нн	%	нн	%	нн	%
Housewife	6	7.4	7	5.0	2	2.2	6	8.6	21	23.2
Agriculture	36	44.4	56	40.0	41	44.1	25	35.7	158	164.2
Small business	24	29.6	40	28.6	24	25.8	19	27.1	107	111.1
Day labour	8	9.9	17	12.1	5	5.4	3	4.3	33	31.7
Private service	6	7.4	3	2.1	8	8.6	6	8.6	23	26.7
Others	0	0	8	5.7	12	12.9	1	1.4	21	20
Fishing	1	1.3	0	0	0	0	0	0.0	1	1.3
Retired	0	0	6	4.3	1	1.1	2	2.9	9	8.3
Government service	0	0	0	0	0	0	5	7.1	5	7.1
Livestock rearing	0	0	3	2.2	0	0	2	2.9	5	5.1
Handicraft	0	0	0	0	0	0	1	1.4	1	1.4
Total	81	100	140	100	93	100	70	100.	384	400

Table 4. Occupation of the respondents

3.1.4 Annual income

The income of the respondents was classified into four categories. (Table 5) shows that, among 384 respondents, 188 char inhabitants have income that is upto 1 lac where 101 char peoples have highest annual income that is more than 3 lac. There are 79 respondents' who have annual income 1-2 lac and 16 respondents who have annual income was 2-3 lac.

3.1.5 Farm size

In Teesta River basin char land, highest landowner were found in Khaga Kharibari union (212 decimals) followed by Tepamadhupur union (170 decimals). The lowest landowner were found in the Gulmond union that was 72 decimals. Moderate landowner was observed in Lakhitari union (160 decimals). Similar results observed by Huda et al. [19].

Annual	_				Stu	Study Unions						
Income	Lakhitari		Tepamadhupur		Khaga Kharibari		Gulmund		Total			
	НН	%	HH	%	НН	%	нн	%	HH	%		
Upto1 lac	55	67.9	41	29.3	58	62.4	34	48.6	188	208.2		
1 lac -2 lac	24	29.6	2	1.4	27	29.0	26	37.1	79	97.1		
2-3 lac	0	0	11	7.9	2	2.2	3	4.3	16	14.4		
More than 3 lac	2	2.5	86	61.4	6	6.5	7	10.0	101	80.4		
Total	81	100	140	100	93	100	70	100	384	400		

Table 6. Farm size

Farm Size		Study Unions							
(Average/HH)	Lakhitari	Tepamadhupur	Khaga Kharibari	Gulmund					
Homestead (Decimal)	120	115	132	62					
Others	40	55	80	10					
(Decimal)									
Total	160	170	212	72					

3.2 Perception on Climate Change

3.2.1 Knowledge about climate change

Climate change, as defined by the Eco life dictionary, is a long-term change in local or worldwide weather patterns. Contrary to global warming, which only refers to one component of climate change a rise in the earth's surface temperature climate change refers to modifications to a region's general weather patterns, such as changes in precipitation, temperatures, cloud cover, and other factors. However, according to the respondents, there is no concrete definition or understanding of what constitutes climate change. A total number of 93% of the respondents clearly grasp that climate change refers to changes in long-term typical weather. They were unsure of the time frame. About 3% of respondents have no idea about climate change that will affect the average weather during the next 20 to 30 years (Fig. 3). Similar results were observed by Sultana et al., [20].

3.2.2 Reasons of climate change

The Teesta River basin char dwellers were asked to find out the different reasons of climate change. According to perception of the respondents, it was found that 42% of the people believed that global warming is the main reason of climate change followed by cutting down of trees (23%). About 22% of inhabitants observed that overpopulation was the third

reason of climate change followed by different pollution (11%). Industrialization was another reason of climate change, and it was perceived by 1% of the respondents of Teesta River basin area. In that area 1% respondents also perceived that climate change occurred naturally (God gifted) (Fig 4).

3.2.3 Threat of climate change

It was observed that ongoing climate change has 78% impact on agriculture followed by health of peoples and other animals that is 20% (Fig 5). Impact of climate was 2% on biodiversity quality and sustainability in Teesta River basin char land. According to the perception of peoples, there was no impact of climate change on transportation communication, business, and instigating disaster. Similar results observed by Haque et al., [14].

3.3 Climate Change Pattern according to the Perception of Char Inhabitants

According to perception of Teesta River basin char people data was collected on three seasons for climate change pattern. In the rainy season they observed increase in lightning from sun as the highest effect of climate change that has a WAI of 1.77. Teesta river basin people perceived dry spell frequency as the second highest extent of climate change that had a WAI of 1.74 followed by timing of rain offset, uneven distribution of rainfall and timing of rain onset (Table 7).



Fig. 3. Knowledge about climate change





Fig. 4. Reasons of climate change



Fig. 5. Threat of climate change

The highest extent of climate change was found in the minimum temperature in winter season and it had highest WAI of 1.72. The second highest extent of climate change was perceived as the late start of winter followed by number of cool days, chilling injury in crop, cold intensity, ending of winter, winter duration and maximum temperature. The lowest extent of climate change was perceived as winter rainy days with WAI of 1.02.

In summer season, the highest extent of climate change was identified as maximum temperature with WAI of 1.82. The second highest extent of change was observed as duration of summer season was increased and it had WAI of 1.43 followed by hailstorm, ending of summer, starting of summer, intensity of hot days, summer rainy days. The lowest extent of climate change was found as minimum temperature in summer season with WAI 0.98.

3.4 Impact of Climate Change on Livelihood According to Teesta River Basin Char Land Inhabitants' Perception of Previous 10 years

There are five capitals of livelihood. These are: human capital, natural capital, financial capital,

physical capital, and social capital. Teesta River basin char respondents have long experience about climate change and weather. Since they are permanent residents, they were enquired about the impact of climate change on their livelihood assets. Perceptions of the char respondents regarding impact of climate change were found as per following.

Table 7. Climate	e Change	pattern	according	to the	perception	of char	[·] inhabitants
------------------	----------	---------	-----------	--------	------------	---------	--------------------------

Components of	Extent of	change					WAI	Rank
climate change	No chang	е	Increased	Decre	ased			order
	HH	%						
			HH	%			_	
					HH	%		
Rainy Season								
Lightening	13	3.4	307	79.9	64	16.7	1.77	1
Dry spell	13	3.4	296	77.1	75	19.5	1.74	2
frequency								
Timing of rain	16	4.2	294	76.6	74	19.3	1.72	3
offset								
Uneven	27	7.0	248	64.6	109	28.4	1.58	4
distribution of								
rainfall								
Timing of rain	21	5.5	130	33.9	233	60.7	1.28	5
onset								
Total amount of	20	5.2	121	31.5	243	63.3	1.26	6
precipitation								
Wind speed	30	7.8	119	31.0	235	61.2	1.23	7
Rainy days	18	4.7	85	22.1	281	73.2	1.17	8
frequency								
Season duration	76	19.8	124	32.3	184	47.9	1.13	9
Cloudy weather	33	8.6	55	14.3	296	77.1	1.06	10
Sunshine hours	205	53.4	85	22.1	94	24.5	0.69	11
Winter Season								
Minimum	11	2.9	287	74.7	86	22.4	1.72	1
temperature								
Starting of winter	48	12.5	176	45.8	160	41.7	1.33	2
Number of cool	43	11.2	115	29.9	226	58.9	1.19	3
days								
Chilling injury in	13	3.4	84	21.9	287	74.7	1.18	4
_ crop	_							
Cold intensity	5	1.3	56	14.6	323	84.1	1.13	5
Ending of winter	1/	4.4	41	10.7	326	84.9	1.06	6
Winter duration	26	6.8	48	12.5	310	80.7	1.06	/
Maximum	63	16.4	84	21.9	237	61.7	1.05	8
temperature	_						1.00	
Winter rainy days	8	2.1	15	3.9	361	94.0	1.02	9
Summer Season							1.00	
Maximum	8	2.1	321	83.6	55	14.3	1.82	1
temperature		05.5	005		0.1		4.40	-
Duration of	98	25.5	265	69.0	21	5.5	1.43	2
season	0	0.0	400	00	007	04.7	4.04	0
Hailstorm	9	2.3	138	36	237	61.7	1.34	3
Ending of	131	34.1	215	56.0	38	9.9	1.22	4
Summer	454	00.0	000	F 4 4	0.1	0.0	4.45	-
Starting of	151	39.3	209	54.4	24	0.3	1.15	5
Summer	6	1.6	50	10 E	200	04.0	1 4 0	6
intensity of hot	o	0.1	52	13.5	320	84.9	1.12	ю
Lummor roinu	22	E 7	40	10.4	200	02.0	1.05	7
Summer rainy	22	5. <i>1</i>	40	10.4	322	03.9	1.05	1
<u>uays</u> Minimum	30	10.2	33	86	310	Q1 2	0.09	8
temperatura	39	10.2	55	0.0	312	01.5	0.90	0
lemperature								

3.4.1 Human capital

The knowledge, skills, and health that people invest in and accumulate during their lives make up human capital. Teesta River basin char peoples were asked about impact of climate change on their human capital. According to their perception it was found that great extent of loss was found in skills of char peoples (48.35%) followed by knowledge of char peoples (44.27%). It was also observed that a third great extent of loss was found in char peoples good health (41.67%) followed by ability to work (39.06%) (Table 8). Similar results observed by Haque et al., [14].

3.4.2 Natural capital

Respondents of char land perceived that among all the natural capital assets, climate change has a great impact on soil. Out of all the respondents 166 (43.2%) observed that soil is becoming dry day by day due to climate change followed by rivers becoming dry (37.8%) (Table 8). They found that growing trees have become difficult (45.1%), due to drought underground water is decreasing (61.7%) and air became dry due to climate change, and these are small extent of loss. They were also asked about sunshine hour reducing and they perceived it as nil that means sunshine hour is increasing day by day.

3.4.3 Financial capital

Respondents were asked about the impact of climate change on financial capital. They mentioned that among all the financial capital due to climate change great extent of loss were flash flood damage crop (49.7%), floods reduce crop yield (39.6%), the people migrated to another sources of income (30.5%). A moderate extent of loss was observed, in decreasing of fish production due to lack of pond and river water (62.8%), pressure on people's deposit (36.5%) and cyclone, tornado causes loss of crops fruits, livestock (43.2%). On the other hand, small extent of loss occured like depending dependence on loan on taking loan from GO and NGO sources (37.8%) and livestock loss due to drought (71%) (Table 8).

3.4.4 Physical capital

Teesta River basin char inhabitants observed that climate change impact was found in all physical capital and great extent impact was perceived on people's migration to another place (41.4%) (Table 8). The moderate extent of loss was perceived by 39.6% of the char people and it causes loss in health facilities and loss of property (shops, land, brick field, small cottage industries). 45.1%, 71.9% and 77.6% people perceived that small extent of loss was found due to climate change in cyclone and flood damage home, loss of uses of technologies and health hazard of family members respectively.

3.4.5 Social capital

Climate change impact was observed in social capital. Teesta River basin char land inhabitants were asked about to extend of climate change impact on social capital. 37.8%, 58.6% and 49.5% char inhabitants perceived that small extent of loss occur in social conflict, social community activities and education of family members respectively. There was no loss occur in drop out of children from school (56.5%) due to climate change (Table 8).

3.5 Strategies for Mitigation of Climate Change in Teesta River Basin Char Land According to Opinion/Perception of Char Inhabitants

Teesta River basin char dwellers opined about 24 adaptation strategies to reduce the impact of climate change on livelihood and ecosystem in that area. Among all the adaptation strategies, control of black smoke from vehicles is the most prioritized adaptation strategy of Teesta River basin char areas with an ASI of 1.90.

Drought is increasing day by day in Teesta River basin char land areas. So, digging more canals got the second position among all the strategies according to the perceptions of the peoples with ASI 1.84 followed by uses of new technologies and ideas in agriculture that means smart agriculture with ASI 1.63. Rainwater harvesting was the 4th adaptation strategy with ASI 1.50 followed by building embankments around the river and canals to avoid flash flood and it was ASI 1.32. The study observed other adaptation strategies index was as like gas emission mitigation from big industries was 1.30, building river dams was 1.21, control over population growth was 1.21, installing deep tube wells was greenhouse effect was 1.10, food 1.14. processing for livestock was 1.08, stopping tree cutting was 1.04, self-resilience was 0.95, assistance from GOs and NGOs was 0.46, more

tree plantation was 0.42, assistance from international donners was 0.40, increasing awareness was 0.32, assistance from the society/community level was 0.30, social forestry was 0.28, training on climate adaptation was

0.19, adopting agroforestry was 0.16, increased educational coverage was 0.16, no adaptation method use was 0.09 and more use of surface water was 0.03 (Table 9). Results has similarity with Rahman et al., [4]. and Haque et al., [14].

Statements	Extent	of loss								
	Great	extent	Mode exter	erate nt	Small	Extent	Nil		WAI	Rank
	HH	%	HH	%	HH	%	HH	%		
Human Capital										
Skill of char	178	46.35	124	32.29	57	14.84	25	6.51	2.18	1
inhabitants										
Knowledge of char	170	44.27	107	27.86	80	20.83	27	26.11	2.09	2
peoples										
Good health	160	41.67	130	33.85	85	22.14	19	4.95	2.07	3
Ability to work	150	39.06	117	30.47	75	19.53	32	8.33	2.05	4
Natural Capital										
Soil become dry	166	43.2	147	38.3	44	11.5	27	7.0	2.18	1
River become dry	145	37.8	106	27.6	122	31.8	11	2.9	2.00	2
Trees growing	81	21.1	99	25.8	173	45.1	31	8.1	1.60	3
become difficult										
Due to drought	54	14.1	55	14.3	237	61.7	38	9.9	1.33	4
Underground										
water decreasing										
Air become dry	71	18.5	10	2.6	276	71.9	27	7.0	1.33	5
Sunshine hrs	8	2.1	19	4.9	190	49.5	167	43.5	0.66	6
reducing										
Financial Capital										
Flash flood	191	49.7	109	28.4	75	19.5	9	2.3	2.26	1
damage crop										
Decreased fish	103	26.8	241	62.8	38	9.9	2	.5	2.16	2
production due to										
lack of pond and										
river water	450	20.0	405	22.0	400	00.0	4	1.0	0.44	2
Floods reduce	152	39.0	125	32.0	103	20.8	4	1.0	2.11	3
 Dreasure on	110	20 6	140	26.5	122	24.4	2	5	1.02	1
doposit	110	20.0	140	30.5	132	34.4	Z	.5	1.95	4
Depends on taking	122	31.8	106	27.6	1/5	37.8	11	20	1.88	5
loan from GO and	122	51.0	100	27.0	145	57.0		2.5	1.00	5
NGO sources										
Cyclone tornado	44	11.5	166	43.2	147	38.3	27	7.0	1.59	6
causes loss of		11.0	100	10.2		00.0		1.0	1.00	Ū
crops fruits.										
livestock										
Migration to	117	30.5	77	20.1	78	20.3	112	29.2	1.52	7
another income										
source										
Livestock loss due	10	2.6	71	18.5	276	71.9	27	7.0	1.17	8
to drought										
Physical Capital										
Health facilities	95	24.7	152	39.6	104	27.1	33	8.6	1.80	1
Migration to	159	41.4	68	17.7	34	8.9	123	32.0	1.68	2
another place										
Cyclone and flood	81	21.1	99	25.8	173	45.1	31	8.1	1.60	3
damage home										
Loss of property	33	8.6	152	39.6	104	27.1	95	24.7	1.32	4
(shops, land, brick										

Table 8. Impact of climate change on livelihood of the respondents

Parvin et al.; Int. J. Environ. Clim. C	Change, vol. 14, no. 2, pp.	407-422, 2024; Article no.IJECC.112900
---	-----------------------------	--

Statements	Exten	t of loss								
	Great extent		Mode	Moderate		Small Extent		Nil		
			exter	extent					WAI	Rank
	HH	%	HH	%	HH	%	HH	%		
field, small cottage industries)										
Uses of technologies	10	2.6	71	18.5	276	71.9	27	7.0	1.17	5
Health hazard of family members	10	2.6	51	13.3	298	77.6	25	6.5	1.12	6
Social Capital										
Social conflicts	122	31.8	106	27.6	145	37.8	11	2.9	1.88	1
Social community activities	13	3.4	15	3.9	225	58.6	131	34.1	0.77	2
Education of family members disturbed	8	2.1	19	4.9	190	49.5	167	43.5	0.66	3
Increased drop out of children from school	16	4.2	42	10.9	109	28.4	217	56.5	0.63	4

Table 9. Strategies for mitigation to Climate change in Teesta River basin char land

Strategies	Not applicable		Name of intensification							
Ū			Highly intensified		Moderately intensified		No intensification		ASI	Dank
	НН	%	HH	%	НН	%	НН	%	-	Nalik
Control of black smoke from vehicles	98	25.5	193	50.3	59	15.4	34	8.9	1.90	1
Digging canals	115	29.9	179	46.6	81	21.1	9	2.3	1.84	2
Uses of new technologies and idea in agriculture (smart agriculture)	152	39.6	175	45.6	43	11.2	14	3.6	1.63	3
Rainwater harvesting	144	37.5	131	34.1	74	19.3	35	9.1	1.50	4
Building embankments	202	52.6	147	38.3	29	7.6	6	1.6	1.32	5
Gas emission mitigation from big industries	129	33.6	87	22.7	70	18.2	98	25.5	1.30	6
Building river dams	217	56.5	137	35.7	22	5.7	8	2.1	1.21	7
Control over population growth	175	45.6	108	28.1	17	4.4	84	21.9	1.15	8
Installing deep tube wells	220	57.3	115	29.9	42	10.9	7	1.8	1.14	9
Greenhouse effect	94	24.5	34	8.9	63	16.4	193	50.3	1.10	10
Food processing for livestock	236	61.5	133	34.6	2	.5	13	3.4	1.08	11
Stopping tree cutting	241	62.8	125	32.6	6	1.6	12	3.1	1.04	12
Self-resilience	57	14.9	17	4.4	2	0.6	308	80.2	0.95	13
Assistance from GOs and NGOs	312	81.3	45	11.7	13	3.4	14	3.6	0.46	14
More tree plantations	323	84.1	50	13.0	2	.5	9	2.3	0.42	15
Assistance from international donors	331	86.2	49	12.8	1	.3	3	.8	0.40	16
Increasing awareness	340	88.5	39	10.2	1	.3	4	1.0	0.32	17
Assistance from the society/community level	344	89.6	37	9.6	1	.3	2	.5	0.30	18

Strategies	Not applicable		Name of intensification							
-			Highly intensified		Moderately intensified		No intensification		ASI	Rank
	HH	%	HH	%	HH	%	HH	%	-	
Social forestry	343	89.3	32	8.3	2	.5	7	1.8	0.28	19
Training on climate adaptation	356	92.7	17	4.4	10	2.6	1	.3	0.19	20
Adopting agroforestry	357	93.0	16	4.2	1	.3	10	2.6	0.16	21
Increased educational coverage	356	92.7	17	4.4	1	.3	10	2.6	0.16	22
No adaptation method used	367	95.6	9	2.3	1	.3	7	1.8	0.09	23
More use of surface water	379	98.7	3	.8	1	.3	1	.3	0.03	24

4. CONCLUSION

The research study was conducted in the Union of Lakhitari in Gangachara upazila and Union of Topa Madhupur in Kaunia Upazila of the Rangpur district and in the Union of Kha Khaqibari of the Dimla Upazila and Union of Gulmonda in Jaldhaka Upazila in the Nilphamari district. Data was collected with the help of a pretested questionnaire. A total number of 93% of the respondents clearly grasp that climate change refers to changes in long-term typical weather. They were unsure of the time frame. About 3% of respondents have no idea about climate change that will affect the average weather during the next 20 to 30 years. It was observed that ongoing climate change has 78% impact on agriculture followed by health of peoples and other animals that is 20%. In the rainy season they observed increase in lightning from sun as the highest effect of climate change that has a WAI of 1.77. Teesta river basin people perceived dry spell frequency as the second highest extent of climate change that had a WAI of 1.74 followed by timing of rain offset, uneven distribution of rainfall and timing of rain onset. According to their perception it was found that great extent of loss was found in skills of char peoples (48.35%) followed by knowledge of char peoples (44.27%). It was also observed that a third great extent of loss was found in char peoples good health (41.67%) followed by ability to work (39.06%). Out of all the respondents 166 (43.2%) observed that soil is becoming dry day by day due to climate change followed by rivers becoming dry (37.8%) (Table 8). They found that growing trees have become difficult (45.1%), due to drought underground water is decreasing (61.7%) and air became dry due to climate change, and these are small extent of loss. They mentioned that among all the financial capital due to climate change great extent of loss were

flash flood damage crop (49.7%), floods reduce crop yield (39.6%), the people migrated to another sources of income (30.5%). A moderate extent of loss was observed, in decreasing of fish production due to lack of pond and river water (62.8%), pressure on people's deposit (36.5%) and cyclone, tornado causes loss of crops fruits. livestock (43.2%). On the other hand, small extent of loss occured like depending dependence on loan on taking loan from GO and NGO sources (37.8%) and livestock loss due to drought (71%). Char inhabitants observed that climate change impact was found in all physical capital and great extent impact was perceived on people's migration to another place (41.4%) (Table 8). The moderate extent of loss was perceived by 39.6% of the char people and it causes loss in health facilities and loss of property (shops, land, brick field, small cottage industries). 45.1%, 71.9% and 77.6% people perceived that small extent of loss was found due to climate change in cyclone and flood damage home, loss of uses of technologies and health hazard of family members respectively. About 37.8%, 58.6% and 49.5% char inhabitants perceived that small extent of loss occur in social conflict. social community activities and education of family members respectively. There was no loss occur in drop out of children from school (56.5%) due to climate change. Teesta River basin char dwellers opined about 24 adaptation strategies to reduce the impact of climate change on livelihood and ecosystem in that area. Among all the adaptation strategies, control of black smoke from vehicles is the most prioritized adaptation strategy of Teesta River basin char areas with an ASI of 1.90.

5. RECOMMENDATIONS

In view of the importance of respondent's involvement in climate change, it is

recommended that all efforts should make to increase the motivation of farmers towards the adoption for climate technology change adaptation in a proportionate way. Experts GO and NGO representatives in collaboration with the farmers might to have launching intensive providina motivation campaign and by technological support in form of training, supply of required inputs in this regard. Suitable combination of technologies is necessary for climate change adaptation that can ensure food production with high quality. Extension workers should have to keep this in mind and put-forth greater effort to convince farmers about the benefits of different technologies used for climate change adaptation. Research should also be undertaken to identify factors and technologies which can improve the livelihood status of the char inhabitants.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

- Un CA, Rodríguez A. Local and global knowledge complementarity: R&D collaborations and innovation of foreign and domestic firms. Journal of International Management. 2018;24(2):137-152.
- Atta-ur-Rahman AS, Parvin GA, Shaw R. Impact of Urban Expansion on Farmlands: A Silent Disaster 7. Urban disasters and resilience in Asia. 2016;91.
- Muhammad N, Mondol MAS, Hasan MF. Effectiveness of union information and service center in utilization of farm information. International Journal of Agricultural Extension. 2015;3(1):37-45.
- Rahman MS, Mony MA, Mondol MAS, Amin MHA, Mohammad N. Need assessment for capacity building of rural women in practicing post harvest activities of potato. Journal of Science and Technology. 2017;44:50.
- 5. Huq S, Ayers J. Taking steps: mainstreaming national adaptation; 2008.
- 6. IPCC. Climate change 2014 synthesis report. IPCC: Geneva, Szwitzerland. 2014;1059-1072.
- 7. Khatun A, Khatun R, Muhammad N, Uddin, MY, Talukder MAI. effect of flooding on human life and environment in sirajganj

district of bangladesh. Journal of Science and Technology. 2015;93:97.

- Mahmood R, Pielke Sr RA, Hubbard K. G., Niyogi D, Dirmeyer PA, McAlpine C, Fall, S. Land cover changes and their biogeophysical effects on climate. International journal of climatology. 2014;34(4):929-953.
- 9. Paul B, Rashid H. Climatic hazards in coastal Bangladesh: non-structural and structural solutions. Butterworth-Heinemann; 2016.
- 10. Harmeling S, Eckstein D. Global climate risk index; 2013.
- 11. Baldacchino G. Islands, island studies, island studies journal; 2006.
- 12. Paul S, Islam MR. Ultra-poor char people's rights to development and accessibility to public services: A case of Bangladesh. Habitat International. 2015; 48:113-121.
- 13. Zaman MQ. Social structure and process in char land settlement in the Brahmaputra-Jamuna floodplain Man. 1991;673-690.
- 14. Haque ME, Islam MN, Alam MJ, Uddin M Y, Haque MM, Islam MR, Turin MZ. Determining the Coping Strategies towards Household Food Security Practiced by the Farmers in Flood Prone Areas. Asian Research Journal of Agriculture. 2021; 14(4):76-81.
- Huda S, Kabir MR, Tabassum S, Sonet M RAF, Muhammad N. Information Needs of the Female Farmers in Agricultural Activities. Asian Journal of Agricultural Extension, Economics & Sociology. 2019; 30(2):1-11.
- Amin MR, Huda S, Karim MR, Molla AM, Muhammad N, Noman MRF. Effect of microcredit on social empowerment of the asa women beneficiaries. Indian Journal of Social Research. 2016;57(5):637-652.
- Cochran WG. Some effects of errors of measurement on multiple correlation. Journal of the American Statistical Association. 1970;65(329):22-34.
- Karim MR, Ampony D, Muhammad N, Noman MRF, Rahman MS. Factors affecting adopting of local adaptation options to climate change vulnerability. Journal of Science and Technology. 2017;15:1-8.
- Huda S, Hasan MR, Rahman MS, Noman MRF, Muhammad N. Effect of Mobile Phone Technology in Improving Small

Farm Productivity. Advances in Computer Science and Information Technology. 2017;4(3):193-19. sustainable development. International Journal of Development Research. 2018; 8(2):18902-18906.

20. Sultana R, Muhammad N, Zakaria AKM. Role of indigenous knowledge in

© 2024 Parvin et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (http://creativecommons.org/licenses/by/4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history: The peer review history for this paper can be accessed here: https://www.sdiarticle5.com/review-history/112900