



A Comparative Diagnosis of Urinary Schistosomiasis in Patients from Primary Healthcare Centers in Keffi, North Central Nigeria, Using Microscopy and PCR

Rosemary Edward ^a, Isa H. Mohammed ^{a*}, Ismaila Ibrahim ^b,
Bawonda E. Omenyi ^c, David Ishaleku ^a, Catherine E. Okoye ^d,
Rita M. Ezeadua ^d and Achimugu Isaiah ^e

^a Department of Microbiology, Nasarawa State University, P.M.B. 1022, Keffi, Nasarawa State, Nigeria.

^b Molecular Biology Unit, Medical Laboratory Department, Federal Medical Centre, P.M.B. 004, Keffi, Nasarawa State, Nigeria.

^c Department of Microbiology and Parasitology, University of Uyo, P.M.B. 1017, Uyo, Akwa Ibom State, Nigeria.

^d Department of Microbiology, Nnamdi Azikwe University, P.M.B. 5025, Awka, Anambra State, Nigeria.

^e Department of Microbiology, Medical Laboratory Services, Federal Medical Centre, P.M.B. 004, Keffi, Nasarawa State, Nigeria.

Authors' contributions

This work was carried out in collaboration among all authors. Authors RE and IHM designed the study, collected samples, performed laboratory and statistical analyses. Authors CEO, RME and AI wrote the first draft of the manuscript. Authors DI, BEO and II conceptualised the study, conducted literature searches, wrote the protocols and led the analyses of the study. All authors read and approved the final manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/113649>

Original Research Article

Received: 27/12/2023

Accepted: 01/03/2024

Published: 22/03/2024

*Corresponding author: E-mail: harunaisamohammed@nsuk.edu.ng, bretp20@yahoo.com;

ABSTRACT

Aims: This study was conducted to compare the diagnosis of urinary schistosomiasis by microscopy and PCR in patients in some selected primary healthcare centres in Keffi, North Central Nigeria.

Study Design: The study was a cross sectional study.

Place and Duration of Study: Keffi, Nasarawa State, between March 2021 and September 2021.

Methodology: Urine samples were collected from 200 patients (29 samples from 29 patients, each from Gidan Zakara, Sabon Gida, Jigwada, Angwan Jaba and 28 samples from 28 patients, each from Kowa, Kaibo Mada and Yarkadai PHCs). Patients' demographic information was obtained using structured questionnaires. The ova of *S. haematobium* were detected microscopically in the samples using the standard sedimentation technique. The DNA of *S. haematobium* was extracted from the samples, amplified and detected by conventional PCR technique using type-specific primers and analysed in a 1.5% agarose gel stained with ethidium bromide. The collected data were analysed using Smith's Statistical Package (version 2.8, California, USA), with a P value of ≤ 0.05 considered statistically significant, and Cohen's Kappa (κ) used to assess the level of agreement between the two diagnostic methods.

Results: Out of the 200 samples screened, 86 (43.0%) were males while the remaining 114 (57.0%) were females. Of them, 6 (3.0%) and 15 (7.5%) were positive for urinary schistosomiasis by microscopy and PCR respectively indicating fair agreement between the two diagnostic methods ($\kappa=0.3142$). There was significant association between gender and urinary schistosomiasis ($P<0.05$), as more males were infected (microscopy: 5.8%, PCR: 15.1%) than their female counterparts (microscopy: 0.9%, PCR: 1.8%). Although both age and occupation of patients were not significantly associated with the infection ($P>0.05$), pupil/students (microscopy: 9.1%, PCR: 29.5%) aged ≤ 14 years (microscopy: 8.0%, PCR: 18.0%) were mostly infected.

Conclusion: Our results show variations between the two diagnostic techniques, with PCR detecting higher rates of urinary schistosomiasis confirming it to be a more sensitive method for diagnosing the infection. Although the findings suggest fair agreement between the two methods, continued efforts to improve diagnostic accuracy are essential for effective disease management.

Keywords: Schistosomiasis; microscopy; PCR; Keffi; Nigeria.

1. INTRODUCTION

Urinary schistosomiasis, caused by the parasitic flatworm *Schistosoma haematobium*, is a pervasive global public health issue most commonly found in South America, Asia and many parts of Sub-Saharan Africa, including Nigeria where an estimated 101.3 million people are at risk of infection and 29 million of the people being infected [1,2,3]. This Neglected Tropical Disease (NTD) continues to pose a substantial burden on the health and well-being of communities, particularly in regions with inadequate water and sanitation infrastructure, where individuals are frequently exposed to freshwater harbouring the larvae of *S. haematobium* [4,5]. Keffi, located in north-central Nigeria, where reliance on untreated water sources is common, is one such region grappling with the challenges posed by urinary schistosomiasis [1,6].

Most Primary Healthcare Centres (PHCs) in Nigeria, like those found in Keffi, Nasarawa

State, where schistosomiasis is prevalent, cannot perform standard laboratory diagnosis [1,7]. This is mainly due to poor funding, limited resources, inadequate training, and infrastructure constraints that lead to reliance on presumptive tests that are uncertain in accuracy and lack guaranteed sensitivity and specificity [8,9]. Consequently, this challenges the effectiveness of measures for preventing and controlling schistosomiasis [3].

The diagnosis of urinary schistosomiasis involves a comprehensive approach, utilizing both traditional microscopic examination and advanced molecular techniques such as polymerase chain reaction (PCR). Clinical evaluation plays a key role in identifying symptoms associated with the disease, including haematuria and dysuria [10]. However, due to the overlap of symptoms with other conditions, a definitive diagnosis relies on laboratory methods [11]. Microscopy, the conventional diagnostic tool for schistosomiasis, has been the cornerstone of disease detection for decades. However, its

limitations in sensitivity, especially for low-intensity infections, and the inability to provide species-specific identification highlight the need for more advanced diagnostic methods [12]. On the other hand, PCR, a molecular technique, offers a more sensitive and specific approach by detecting the DNA of the parasite [10,12]. Despite its higher accuracy, the practicality and cost-effectiveness of PCR in resource-limited settings need careful consideration.

Our study design involves the collection of urine samples from patients attending selected primary healthcare centres in Keffi, followed by a comparative analysis of diagnostic outcomes using both microscopy and PCR. By evaluating the performance of these methods against each other, we aim to highlight their respective strengths and weaknesses in terms of sensitivity. Additionally, socio-demographic and clinical data were gathered to explore potential associations between the prevalence of urinary schistosomiasis and factors such as age, gender, and occupation of the participants.

The findings of this research hold substantial promise for informing healthcare policies, guiding clinicians in their diagnostic choices, and contributing to the global effort to control and eliminate urinary schistosomiasis. By elucidating the comparative diagnostic accuracy of microscopy and PCR in a real-world setting, this study endeavours to facilitate evidence-based decision-making and foster advancements in the management of urinary schistosomiasis in Keffi, North Central Nigeria, and similar endemic regions worldwide.

2. MATERIALS AND METHODS

2.1 Study Area

The study was conducted in seven (7) selected PHCs (Gidan Zakara, Sabon Gida, Jikwada, Yarkaddai, Angwan Jaba, Kawo, and Kaibo Mada) in Nasarawa State, Nigeria. Keffi is a town in north-central Nigeria, about 68 kilometres from Abuja, the Federal Capital Territory, and 128 kilometres from Lafia, the capital of Nasarawa State. It is 850 metres above sea level and is located between latitude 8° 49' 24" N of the equator and longitude 7° 52' 35" E. The majority of the inhabitants are traders, farmers, students and those who engage in menial jobs [13].

2.2 Study Population

Male and female patients of all ages attending the seven (7) designated PHCs in Keffi for

treatment who agreed to participate in the study comprise the study population. These PHCs were chosen because of their proximity to the stream, where locals go to swim, wash, and undertake other household chores, potentially exposing them to risky water contact due to the infected freshwater snails that host the parasite.

2.3 Sample Size Determination

The sample size in this study was obtained using the formula developed by Pourhoseingholi et al. [14] for sample size calculation in medical studies at the 0.05 level of precision.

2.4 Data Collection, Sample Collection and Processing

A self-administered questionnaire was used to collect the participants' socio-demographic information. Following the distribution of the questionnaire, every participant was given a 30 ml sterile wide-mouth screw-capped plastic container labelled with their identification number and informed on how to collect the urine sample aseptically in private [11]. Between 10:00 a.m. and 2:00 p.m., 200 urine samples were collected (29 from Gidan Zakara, Sabon Gida, Jigwada, Angwan Jaba, and 28 from Kowa, Kaibo Mada, and Yarkadai) and were transferred in a cool box to the Federal Medical Centre, Keffi, and stored at 4°C in the refrigerator until ready for analysis.

2.5 Laboratory Analysis

2.5.1 Macroscopic examination urine samples

All urine samples collected were examined macroscopically for colour, clarity, presence and absence of blood, odour, and specific gravity as previously described by Cheesbrough [11].

2.5.2 Microscopy of urine samples

Direct microscopy using the urine sedimentation method was used to detect *S. haematobium* eggs in the collected samples as previously described [11]. For this purpose, 10 ml of urine was poured into a centrifuge tube and spun at 15,000 rpm for 5 minutes. The supernatant was then discarded, and the sediments were collected using a Weide pipette. Two drops of the sediments were placed on a frosted glass slide and covered with a coverslip. The slides were then examined with a light microscope at x10 and x40 objectives for the presence of *S. haematobium* eggs. Eggs were identified and

distinguished from other schistosomes by the presence of a conspicuous lateral spine; they were counted, and infection intensity was recorded as the number of ova/10 mL urine and categorised into light (≤ 50 ova/10 mL urine) and heavy (≥ 50 ova/10 mL urine) [15].

2.5.3 Genomic DNA extraction

The genomic DNA of *S. haematobium* was extracted and purified directly from urine samples using a commercial genomic DNA extraction kit, Qiagen QIAamp DNA Miniprep Kit (Qiagen Group, Germany), according to the manufacturer's instructions.

2.5.4 DNA amplification and polymerase chain reaction (PCR)

S. haematobium DNA was detected by conventional PCR technique previously described by Tembo et al. [16] using specific primers (ShDra consensus primers) from the work of Lodh et al. [17]. In brief, the extracted DNA was amplified on a programmable thermal cycler (MJ Research Inc., Water- town, USA) using a set of primers, namely ShDra1F (5'-GATCTCACCTATCAGACGAAAC-3') and ShDra1R (5'-TCACAACGATACGACCAAC-3') targeting the ShDra region of *S. haematobium*, according to a previously described PCR protocol [16]. The PCR reaction volume was 20 μ L and consisted of 5 ng/ μ L of DNA template, 3 μ L of nuclease-free water, 10 μ L of One Taq quick load (Biolabs, Durham, North Carolina, USA) and 0.5 μ M of each primer. The following PCR conditions were used: 95 °C for 5 min initial denaturation, followed by 40 cycles of 30 s at 95 °C, 30 s at 56 °C, 1 min at 72 °C and a final extension step of 7 min at 72 °C. The analysis was carried out at the research laboratory of the Nigeria Centre for Disease Control, Gaduwa, Abuja-Nigeria.

2.5.5 Agarose gel electrophoresis

The PCR products were analysed in a 1.5% agarose gel stained with ethidium bromide. The size of the PCR products was estimated from the migration pattern of a 100bp to 1000bp increment plus DNA molecular marker (BIONEER Daejeon, North Korea). The expected amplicon band size of the *S. haematobium* ShDra gene was 298 bp.

2.6 Data Analysis

Data collected were analysed using Smith's Statistical Package (version 2.8, California, USA), with a P value of ≤ 0.05 considered

statistically significant. Cohen's Kappa (κ) was used to assess the level of agreement between the two diagnostic methods, with a κ value of ≤ 0 indicating no agreement, 0.01-0.20 indicating slight agreement, 0.21-0.40 indicating fair agreement, 0.41-0.60 indicating moderate agreement, 0.61-0.80 indicating substantial agreement, and 0.81-1.00 indicating almost perfect agreement.

3. RESULTS AND DISCUSSION

This present study compared the diagnosis of urinary schistosomiasis by microscopy and PCR in patients in some selected primary healthcare centres in Keffi. A total of 200 patients most of them female (114/200), civil servants (87/200) and aged 15 to 34 years (84/200) were recruited and their urine samples analysed for the presence of *S. haematobium* using microscopy and PCR (Table 1).

The results showed that 6 (3.0%) of the participants tested positive by microscopy and 194 (97.0%) tested negative for urinary schistosomiasis, while 15 (7.5%) samples tested positive by PCR and 185 (92.5%) tested negative (Table 2). Application of Cohen's Kappa coefficient to assess the degree of agreement between the two diagnostic methods yielded a value of 0.3142, indicating fair agreement beyond chance (Table 2).

Concerning the sampling area, using microscopy, 4 (13.7%) patients from PHC Sabon Gida were positive for urinary schistosomiasis while the remaining 2 (6.9%) patients were from PHC Angwan Jaba, none was from PHC Gidan Zakara, PHC Jigwada, PHC kawo, PHC Kaibo Mada and PHC Yarkadai. When using PCR, 6 (20.7%) patients from PHC Gidan Zakara, 5 (17.2%) from PHC Sabon Gida, 2 (6.9%) from PHC Angwan Jaba and 1 (3.6%) each from PHC Kawo and PHC Kaibo Mada were positive for urinary schistosomiasis while none was from PHC Jigwada and PHC Yarkadai. (Table 3).

To date, microscopy is the conventional method for diagnosing schistosomiasis [3]. However, there is a high probability of false-negative results, especially if the parasite intensity is low or if it is not performed by an expert [10,12]. This is confirmed by the results of this current study, as the microscopy method gave a lower number of positive cases (3.0%) compared to PCR (7.5%) (Figs. 1, 2 and 3). Interestingly, there were no false-negative results by microscopy

and PCR also did not produce any false-positive results. The calculated Kappa coefficient suggests fair agreement but also indicates room for improvement in diagnostic concordance (Table 2).

These results are consistent with previous studies comparing diagnostic methods for urinary schistosomiasis. For instance, Sow et al. [18] reported varying rates of urogenital

schistosomiasis, as real-time PCR detected more cases of the infection (34.6%) compared to microscopy (20.3%) in pre-school, school-aged children and women of productive age in central Senegal. In another study among children and adults in Zanzibar, qPCR also significantly increased the sensitivity of urinary schistosomiasis diagnosis from 13.3 % to 26.8 % compared to urine filtration microscopy [19].

Table 1. Prevalence and distribution of *S. haematobium* infection in relation to Socio-demographics among patients in some selected primary healthcare centres in Keffi, North Central Nigeria by microscopy and PCR

Socio-demographic	No. Examined (N=200)	Prevalence (%)	
		Microscopy	PCR
Age (Years)			
≤ 14	50	4(8.0)	9(18.0)
15 – 34	84	1(1.2)	3(3.6)
35 – 64	43	1(2.3)	3(6.9)
≥ 65	23	0(0.0)	0(0.0)
P-value		0.1107	0.0850
Gender			
Male	86	5(5.8)	13(15.1)
Female	114	1(0.9)	2(1.8)
P-value		0.0428*	0.0004*
Occupation			
Pupil/Student	44	4(9.1)	13(29.5)
Farmer	87	1(1.1)	2(2.3)
Civil Servant	41	0(0.0)	0(0.0)
Others	28	1(3.6)	0(0.0)
P-value		0.0774	0.1000

*Statistically significant

Table 2. Comparative diagnosis of urinary schistosomiasis by microscopy and PCR in patients in some selected primary healthcare centres in Keffi, North Central Nigeria

Diagnostic method	No. examined	Outcome	
		No. positive(%)	No. negative(%)
Microscopy	200	6(3.0)	194(97.0)
PCR	200	15(7.5)	185(92.5)

Kappa coefficient(κ)= 0.3142

Table 3. Prevalence and distribution of *S. haematobium* infection in relation to Sampling Area among patients in some selected primary healthcare centres in Keffi, North Central Nigeria by microscopy and PCR

Sampling Area	No. Examined (N=200)	Prevalence (%)	
		Microscopy	PCR
PHC Gidan Zakara	29	0(0.0)	6(20.7)
PHC Sabon Gida	29	4(13.7)	5(17.2)
PHC Jigwada	29	0(0.0)	0(0.0)
PHC Angwan Jaba	29	2(6.9)	2(6.9)
PHC Kawo	28	0(0.0)	1(3.6)
PHC Kaibo Mada	28	0(0.0)	1(3.6)
PHC Yarkadai	28	0(0.0)	0(0.0)
Total	200	6(3.0)	15(7.5)




Fig. 1. Agarose gel electrophoretogram of *S. haematobium* ShDra1 amplified gene. Samples 1, 2, 6, 7, 17, 18, and 20 were positive for *S. haematobium* DNA while samples 3, 4, 5, 8, 9, 10, 11, 12, 13, 14, 15, 16 and 19 were negative. *M represents the molecular ladder, '-' is the negative control while '+' is the positive control




Fig. 2. Agarose gel electrophoretogram of *S. haematobium* ShDra1 amplified gene. Samples 21, 24, 25 and 31 were positive for *S. haematobium* DNA while samples 22, 23, 26, 27, 28, 29, 30, 32, 33, 34, 35, 36, 37, 38, 39 and 40 were negative. *M represents the molecular ladder, '-' is the negative control while '+' is the positive control




Fig. 3. Agarose gel electrophoretogram of *S. haematobium* ShDra1 amplified gene. Samples 44, 49, 52, 54 and 59 were positive for *S. haematobium* DNA while samples 41, 42, 43, 45, 46, 47, 48, 50, 51, 53, 55, 56, 57, 58 and 60 were negative. *M represents the molecular ladder, ‘-’ is the negative control while ‘+’ is the positive control

The higher infection rates detected with the PCR technique indicate that this technique is more accurate and robust in detecting schistosomiasis even at low levels that might have gone undetected with microscopy.

Interestingly, the overall prevalence of 7.5% of *S. haematobium* infection by PCR recorded in this current study is lower than the reports of most previous researchers who detected the parasite at the same study location using microscopic methods. For example, Ezhim et al. [20] and Haruna Mohammed & Buhari [1] reported 53% and 12.0% prevalence of schistosomiasis in urine of primary school pupils in Keffi, respectively. In addition, Abdullahi & Ramatu [21] also recently reported 15.3% cases of the parasitic infection among pupils of Angwan Lambu primary school in Keffi. The present study has reconfirmed the endemicity of this Neglected Tropical Disease (NTD) in Keffi Nasarawa State, which was previously associated with extreme poverty, inadequate or lack of social amenities such as portable water for drinking and other domestic activities [1, 2, 22].

The decreased rate of urinary schistosomiasis observed in this present study compared to other previous studies conducted in the same locality could be due to the integrated prevention and

control measures introduced in public primary schools in Nigeria, including mass drug administration to infected individuals and health education on the predisposing factors responsible for the transmission of the parasitic infection [23]. It could also be due to the fact that most of the previous studies conducted at the study location such as those by Haruna Mohammed & Buhari [1], Ezhim et al. [20] and Abdullahi & Ramatu [21] were all carried out among school-aged children who are known to be the most susceptible groups to schistosomiasis [2, 22], whereas the current study surveyed patients of all age groups.

There is no significant association between urinary schistosomiasis and the age of the participants in the present study ($P > 0.05$) (Table 1). However, patients aged ≤ 14 years were most frequently infected using both microscopy (8.0%) and PCR (18.0%). This is to be expected as school-aged children (SAC), who are usually under 14 years of age, have been shown to be the most vulnerable group for schistosomiasis [7, 24, 25, 26] as they are more likely to engage in activities that may expose them to contaminated freshwater, like playing in infested water, swimming, washing, etc [4, 27]. Other previous studies also reported similar findings [1, 18, 28].

Infection with *S. haematobium* is significantly associated with gender in this study ($P < 0.05$), as more male patients were infected with the parasites both microscopically (5.8%) and by PCR (15.1%) than their female counterparts (0.9% and 1.8% for microscopy and PCR, respectively) (Table 1). This could be due to the fact that most males, especially in rural areas, engage in outdoor jobs, such as farming and fishing, where they come into contact with freshwater sources, which increases their risk of infection with *S. haematobium*. Most other researchers have also made similar observations [1, 24, 26, 29].

It is interesting to note that most Nigerian farmers live in rural areas associated with extreme poverty, inadequate or lack of social amenities such as portable water for drinking and other domestic activities [2,22], and residents of such areas are usually at risk of many infections, including schistosomiasis [1, 7,30]. Surprisingly, however, the patient's occupation is not associated with urinary schistosomiasis in the present study ($P > 0.05$), and pupils/students were more frequently infected with the parasite using both microscopy (9.1%) and PCR (29.5%) than farmers (1.1% and 2.3% for microscopy and PCR, respectively) (Table 1). This observed result is most likely due to the fact that the schools in the surveyed areas might be located close to water bodies contaminated with schistosome and pupils/students may indulge in play activities involving contact with the water, making them susceptible to schistosomiasis.

4. CONCLUSION

In the present study, PCR demonstrated greater sensitivity with fair agreement compared to microscopy in detecting urinary schistosomiasis. The results suggest that PCR could be a more reliable diagnostic tool for identifying cases of the disease in the population studied. Therefore, further efforts are needed to improve the diagnostic accuracy of the infection to enhance detection and treatment efficacy in North Central Nigeria and beyond.

CONSENT AND ETHICAL APPROVAL

The Keffi Local Government Council Research Ethics Committee granted permission to undertake this study (KLG/WELL/227/VOL.I/XXX). The heads of each PHC where urine samples were taken also gave their approval. Furthermore, all participants in

this study fully completed and signed an informed consent form.

ACKNOWLEDGEMENT

The study team would like to thank the management of the Primary Healthcare Centres, Gidan Zakara, Sabon Gida, Jikwada, Yarkaddai, Angwan Jaba, Kawo and Kaibo Mada for their kind permission to conduct the research. We would also like to thank all the patients or their parents/guardians who voluntarily participated in the study. However, this research was not supported in any way by governmental or non-governmental organisations.

COMPETING INTERESTS

The authors have declared that no competing interests exist.

REFERENCES

1. Haruna Mohammed I, Buhari S. Prevalence and determining factors of urinary schistosomiasis among primary school pupils in North Central Nigeria; 2020;61175.
2. Hotez PJ, Savioli L, Fenwick A. Neglected tropical diseases of the Middle East and North Africa: Review of their prevalence, distribution, and opportunities for control. PLoS Neglected Tropical Diseases. 2012;6(2):e1475. Available: <https://doi.org/10.1371/journal.pntd.0001475>
3. World Health Organization: Schistosomiasis: Progress- Google Scholar. (n.d.). Retrieved August 18, 2023. Available: https://scholar.google.com/scholar_lookup?title=Schistosomiasis:+progress+report+2001-2011+and+strategic+plan+2012-2020&publication_year=2013&
4. Hajissa K, Muhajir AEMA, Eshag HA, Alfadel A, Nahied E, Dahab R, Ali SM, Mohammed M, Gaafar M, Mohamed Z. Prevalence of schistosomiasis and associated risk factors among school children in Um-Asher Area, Khartoum, Sudan. BMC Research Notes. 2018; 11(1): 779. Available: <https://doi.org/10.1186/s13104-018-3871-y>
5. Singh K, Muddasiru D, Singh J. Current status of schistosomiasis in Sokoto,

- Nigeria. Parasite Epidemiology and Control. 2016;1(3):239–244. Available:<https://doi.org/10.1016/j.parepi.2016.08.003>
6. Donuma KU, Ma L, Bu C, George LY. Impact and health risk assessment of groundwater in the vicinity of dumpsites in Keffi Metropolis, Nigeria. *Journal of Geoscience and Environment Protection*. 2023;11(8). Available:<https://doi.org/10.4236/gep.2023.118006>
 7. Pam DD, Pam VA, Nuhu EI. Prevalence of schistosoma mansoni infection among primary school pupils in Keffi Town, Keffi local government area, Nasarawa State, Nigeria. *Nigerian Journal of Parasitology*. 2016;37(2). Available:<https://doi.org/10.4314/njpar.v37i2>
 8. Oyekale AS. Assessment of primary health care facilities' service readiness in Nigeria. *BMC Health Services Research*. 2017;17(1):172. Available:<https://doi.org/10.1186/s12913-017-2112-8>
 9. Paul. Factors militating against effective implementation of primary health care (PHC) system in Nigeria. *Annals of Tropical Medicine and Public Health (ATMPH)*; 2023. Available:<https://www.atmph.org/factors-militating/>
 10. Nigo MM, Salieb-Beugelaar GB, Battagay M, Odermatt P, Hunziker P. Schistosomiasis: From established diagnostic assays to emerging micro/nanotechnology-based rapid field testing for clinical management and epidemiology. *Precision Nanomedicine*. 2019;3(1):439–458. Available:[https://doi.org/10.33218/prnano3\(1\).191205.1](https://doi.org/10.33218/prnano3(1).191205.1)
 11. Cheesbrough M. *District Laboratory Practice in Tropical Countries*; 2009.
 12. Utzinger J, Becker SL, Lieshout L van, Dam GJ van, Knopp S. New diagnostic tools in schistosomiasis. *Clinical Microbiology and Infection*. 2015; 21(6): 529–542. Available:<https://doi.org/10.1016/j.cmi.2015.03.014>
 13. Yakubu IO. Geography of Keffi, Nassarawa State, Nigeria; 2013. Retrieved August 20, 2023. Available:https://www.academia.edu/13014831/Geography_of_Keffi_Nassarawa_State_Nigeria_Published_online_in_2013_
 14. Pourhoseingholi MA, Vahedi M, Rahimzadeh M. Sample size calculation in medical studies. *Gastroenterology and Hepatology From Bed to Bench*. 2013;6(1): 14–17.
 15. WHO expert committee. Prevention and control of schistosomiasis and soil-transmitted helminthiasis. *World Health Organization Technical Report Series*. 2002;912(i–vi):1–57.
 16. Tembo R, Muleya W, Yabe J, Kainga H, Nalubamba KS, Zulu M, Mwaba F, Saad SA, Kamwela M, Mukubesa AN, Monde N, Kallu SA, Mbewe N, Phiri AM. Prevalence and molecular identification of schistosoma haematobium among children in Lusaka and Siavonga districts, Zambia. *Tropical Medicine and Infectious Disease*. 2022;7(9). Available:<https://doi.org/10.3390/tropicalmed7090239>
 17. Lodh N, Naples JM, Bosompem KM, Quartey J, Shiff CJ. Detection of parasite-specific dna in urine sediment obtained by filtration differentiates between single and mixed infections of schistosoma mansoni and s. haematobium from endemic areas in Ghana. *PLoS ONE*. 2014;9(3): e91144. Available:<https://doi.org/10.1371/journal.pone.0091144>
 18. Sow D, Sylla K, Dieng NM, Senghor B, Gaye PM, Fall CB, Goumballa N, Diallo A, Ndiaye J. LA, Parola P, Sokhna C, Doucouré S, Faye B. Molecular diagnosis of urogenital schistosomiasis in pre-school children, school-aged children and women of reproductive age at community level in central Senegal. *Parasites & Vectors*. 2023;16(1). Available:<https://doi.org/10.1186/s13071-023-05671-x>
 19. Keller D, Rothen J, Dangy JP, Saner C, Daubenberger C, Allan F, Ame SM, Ali SM, Kabole F, Hattendorf J, Rollinson D, Seyfarth R, Knopp S. Performance of a real-time PCR approach for diagnosing Schistosoma haematobium infections of different intensity in urine samples from Zanzibar. *Infectious Diseases of Poverty*. 2020;9(1):128. Available:<https://doi.org/10.1186/s40249-020-00726-y>
 20. Ezhim M, Oti V, Goriya K, Gloria Z, Dlama J. Prevalence of urinary schistosomiasis

- among primary school children in a Northern Nigerian population. *International Journal of Advanced Research*. 2015;3(11):511 – 519.
21. Abdullahi M, Ramatu MA. Prevalence of urinary schistosomiasis among pupils attending Angwan Lambu Primary School, Keffi, Nasarawa State, Nigeria. *FUDMA Journal of Sciences*. 2022;6(6). Available:<https://doi.org/10.33003/fjs-2022-0606-1134>
22. Uneke CJ, Ugwuoke-Adibuah S, Nwakpu KO, Ngwu BA. An Assessment of schistosoma haematobium infection and urinary tract bacterial infection amongst school children in rural eastern Nigeria. *The Internet Journal of Laboratory Medicine*. 2009; 4(1). Available:<http://ispub.com/IJLM/4/1/3651>
23. Oyeyemi OT. Schistosomiasis control in Nigeria: Moving round the circle? *Annals of Global Health*. 2020;86(1):74. Available:<https://doi.org/10.5334/aogh.2930>
24. Dogara MM, Ahmad S, Balogun BJ, Dawaki SS, Mustapha MB, Abdurrahman AU, Bala L, Zakari A, Livingstone DU. Schistosomiasis and associated risk factors among school-aged children in northern Nigeria. *International Journal of Translational Medical Research and Public Health*. 2020;4(2). Available:<https://doi.org/10.21106/ijtmrph.146>
25. Ojo JA, Adedokun SA, Akindele AA, Olorunfemi AB, Otutu OA, Ojurongbe TA, Thomas BN, Velavan TP, Ojurongbe O. Prevalence of urogenital and intestinal schistosomiasis among school children in South-west Nigeria. *PLOS Neglected Tropical Diseases*. 2021;15(7):e0009628. Available:<https://doi.org/10.1371/journal.pntd.0009628>
26. Sady H, Al-Mekhlafi HM, Mahdy MAK, Lim YAL, Mahmud R, Surin J. Prevalence and associated factors of schistosomiasis among children in Yemen: Implications for an effective control programme. *PLOS Neglected Tropical Diseases*. 2013; 7(8): e2377. Available:<https://doi.org/10.1371/journal.pntd.0002377>
27. Alade T, Ta-Tang TH, Nassar SA, Akindele AA, Capote-Morales R, Omobami TB, Berzosa P. Prevalence of schistosoma haematobium and Intestinal Helminth Infections among Nigerian School Children. *Diagnostics*. 2023;13(4):759. Available:<https://doi.org/10.3390/diagnostics13040759>
28. Esiere RK, Ibeneme EO, Effanga EO, Imalele EE, Esiere MK, Inyang-Etoh PC, Alaribe AAA. Detecting schistosoma haematobium infection by microscopy and polymerase chain reaction (pcr) in school children in three senatorial districts of cross river state, Nigeria. *Journal of Parasitic Diseases: Official Organ of the Indian Society for Parasitology*. 2022;46(1):272–279. Available:<https://doi.org/10.1007/s12639-021-01446-2>
29. Dawaki S, Al-Mekhlafi HM, Ithoi I, Ibrahim J, Abdulsalam AM, Ahmed A, Sady H, Atroosh WM, Al-Areeqi MA, Elyana FN, Nasr NA, Surin J. Prevalence and risk factors of schistosomiasis among Hausa communities in Kano State, Nigeria. *Revista Do Instituto de Medicina Tropical de São Paulo*. 2016;58:54. Available:<https://doi.org/10.1590/S1678-9946201658054>
30. Jibril H, Mohammed Y, Idris A, Manu A, Umar A, Ahmad IT. Incidence of urinary schistosomiasis among rice farmers in some selected villages of kura local government area, Kano, Nigeria. *Nigerian Journal of Experimental and Clinical Biosciences*. 2020;8(2): 65–65.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/113649>