



Millets for Food and Nutritional Security in the Context of Climate Resilient Agriculture: A Review

Lalit Kumar ^a, R. K. Naresh ^b, Himanshu Tiwari ^{b*}, S. K. Kataria ^c,
Sachin Saharan ^d, B. Rajeswara Reddy ^e, Omkar Singh ^f,
Sidra Qidwai ^{g#} and Ram Pratap Singh ^{g†}

^a Krishi Vigyan Kendra Moradabad, Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut, Uttar Pradesh, India.

^b Department of Agronomy, Sardar Vallabhbhai Patel University of Agriculture and Technology, Meerut, Uttar Pradesh, India.

^c Department of Agronomy, Gochar Mahavidyalay Rampur Maniharan, C. C.S. University Meerut, Saharanpur, Uttar Pradesh, India.

^d Department of Agronomy, CSK Himachal Pradesh Agricultural University, Palampur, Himachal Pradesh, India.

^e Agronomy, School of Agricultural Sciences, Nagaland University, Nagaland, India.

^f ICAR-Indian Institute of Farming Systems Research, Modipuram, Meerut-250110, Uttar Pradesh, India.

^g Department of Agronomy, ANDUAT, Kumarganj, Ayodhya- 224 229, Uttar Pradesh, India.

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJPSS/2022/v34i232504

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/92823>

Review Article

Received 20 August 2022
Accepted 27 October 2022
Published 31 October 2022

ABSTRACT

The world is dealing with both agrarian and nutritional issues. We must concentrate on dry lands in order to further increase grain production because agricultural lands with irrigation facilities have been fully utilised. It is difficult to use dry lands to produce enough high-quality grains because of their low fertility. Millets, a crop that complies with climate change regulations, outperform other

^o Senior Research Fellow;

[#] Ph.D. Scholar (Agronomy);

[†] S.M.S. (Agronomy);

*Corresponding author: E-mail: himanshu1996nduat@gmail.com;

grains like wheat and rice in terms of poor growing conditions and high nutritional value. Sustainable food systems aim to provide sufficient and nutritious food, while maximising climate resilience and minimizing resource demands as well as negative environmental impacts. We perform a series of optimizations to maximize nutrient production (i.e., protein and iron), minimize greenhouse gas (GHG) emissions and re-source use (i.e., water and energy), or maximize resilience to climate extremes. We find that increasing the area under coarse cereals (i.e., millets, sorghum) improves nutritional supply (on average, +1% to +5% protein and +5% to +49% iron), increases climate resilience (1% to 13% fewer calories lost during an extreme dry year), and reduces GHGs (-2% to -13%) and demand for irrigation water (-3% to -21%) and energy (-2% to -12%) while maintaining calorie production and cropped area. The extent of these benefits partly depends on the feasibility of switching cropped area from rice to coarse cereals.

Climate-resilient millets are regarded as "Miracle Grains" because of their ability to adapt to a wide range of ecological conditions while using less water for irrigation and producing more effectively in low-nutrient soils. They exhibit little vulnerability to environmental stresses and only minimal demand for artificial fertilisers. Reviving interest in millet groups as nutritious foods that can improve food and nutritional security and reduce malnutrition is necessary. Two main groups of millets are great millets (Sorghum and Pearl millet) and Small millets (Finger millet, Foxtail millet, Little millet, Proso millet, Barnyard millet, Kodo millet and Brown top millet) classified based on the grain size. Both great and small millets have traditionally been the main components of the food basket of the poor people in India. India stands first in area of millets with 90.94 lakh Hectare followed by Niger with 69.99 lakh Hectare. Millets area of the entire world accounts for 312.44 lakh Hectare. India also stands first in production of millets with 115.6 lakh tonnes followed by Niger with 37.9 lakh tonnes. Millets Production of the entire world accounts for 284.59 lakh tonnes. Uzbekistan stands first in yield of millets with 7563 kg per ha followed by Switzerland with 4236 kg per Hectare, yield of the entire world accounts for 910 kg per ha (Food and Agricultural Organization, 2017).

Millets contain high amounts of proteins, fiber, niacin, thiamine and riboflavin, methionine, lecithin and little of vitamin E. They are rich in minerals like iron, magnesium, calcium and potassium also. Due to their nutritional value, millets may help prevent cancer, reduce the risk of cardiovascular disease, stop the growth of tumours, lower blood pressure, lower the rate at which fat is absorbed, delay gastric emptying, and increase gastrointestinal bulk. Value-adding millet grains as ready-to-eat and ready-to-cook foods provides farmers with a good opportunity to increase income generation, promotes production, and fosters marketing, all of which lead to the creation of jobs, income, and nutritional security. However, the successful harvest of small millets justifies the incorporation of tried-and-true and climate-smart technologies for the satisfaction of the population's future needs. The review paper focused on all these aspects. Moreover, the research scope mentioned in the review paper implies future directions for enhancing millet-based agriculture viable in diversifying food baskets and achieving food and nutritional security in a hunger-free society.

Keywords: Millets; food security; nutritional value; climate-resilient crops.

1. INTRODUCTION

"Tackling hunger and feeding the world population are two of the biggest challenges of the modern world. Reasons contributing to this issue range from deficiencies in the supply of micro- and macro-nutrients, shortage in production of foods leading to supply-demand imbalances, and conflicts destabilize various parts of the world. Although several of these triggers for hunger can be addressed leading to a slight reduction in the population suffering from hunger and malnutrition from almost one billion in 1990-1992 to 850 million in 2010-2012, the

threat of climate change and global warming still lingers" [1]. "Estimates show that the reduction in food production rates along with the added pressure of feeding a population exceeding 9 billion by 2050 could lead to 2-3 billion people suffering from hunger, food and nutritional insecurities" [2,3]. Millets are multipurpose: They consume 70 percent less water than rice; grow in half the time of wheat; and require 40 percent less energy in processing. They are one-stop solution in the wake of climate change, water scarcity, and drought conditions along with high nutritive value to provide sustainable food security (Fig.1).

Chart 1. Millets: an approach for sustainable agriculture and healthy world

↓	↓	↓	↓
Food Security	Nutritional Security	Safety from diseases	Economic security
Sustainable food source for combating hunger in changing world climate Resistant to climate stress, pests and diseases	Rich in micronutrients like calcium, iron, zinc, iodine etc. Rich in bioactive compounds Better amino acid profile	Gluten free: a substitute for wheat in celiac diseases Low GI: a good food for diabetic persons Can help to combat cardiovascular diseases, anemia, calcium deficiency etc.	Climate resilient crop Sustainable income source for farmers Low investment needed for production Value addition can lead to economic gains




Fig. 1. Current and optimized shares of monsoon cereal production [Source: 4]

Diets with a low environmental impact are protective of bio diversity and ecosystems, which helps ensure access to food and nutrition. Without sacrificing nutritive value, crop diversification through the use of more coarse cereals, like millets, can increase food production, lower greenhouse gas (GHG) emissions, and improve climate resilience. Millets were identified as a viable option for ensuring food security and environmental resilience in a quantitative analysis of changing monsoon cereal production in India. The Fig. 2 indicates the contribution to calories from monsoon cereals with maximum supply of protein and iron, maximum savings in water, energy, green house gases, and climate resilience.

Millets are a smart food because they are good for your health, the environment because they

require less water to grow and have a low carbon footprint, and the farmer because they are more tolerant of changing weather patterns. Millets are a source of food for more than 90 million people in Asia and Africa. Wheat, rice, and maize, in contrast, are staple foods for 4 billion people. These three major cereals provide 51% of the world’s calorie intake. In the past, millets were poor farmer’s insurances against the vagaries of the Indian monsoon. In the future, millets can be our insurance in times of climate change. Millets are resilient to extreme conditions including high temperatures and drought. They can grow in the harshest, most arid regions. Currently, around 55% of millets are grown in arid regions of Africa, 40% in Asia, and 3% in Europe. In India, the demand for millets has grown by 140% but the production is less than 50%.

A farmer gets just 600 kilograms of millet per hectare with 400 millimeters of rain. Currently, the yield of rice in India is 2.7 metric tons per hectare and wheat is 3.5 metric tons. In China, this is 4.9 and 5.6 metric tons respectively but with investments in input technology, such as high-quality seeds and irrigation, per hectare yield of small millets can reach 4.5 tons, and pearl millet and sorghum can touch 9 tons. Millets have a diverse genetic makeup and require few cultivation inputs. They are also more nutrient-dense than traditional grains [5]. Millets, which are grown for both food and fodder, provide food and livelihood stability to millions of people while also assisting in efficient farming especially for small/marginal farmers of rain-fed areas [6]. Malnutrition is pervasive in many developing nations; for example, India is one of many countries where child malnutrition is severe [7]. Minor millets are essential food group that has been missing from the food basket in recent years. Minor millets are well-known for their health advantages.

Because millets are so nutrient-dense, there is a need to promote them as a reliable source of nutrition. In human history, these have been significant food staples, especially in Asia and Africa. Over the past three decades, consumption of sorghum and other millets as direct food has significantly decreased. The production of millets has significantly decreased in India as a result of the decline in demand. Production of sorghum in India has come down from 7 million tonnes during 2010-11 to 4.2 million tonnes during 2015-16; bajra production was reduced from 10.4 million tonnes to 8.1 million tonnes, production of ragi reduced from 2.2 million tonnes to 1.8 million tonnes and small millets production came down to 0.39 million tonnes from 0.44 million tonnes during the same period.

Davis et al. [4] also found that “the potential for substantial increases in iron supply, modest improvements in protein supply, reductions in water and energy demands and GHG emissions, and enhancements of climate resilience” (Fig. 2) all while maintaining calorie production from cereals within each state. National iron supply from cereals could increase by 49% on average (+737 tons annually). Changes in protein supply were relatively modest in comparison (+5%) due to the overall similarity in protein content between cereals. Reductions in total water demand were also modest (−8%), while monsoon irrigation water demand could be

reduced by 16 km³ H₂O on average (or −21%) and in many places that currently experience declining groundwater tables. Energy demand reduced by an average 12% and GHG emissions were reduced by an average 21 M tonne CO₂eq (−13%). Further, by increasing areas cultivating coarse cereals, the loss of calories under an extremely dry year would reduce by 13% on average.

“These estimated changes are in large part a reflection of the nutritional and environmental characteristics of rice in comparison to coarse grains [i.e., comparable protein content; lower iron content; less efficient intensities of water use energy use, and GHG emissions” [8], and higher climate sensitivity [4].

There is a big shortfall between the amount of food we produce today and the amount needed to feed everyone in 2050. There will be nearly 10 billion people on Earth by 2050—about 3 billion more mouths to feed than there were in 2010.

Feeding 10 billion people *sustainably* by 2050, then, requires closing three gaps:

- **A 56 percent food gap** between crop calories produced in 2010 and those needed in 2050 under “business as usual” growth;
- **A 593-million-hectare land gap** (an area nearly twice the size of India) between global agricultural land area in 2010 and expected agricultural expansion by 2050; and
- **An 11-gigaton GHG mitigation gap** between expected agricultural emissions in 2050 and the target level needed to hold global warming below 2°C (3.6°F), the level necessary for preventing the worst climate impacts.

Millets are a generally excellent source of fibre, minerals, and B-complex vitamins. Millets' high levels of fibre and the presence of certain antinutritional elements like phytates and tannins have an impact on the bioavailability of minerals. “Millets are also rich in health promoting phytochemicals like polyphenols, lignans, phytosterols, phytoestrogens, phytocyanin's. These function as antioxidants, immune modulators, detoxifying agents etc. and hence protect against age-related degenerative diseases like cardiovascular diseases (CVD), diabetes, cancer etc.” (Rao et al. 2011). “Some of the known nutrient's vitamins, minerals,

essential fatty acids also have benefits in terms of prevention of degenerative diseases besides their known functions of preventing nutritional deficiency diseases. Being non-glutinous, millets are safe for people suffering from gluten allergy and celiac disease. They are non-acid forming, easy to digest and non-allergenic” [9]. “Millets have potential for protection against age-onset degenerative diseases. Consumption of millets reduces risk of heart disease, protects from diabetes, improves digestive system, lowers the risk of cancer, detoxifies the body, increases immunity in respiratory health, increases energy levels and improves muscular and neural systems and are protective against several degenerative diseases such as metabolic syndrome and Parkinson’s disease” (Chandrasekara and Shahidi, 2012). “The important nutrients present in millets include resistant starch, oligosaccharides, lipids, antioxidants such as phenolic acids, avenanthramides, flavonoids, lignans and phytosterols which are believed to be responsible for many health benefits” (Edge et al. 2005).

Among these, “millets are known for their climate-resilient features including adaptation to a wide range of ecological conditions, less

irrigational requirements, better growth and productivity in low nutrient input conditions, less reliance on synthetic fertilizers, and minimum vulnerability to environmental stresses. The achieves inclusive and fair growth and development in our country, the second green revolution must focus primarily on nutrition, which was overlooked in the first green revolution, which was focused on production” (Negin et al. 2009). In our country, inner invisible hunger (micronutrient insufficiency) is a major issue. Eradication of extreme poverty and hunger is the first of the Millennium Development Goals (MGDs) proposed by the United Nations in the year 2000. India is far away from achieving this goal [10]. Further, cultivation of millets addresses some of the Sustainable Development Goals (SDG) such as SDG 1 (no poverty), SDG 2 (zero hunger), SDG 3 (good health and wellbeing) and SDG 15 (life on land) (UN, 2021). “Tackling these challenges necessitates a paradigm shift from the existing incremental adaptation strategies toward transformative substitutes that emphasize human health, nutrition, and environmental sustainability. The current natural disasters make it even more imperative to shift toward a climate-resilient agriculture system” [11], (Rasul, 2021).




Fig. 2. Outcomes of optimizations for nutrient supply, environment, and climate resilience [Source: Davis et al. 2019]

2. MILLETS FOR NUTRITIONAL IMPORTANCE

World is in the clinch of several health disorders and chonical diseases. A nutrient imbalanced diet is responsible for most of these diseases. According to the estimates of United Nations Food and Agriculture Organization, about 795 million people (10.9% of world population in 2015) were reported undernourished. While on the other hand more than 1.9 billion (39% of world's population) adults \geq 18 years of age were over -weight and further 13% were reported to be obese [12]. India is the home of world's largest undernourished population. About 194.6 million people, i.e., 15.2% of total population of India, are undernourished. According to the 2017 Global Hunger Index report, India ranked 100th among 119 countries. The score of India is even poorer than Nepal, Sri Lanka and Bangladesh [13]. Protein energy malnutrition (PEM) was reported to result in 4, 69,000 deaths with 84,000 deaths from the deficiency of other vital nutrients such as iron, iodine and vitamin A [14]. Obesity is also a major health concern in India with the prevalence rate of 11% in men and 15% in women.

60 million children under-weight (highest in world), 30% low birth weight babies, 75% pre-school children suffer from iron deficiency anaemia, 85% districts have endemic iodine deficiency [15] and 35.7% of children under five are under weight; 58.4% of children between 6 and 59 months are anaemic; 53% of (non-pregnant) women are anaemic. 51 million people suffer from diabetes which is expected to increase to 79.4 million by 2030 (the increasing consumption of highly polished rice grains and decreasing consumption of coarse cereals contributes to this trend) [16] 18.5% children over weight 5.3% children obese.

Choi et al. [17] studied "the effect of dietary protein of Korean fox-tail millet and found its importance in increasing insulin sensitivity and cholesterol metabolism". Lee et al. [18] investigated "the effect of millet consumption on lipid levels and C-reactive protein concentration; it was found that hyperlipidemic rats fed with foxtail millet had decreased levels of triglycerides". "Aqueous and ethanolic extracts of kodo millet have been reported to produce a dose-dependent fall in fasting blood glucose" [19]. Further millets are gluten free and might have anti-carcinogenic properties [20].

"The millets contain as high as 13-38% of total dietary fiber that could be considered in the management of disorders like diabetes mellitus, obesity, hyperlipidemia, etc. The glycemic load lowering effects of barnyard millet is highest among all the millets" [21]. "Foxtail millet exhibits antihyperglycemic and antilipidemic activities. An aqueous extract of 300 mg of foxtail millet per kg body weight of rats exhibited 70% reduction in blood glucose level in streptozotocin induced diabetic rats" [22]. Millets are also a good source of carotenoids (78-366mg/100g) and possess higher antioxidant capacity [23]. The antioxidant activity of millets is also attributed to their tocopherol content (Fig. 3).

The barnyard millet grain contains about 49-65% carbohydrates [24]. The carbohydrates are mainly composed of dietary fibre. The high content of dietary fibre helps in the prevention of constipation, reducing glycaemic load and lowering of blood cholesterol level. It has the highest content of crude fibre (14.7g/100g) and minerals (4.0g/100g) among all the millets. The iron content (18.6 mg/100g) of Barnyard millet is higher among all the millets and protein content is from 11.1% to 13.9%.

"Proso millet the fat content is 2nd highest among millets i.e., 4.0g/100g and the predominant fatty acids in the free lipids are linoleic, oleic, and palmitic acids" [25]. "The protein content is 10.6 g/100g. The protein of proso millet is gluten free and can be used for foods for people with gluten intolerance or celiac disease. It is also a good source of Vitamin B2, B3 and B9. It is also rich in fibre and minerals i.e., 12g and 2.9 per 100g, respectively. Proso millet is also a good source of dietary fibre and has a lower glycaemic index" [26].

"About 5-8% of protein is present in finger millet, 65-75% carbohydrates, 15-20% dietary fiber and 2.5-3.5% minerals" [28]. "The Naked caryopsis of finger millet with brick red coloured seed coat is generally used in the form of whole meal in the traditional food preparations such as roti, muddle and ambali (thin porridge). Regular consumption of whole grain cereals and their products have shown in epidemiological studies that they can protect against risk of diabetes mellitus, gastrointestinal diseases and cardiovascular risks" [29]. "The use of millets as whole grain makes the essential nutrients such as dietary fiber, minerals, phenolics and vitamins concentrated in the outer layer of the grain or the seed coat form the part of the food and offer their nutritional and health benefits" [30].




Fig. 3. Nutrient Composition of different millets
[Source: 27]

“Sorghum has 11.9 per cent of moisture and about 10.4 per cent of protein and a lower fat content of 1.9 per cent. The fibre and mineral content of grain sorghum is essentially similar, and is 1.6 per cent. It is a good source of energy and provides about 349 K cal and gives 72.6 per cent of carbohydrates” (Gopalan et al. 1996). “Starch is the major carbohydrate of the grain. The other carbohydrates present are simple sugars, cellulose and hemicellulose. The amylose content of starch varies from 21.28 per cent. Sorghum is also rich in dietary fibre (14.3%). Calcium, phosphorous and iron content of sorghum is 25 mg, 222 mg and 4.1 mg (per 100 g of edible portion), respectively” (Hosmani and Chittapur, 1997).

Millets like Jowar has protein content of 10.4g, Bajra 11.6g, the 12.5g of proso millets, 12.3g of foxtail millet, and 11.6 of barnyard millet is equal to wheat’s 11.8g and significantly greater than

rice’s 6.8g. In comparison to wheat and rice, finger millet has lower protein content (7.3g), but it is higher in mineral matter and calcium. Millets in general have more fibre than fine cereals. In compared to wheat 1.2g and rice 0.2g, tiny millets such as barnyard millet 14.7g, kodo millet 9g, little millet 8.6g, and foxtail millet 8.0g are the richest in fibre. As a result, millets are now referred to as "Miracle grains/ Adbhut Anaj and nutria-cereals"[31]. “Finger millet has 16 times the calcium content of maize, and some believe it could eventually replace rice as a staple diet, which is especially important to humans due to the availability of key minerals” [32].

“Consuming little amounts of millet on a daily basis reduced the chance of developing type 2 diabetes. This happens because whole grains like minor millet are high in magnesium, which serves as a co-factor in a variety of biochemical

activities in the body and hence aids glucose and insulin secretion” [33].

3. MILLETS FOR CLIMATE SMART AGRICULTURE

“Future agriculture will face some common environmental changes like enhanced temperature, uncertainties in rainfall, elevated CO₂ and GHGs levels, and more frequency in natural calamities. Under these conditions, climate resilient agriculture should be adopted in which cultivation of climate smart crops will play a pivotal role. There is no doubt that millets are the climate smart crops which can simultaneously mitigate the ill effects of climate change and adapt to the changed and wider agro-climatic conditions” [34]. “Millets have some efficient morphological, physiological, molecular and biochemical traits which can withstand abiotic stresses. As millets are short-duration crops, thus completing a cycle within a short time span, they can escape the possibility of environmental stress conditions under early or late sowing conditions. Besides, millets possess less leaf area a thickened cell wall and a dense fibrous root system which facilitates their capability to tolerate abiotic stress” [35]. “Being C₄ plants, millets can utilize more atmospheric CO₂ and by the process of photosynthesis can

produce more assimilates, even under elevated CO₂ levels into the atmosphere” [36]. Furthermore, “water use efficiency (WUE) of millets is higher than major cereals and, in the future, under the crucial situation of water deficit in a major portion of the world, millets will automatically be chosen to combat water shortage. For example, foxtail millet needs only 257 g of water to produce 1 g of dry biomass and the water requirement is half that of maize (470 g) and wheat (510 g) to produce the same quantity of dry biomass” (Bandopadhyay et al. 2017). Wheat and rice have a global warming potential of around 4 tons CO₂ eq/ha and 3.4 tons CO₂ eq/ha, respectively. Wheat, rice and maize also have high carbon equivalent emissions of 1000, 956 and 935 kg C/ha for wheat, rice and maize, respectively (Jain et al. 2016). The carbon footprint of millets is comparatively less than major cereals and therefore cultivation of millets can reduce the carbon footprint [37]. Furthermore, in the process of chemical nitrogen fertilizer production, CO₂ is generated and as per an estimate to fulfill the present requirement of chemical nitrogen fertilizers in the world, annually 300 tera-gram (Tg = 10¹²g) of CO₂ is released into the atmosphere [38]. Millets are less nutrient demanding crops and promotion of millet cultivation will indirectly save the environment.




Fig. 4. Unique properties of millets for climate smart agriculture, ensuring human health, food and nutritional security [Source: 39]

“The demand for food will increase proportionally with a growing population. While maize, rice and wheat have been adopted as the major staple cereals, millets and other orphan crops are lagging behind. There is a lesser possibility of crop improvement production as the world is already facing the challenges of drylands expansion, soil degradation, and groundwater scarcity” [11]. “Millets are the best choice among orphan crops and their cultivation can solve this problem as they can be grown on shallow, low fertile soils with a varied (ranging from 4.5 to 8.0) pH” [39]. There are many small types of millet such as finger, foxtail, proso, barnyard, Kodo, little, guinea, brown top, teff, and fonio. Millets can be an easy replacement for wheat and rice. Further, millets like pearl and finger millet can grow up to a soil salinity of 11–12 dS/m, while rice has poor growth and productivity on a soil having salinity higher than 3 dS/m [39]. They are considered as a poor man’s crop due to their significant contributions to a resource-limited population diet offering several opportunities for their cultivation in developing countries (Fig. 4).

4. DROUGHT AND MILLETS: IMPACT AND ADAPTATION

Crop productivity is affected by a number of biotic and abiotic stresses. Rapid changes in climate led to major losses of arable land used for crop production and imposing abiotic stresses during critical plant growth and development stages causing yield losses. In semi-arid and arid regions, abiotic stress such as drought, extreme temperature (cold, frost and heat), flooding, salinity etc. are the major yield limiting stressors. Millets encompass numerous morpho-physiological, molecular, and biochemical properties that confer better tolerance to environmental stresses than major cereals. Molecular mechanisms underlying the plants responses to abiotic stresses include multitude of processes including sensing, signaling, transcription, transcript processing, translation and post-translational protein modifications which are governed by both genetic and epigenetic factors (Fig. 5). “Among all major abiotic

stresses, increased drought and heat due to climate change adversely affect current crop production and alone cause more annual losses. The climate change models predict that drought stress would continue as a major abiotic limitation for food production” [40].

“Millets encompass numerous morpho-physiological, biochemical and molecular traits that confer superior adaptation to drought than major cereals (Fig. 6). The rainfall requirement of pearl and proso millet is 20 cm, which is many folds lower than rice as they require more than 120–140 cm” [27]. “The short life-cycle of millets (~10–12 weeks) as compared to other major crops (20–24 weeks) also supports them in stress mitigation. Millets have enhanced photosynthetic rates at warm conditions and confer immediate water and nitrogen use efficiency, which is ~1.5- to 4-fold higher than C_3 photosynthesis” [41]. For instance, *Setaria italica* requires just 257 g of water to produce dry biomass of 1 g, whereas maize and wheat require 470 and 510 g, respectively [42]. Additionally, “ C_4 photosynthesis provides secondary benefits to millets, including better growth and ecological performer in warm temperatures, enhanced flexible allocation patterns of biomass, and reduced hydraulic conductivity per unit leaf area” [43].

“Compared to maize, pearl millet can modulate their membrane dynamics better for water permeability to attain better water status during osmotic stress [34]. An increase in leaf tensile strength and root length was reported in teff and little millet under drought” [44]. “Several biochemical events, e.g., reactive oxygen species (ROS) regulation, enhances ROS scavenging enzymes (catalase and superoxide), and other stress-related proteins. The accumulation of antioxidants and osmolytes has been reported in response to abiotic stresses in millets” [45] These qualities of millets make them a suitable model system holding the potential for research to explore the stress-responsive traits and delineate the mechanism of stress at the physiological, biochemical, and molecular levels.




Fig. 5. A schematic framework of abiotic stress response sensing, signaling and regulation in millet cells. Abiotic stresses alter morpho-physiological, biochemical and molecular processes and also by modulating the rhizosphere properties and functions in millets [Source: 40]




Fig. 6. Drought responses and adaptive strategies linked to numerous morpho-physiological, molecular, and biochemical processes that confer better tolerance to environmental stresses in millets compared to major cereals [Source: 27]

5. INCORPORATION OF MILLETS IN INDIAN FOOD SUPPLY CHAIN

- To feed the population we first need to produce then only we can process millets and enjoy its nutritional benefits [9]. “But the scope for enhancement of productivity under irrigated conditions is limited because of over-exploitation of available resources, but there is ample opportunity for boosting yield in dry-lands by adopting suitable crops and cropping systems. The combination of cereal and legume in intercropping can be a major help to the farmers in subsistence farming targeting livelihood security” [46]. “They also have numerous advantages, such as increased crop productivity, increased resource efficiency, reduced water run-off and soil conservation in erosion-prone areas, prevention of soil nutrient loss, improved soil health, insurance against crop failure due to unusual weather, and a higher monetary return and benefit-cost ratio” [47,48].
- Snacking is becoming a common practice especially in children and adults therefore an attempt to develop some healthy snacks like muffin cakes and biscuits from processed malted finger millet flour to get

the maximum advantage of their nutrient content in terms of bioavailability. Malting Finger millet malt (FMM) can be prepared by different methods given by with slight modifications that is by steeping finger millet grains for 18-24 h at room temperature and allowed to sprout for 48-120 h. Grains were dehydrated for 60 min at 60°C. Further rootlets of grains were removed and powdered to obtain flour for later usage. Also, the above process can be repeated by blanching, pressure cooking or roasting the grains field [49].

- After fermenting and cooking the ready-made mix to create idli and dosa, germinated powders of minor millets were blended and incorporated with other fundamental traditional components like rice powder and de-husked black gramme powder in defined proportions. In comparison to rice-based idli, high proportions of protein (15-18%), fat (8.5-9.8), and carbohydrate (69-72%) were determined for dosa. In millet-based meals, processing stages such as decortication, germination, and fermentation also considerably reduced anti nutrients such phytic acids (69%) and tannin (78%) concentration [50].




Fig. 7. Incorporation of millets in Indian food supply chain

- “A key caveat in achieving the estimated benefits of cereal diversification is the extent to which agronomic characteristics will permit switches between crops. On one hand, historical policy regimes have promoted the widespread cultivation of crops in places that may not have otherwise been agro-ecologically suitable or sustainable (e.g., rice in northern India). On the other hand, certain areas where rice is currently grown may not be able to support the cultivation of coarse cereals. Assessments quantifying the range of biophysical conditions that can support the cultivation of each cereal will therefore be essential for understanding the potential magnitude of co-benefits from increased coarse cereal production” [51]
- Increases in coarse cereal production largely occurred in or close to the places where the cultivation of these cereals is currently centered. This is encouraging from a farmer perspective as the local knowledge of effective crop management practices may be more readily available [52].

6. CONCLUSIONS

Climate change has already had obvious effects on the environment and put challenges for the agricultural sector globally. These extreme climatic events (temperatures rise, rainfall variation, drought etc) add to pressures on agricultural and food systems. This outbreak has disrupted many agriculture and supply chain activities, compounding food and nutrition security challenges, and sustaining livelihoods. Their impacts have the most negative effects on developing countries by adding resource problems, such as water scarcity, pollution, and soil degradation. There are huge genetic resources in terms of landraces, varieties are available in different millets but not much attention has been given to work the responsiveness of varieties under different soil types.

Millets have low glycaemic index, abode gluten-free protein and are rich in minerals (calcium, iron, copper, magnesium, etc.), B-vitamins and antioxidants. These extraordinary traits make them nutritious and climate change compliant crops. These can not only serve as an income crop for farmers but also improve the health of the community as a whole. Therefore, we advocate using the term “climate smart/resilient”

agriculture considering several elements such as: (i) more prominent role for locally produced important millet crops, (ii) increase in production of nutritious food and value addition to help the human immune system, (iii) creation of low-cost farming systems with less dependence on water and chemicals, (iv) leveraging agro-ecology to synthesize the environment with farming, and (v) nudging consumers’ demand toward climate-resistant grains, and making farming more feasible for marginal farmers.

In India, diversifying crop production to include more coarse cereals, such as millets and sorghum, can make food supply more nutritious, reduce resource demand and greenhouse gas emissions, and enhance climate resilience without reducing calorie production or requiring more land. Millets can contribute to sustainable food systems under climate change. Their resilient nature and outstanding potential to survive under low water availability and stressful environments serves as best alternative to staple cereal crops.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Food and Agriculture Organization of the United Nations [technical note]. FAO methodology to estimate the prevalence of undernourishment. Rome, Italy: Food and Agriculture Organization; 2012.
2. Godfray HCJ, Beddington JR, Crute IR, Haddad L, Lawrence D, Muir JF et al. Food security: the challenge of feeding 9 billion people. *Science*. 2010;327(5967): 812-8.
3. Wheeler T, Von Braun J. Climate change impacts on global food security. *Science*. 2013;341(6145):508-13.
4. Davis KF, Chhatre A, Rao ND, Singh D, DeFries R. Sensitivity of grain yields to historical climate variability in India. *Environ Res Lett*. 2019;14(6):064013.
5. Banerjee P, Maitra S. The role of small millets as functional food to combat malnutrition in developing countries. *Indian J Nat Sci*. 2020;10(60):20412-7.
6. Reynolds TW, Waddington SR, Anderson CL, Chew A, True Z, Cullen A. Environmental impacts and constraints associated with the production of major

- food crops in Sub-Saharan Africa and South Asia. *Food Sec.* 2015;7(4):795-822.
7. Pasricha SR, Biggs BA. Under-nutrition among children in south and South-East Asia. *J Paediatr Child Health.* 2010;46(9): 497-503.
 8. Rao ND, Poblete-Cazenave M, Bhalerao R, Davis KF, Parkinson S. Spatial analysis of energy use and GHG emissions from cereal production in India. *Sci Total Environ.* 2019;654:841-9.
 9. Saleh ASM, Zhang Q, Chen J, Shen Q. Millet grains: nutritional quality, processing, and potential health benefits. *Compr Rev Food Sci Food Saf.* 2013;12(3):281-95.
 10. Patwari AK. Millennium Development Goals and child under nutrition. *Indian Pediatr.* 2013;50(5):449-52.
 11. Bisoffi S, Ahrné L, Aschemann-Witzel J, Báldi A, Cuhls K, DeClerck F, et al. COVID-19 and sustainable food systems: what should we learn before the next emergency. *Front Sustain Food Syst.* 2021;5:650987.
 12. FAO. The state of food insecurity in the world 2015. *Food Agric Organ UN.* 2015.
 13. Von Grebmer K, Bernstein J, Hossain N, Brown T, Prasai N, Yohannes Y. Global hunger index: the inequalities of hunger. *International Food Policy Research Institute;* 2017.
 14. Lozano R, Naghavi M, Foreman K, Lim S, Shibuya K, Aboyans V et al. Global and regional mortality from 235 causes of death for 20 age groups in 1990 and 2010: a systematic analysis for the Global Burden of Disease Study 2010. *Lancet.* 2012;380(9859):2095-128.
 15. Gragnoliata M, Shekarb M, Gupta MC, Bredenkampd C, Lee Y. India's undernourished children: A call for reform and action, health, nutrition and population (HNP) discussion paper. *Washington, DC: World Bank;* 2005.
 16. Kaveeshwar SA, Cornwall J. The current state of diabetes mellitus in India. *Australas Med J.* 2014;7(1):45-8.
 17. Choi YY, Osada K, Ito Y, Nagasawa T, Choi MR, Nishizawa N. Effects of dietary protein of Korean foxtail millet on plasma adiponectin, HDL-cholesterol, and insulin levels in genetically type 2 diabetic mice. *Biosci Biotechnol Biochem.* 2005;69(1):31-7.
 18. Lee SH, Chung IM, Cha YS, Park Y. Millet consumption decreased serum concentration of triglyceride and C-reactive protein but not oxidative status in hyperlipidemic rats. *Nutr Res.* 2010;30(4): 290-6.
 19. Hegde PS, Chandrakasan G, Chandra TS. Inhibition of collagen glycation and cross linking in vitro by methanolic extracts of finger millet (*Eleusine coracana*) and kodo millet (*Paspalum scrobiculatum*). *J Nutr Biochem.* 2002;13(9):517.
 20. Dykes L, Rooney LW. Review sorghum and millet phenols and antioxidants. *J Cereal Sci.* 2006;44(3):236-51.
 21. Krishna Kumari SK, Thayumanavan B. Characterization of starches of proso, foxtail, barnyard, kodo, and little millets. *Plant Foods Hum Nutr.* 1998;53(1):47-56.
 22. Sireesha Y, Kasetti RB, Nabi SA, Swapna S, Apparao C. Antihyperglycemic and hypolipidemic activities of *Setaria italica* seeds in STZ diabetic rats. *Pathophysiology.* 2011;18(2):159-64.
 23. Devi PB, Vijayabharathi R, Sathyabama S, Malleshi NG, Priyadarisini VB. Health benefits of finger millet (*Eleusine coracana* L.) polyphenols and dietary fiber: a review. *J Food Sci Technol.* 2014;51(6): 1021-40.
 24. Sood S, Khulbe RK, Gupta AK, Agrawal PK, Upadhyaya HD, Bhatt JC. Barnyard millet—a potential food and feed crop of future. *Plant Breed.* 2015;134(2):135-47.
 25. Amadou I, Gounga ME, Le GW. Millets: nutritional composition, some health benefits and processing-A review. *Emirates J. Food Agric.* 2013:501-8.
 26. Park KO, Ito Y, Nagasawa T, Choi MR, Nishizawa N. Effects of dietary Korean proso-millet protein on plasma adiponectin, HDL cholesterol, insulin levels, and gene expression in obese type 2 diabetic mice. *Biosci Biotechnol Biochem.* 2008;72(11): 2918-25.
 27. Kumar A, Tomer V, Kaur A, Kumar V, Gupta K. Millets: a solution to agrarian and nutritional challenges. *Agric & Food Secur.* 2018;7(1):31.
 28. Chethan S, Malleshi NG. Finger millet polyphenols: optimization of extraction and the effect of pH on their stability. *Food Chem.* 2007;105(2):862-70.
 29. McKeown NM, Meigs JB, Liu S, Wilson PW, Jacques PF. Whole-grain intake is favorably associated with metabolic risk factors for type 2 diabetes and cardiovascular disease in the Framingham Offspring Study. *Am J Clin Nutr.* 2002; 76(2):390-8.

30. Antony U, Sripriya G, Chandra TS. Effect of fermentation on the primary nutrients in finger millet (*Eleusine coracana*). J Agric Food Chem. 1996;44(9): 2616-8.
31. Senthilvel S, Jayashree B, Mahalakshmi V, Kumar PS, Nakka S, Nepolean T et al. Development and mapping of simple sequence repeat markers for pearl millet from data mining of expressed sequence tags. BMC Plant Biol. 2008;8(1):119.
32. Hassan ZM, Sebola NA, Mabelebele M. The nutritional use of millet grain for food and feed: a review. Agric Food Secur. 2021;10(1):16.
33. Saikia D, Deka SC. Cereals: from staple food to nutra-ceuticals. Int Food Res J. 2011;18(1).
34. Bandyopadhyay T, Muthamilarasan M, Prasad M. Millets for next generation climate-smart agriculture. Front Plant Sci. 2017;8:1266.
35. Li P, Brutnell TP. *Setaria viridis* and *Setaria italica* model genetic systems for the panicoid grasses. J Exp Bot. 2011;62(9):3031-7.
36. Aubry S, Brown NJ, Hibberd JM. The role of proteins in C3 plants prior to their recruitment into the C4 pathway. J Exp Bot. 2011;62(9):3049-59.
37. Saxena R, Vanga SK, Wang J, Orsat V, Raghavan V. Millets for food security in the context of climate change: a review. Sustainability. 2018;10(7):2228.
38. Jensen ES, Peoples MB, Boddey RM, Gresshoff PM, Hauggaard-Nielsen H, J.R. Alves B et al. Legumes for mitigation of climate change and the provision of feedstock for biofuels and biorefineries. A review. Agron Sustain Dev. 2012;32(2): 329-64.
39. Rathinapriya P, Pandian S, Rakkammal K, Balasangeetha M, Alexpandi R, Satish L, et al. The protective effects of polyamines on salinity stress tolerance in foxtail millet (*Setaria italica* L.), an important C4 model crop. Physiol Mol Biol Plants. 2020;26(9):1815-29.
40. Simmons T, Styer AB, Pierroz G, Gonçalves AP, Pasricha R, Hazra AB, et al. Drought drives spatial variation in the millet root microbiome. Front Plant Sci. 2020;11:599.
41. Wang C, Guo L, Li Y, Wang Z. Systematic comparison of C3 and C4 plants based on metabolic network analysis. BMC Syst Biol. 2012;62;Suppl 2:1-14.
42. Nadeem F, Ahmad Z, UI Hassan M, Wang R, Diao X, Li X. Adaptation of foxtail millet (*Setaria italica* L.) to abiotic stresses: a special perspective of responses to nitrogen and phosphate limitations. Front Plant Sci. 2020;11: 187.
43. Lundgren MR, Osborne CP, Christin PA. Deconstructing Kranz anatomy to understand C4 evolution. J Exp Bot. 2014;65(13):3357-69.
44. Balsamo RA, VanderWilligen CV, Bauer AM, Farrant J. Drought tolerance of selected *Eragrostis* species correlates with leaf tensile properties. Ann Bot. 2006;97(6):985-91.
45. Ajithkumar IP, Panneerselvam R. ROS scavenging system, osmotic maintenance, pigment and growth status of *panicum sumatrense* roth. Under Drought Stress. Cell Biochem Biophys. 2014;68(3): 587-95.
46. Maitra S, Pine S, Shankar T, Pal A, Pramanick B. Agronomic Management of foxtail millet (*Setaria italica* L.) in India for Production Sustainability: a review. Int J Bioresour Sci. 2020;7(1):11-6.
47. DeVincentis AJ, Solis SS, Bruno EM, Leavitt A, Gomes A, Rice S et al. Using cost benefit analysis to understand adoption of winter cover cropping in California's specialty crop systems. J Environ Manage. 2020;261:110205.
48. Maitra S, Pine S, Banerjee P, Shankar T. Millets: robust entrants to functional food sector. In: Pirzadah TB, Malik B, Bhat A, Hakeem KR, editors, Bioresource technology: concept, tools and experiences. Wiley Online Library; 2022.
49. Khokhar S, Apenten RKO. Anti-nutritional factors in food legumes and effects of processing. The Role Food Agric For Fish Hum Nutr. 2003;4:82-116.
50. Krishnamoorthy S, Kunjithapatham S, Manickam L. Traditional Indian breakfast (Idli and Dosa) with enhanced nutritional content using millets. Nutr Diet. 2013; 70(3):241-6.
51. DeFries R, Chhatre A, Davis KF, Dutta A, Fanzo J, Ghosh-Jerath S, et al. Impact of historical changes in coarse cereals consumption in India on micronutrient intake and anemia prevalence. Food Nutr Bull. 2018;39(3):377-92.

52. Borelli T, Hunter D, Padulosi S, Amaya N, Meldrum G, de Oliveira Beltrame DM, et al. Local solutions for sustainable food systems: the contribution of orphan crops and wild edible species. *Agronomy*. 2020;10(2): 231. DOI: 10.3390/agronomy10020231.

© 2022 Kumar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/92823>