



Epidemiological Studies of Polyparasitism and Gender Associated Factors in Women in Igede Land

P. O. Uweh ^{a*} and E. A. Omudu ^b

^a Department of Biology, Akawe Torkula Polytechnic, Makurdi, Benue State, Nigeria.

^b Department of Biological Sciences, Benue State University, Makurdi, Nigeria.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJTDH/2022/v43i1630652

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/89902>

Original Research Article

Received 15 June 2022
Accepted 07 August 2022
Published 16 August 2022

ABSTRACT

Introduction: Polyparasitism remains one of the most pressing public health problems with one third of the world population generally estimated to be infected. Also, Women are particularly of concern due to their gender roles which is sensitive since ill health and/or the illness of family members generally represent greater burdens for poor women in comparison to poor men.

Aim: To epidemiologically study polyparasitism in Women of Reproductive age in Igede land of Benue State, Nigeria.

Study Design: A cross-sectional study design was adopted for this study. Fresh stool, urine and blood samples were collected from women in Igede Land of Benue State and processed for internal parasites, urine parasites and malaria parasite respectively from September 2019 to August, 2021 using systematic random sampling technique. The study was restricted to women of reproductive age in Igedeland only. Well-structured questionnaires were further utilized to elicit information on associated factors related to polyparasitism. Positive samples were afterwards processed molecularly using PCR technology and the results documented. Data obtained was analysed using SPSS and comparative analysis was done using chi-square statistical tool.

Results: The overall rate of polyparasitism in this study was 3.50% while a prevalence of 33.50% was recorded for single parasitic infection. The dominant parasite was *Plasmodium* spp. (52.16%) while the least was *Strongyloides stercoralis* (1.44%). Parasite combination involving *Schistosoma haematobium* and Malaria parasite recorded highest prevalence (36.11%) while least prevalence was recorded for coinfection with Hookworm, *Entamoeba histolytica* and *Ascaris lumbricoides*.

(8.33%). Also, associated factors of excretion and water source were significant ($P < 0.05$) while meat consumption and sanitation were not significant ($P > 0.05$).

Conclusion: Polyparasitism among women in Igedeland of Benue State is relatively low with malaria parasite being the most dominant parasite.

Recommendations: Awareness on polyparasitism should be increased among women and antihelminthic and antimalarial drugs should be distributed free of charge to women so as to help reduce the burden of parasite infection in the study area.

Keywords: Polyparasitism; risk factors; Plasmodium; women, Igede.

1. INTRODUCTION

There have been a general renaissance in the epidemiological investigation of polyparasitism, with a particular focus on multiple helminth species and more recently, on *Plasmodium*-helminth co-infections [1,2,3]. Studies across multiple epidemiological settings have shown that polyparasitism is the norm rather than the exception and occurs at different frequencies than would be expected under assumptions of independence [4,5,6]. Interactions between parasites in humans can be synergistic or antagonistic. For example, studies have demonstrated a positive association between intensity and concurrent infection of helminth species, suggesting that individuals harbouring multiple helminth species also harbour the most intense infections [4,5,6]. It is conceivable therefore that polyparasitism may have a greater impact on morbidity than single species infections since morbidity is typically related to infection intensity for most parasite species. Multiple species infections may also increase susceptibility to other infections [3,7].

Although accurate estimate of the magnitude of polyparasitism is unknown, the case is distributed widely in the tropics and sub-tropics [8]. It was found that individuals with polyparasitic infection also harbor the most intense infections [1,6]. Even low-intensity of polyparasitic infections may result in clinically significant morbidity [9,10]. Therefore, polyparasitism may have a greater impact on morbidity than single species infection. It may also increase susceptibility to other infections and adversely affect the clinical outcome of the concomitant diseases such as tuberculosis, HIV/AIDS, and malaria [7,11].

Several findings from different parts of the world have revealed tangible information pertaining to the impact of polyparasitism on general health. It has been reported that infants and pregnant women are the group of individuals most likely to harbour multiple parasite infections [12]. It is

opined that Malaria, hookworm, schistosomiasis and, to a lesser extent, *Trichuris trichiura* are associated with iron loss and imbalances predisposing to anaemia among nutritionally vulnerable populations, with risk being closely correlated with infection intensity [13]. Also, high iron demands of infant growth and pregnancy means that anaemia is most common and severe among children and pregnant women [14]. The gut protozoa such as *Entamoeba histolytica*, *Giardia lamblia* and *Cryptosporidium* spp. are also highly prevalent and of major importance in children, pregnant women and in HIV-infected patients. Furthermore, polyparasitism is higher among younger children than adults. They attributed this to factors such as environment and personal hygiene [15].

While existing literature reveals that both poor men and women suffer greater ill health than their more well off counterparts, it also indicates that ill health and/or the illness of family members generally represent greater burdens for poor women in comparison to poor men [16,17]. This is because in addition to the resource constraints that affects the poor in general, there are gender inequalities and inequities that can place women in particular at a further disadvantage. The reasons for this are multiple, and involve the interplay of many factors. First of all, in all cultures, ethnic and age groups, men and women are assigned different roles and responsibilities. This assignment shapes the development of different skills and abilities and channels their application to specific life spaces or spheres of activity. This differential value has direct implications for the level of access to and control of resources available to men and women to protect their health, the health of their families. In sum, socially constructed characteristics that are attributed to one sex or the other at the individual level, or that are associated with male and female sphere of activity at the societal level, give rise to gender inequalities. They also perpetuate gender that can enhance or deter the

health and development of men and women, as individuals and as population sub-groups [18,19].

The statements made by the above researchers as well as findings from previous research have formed a good background to the study of polyparasitism over the years. However, little is known about the impact of polyparasitism on the health of women in the world, North central Nigeria and Benue State in particular. This is therefore the driving force behind this study.

2. MATERIALS AND METHODS

2.1 Study Area and Population

The study was carried out in Oju and Obi local government areas in the southern part of Benue State. Oju is located between latitude $6^{\circ}6'N$ and

$6^{\circ}9'N$ and Longitude $8^{\circ}10'E$ and $8^{\circ}25'E$ in the south eastern part of Benue State. Oju local government area covers an area of 1,283 km². It is bounded in the south by Cross River State, in the east by Konshisha local government area, in the north by Obi and Gwer East local government areas, in the west by Ado local government area, and in the south-west by Ebonyi State. The widely disputed result of the 2006 national population and housing census put the population of Obi local government area of Benue State at 98,707, with 49,143 males and 49,564 females. Obi local government area covers an area of 423 km². It is bounded in the north-east by Gwer East local government area, in the south by Oju local government area, in the west by Ado local government area, and in the north by Otukpo local government area [16].




Fig. 1. Map of Benue State showing Location of Oju and Obi Local Government Area [16]

2.2 Experimental Design

A Cross sectional study design was adopted for the study. All women in the areas visited were given equal opportunity to participate in the study. Women were briefed on the health implication of being infected with parasites, the dangers of parasitism and the relevance of the study to each participant. Participation was made voluntary. Volunteers were made to fill a written consent form indicating their willingness to participate in the study.

Study Period: The study was conducted from September 2019 to August, 2021.

Eligibility Criteria: Only women of reproductive age were allowed to participate in the study. This includes women who are 18years and above and who consent to participate in the study.

Sample Size: The sample size used for the study (1037) was determined using the formula below [19].

$$n = \frac{z^2 p(1 - p)}{d^2}$$

Where

n = sample size

z = statistic for a level of confidence, in this case the level of confidence will be 95% (1.96)

p = expected prevalence [20]

d = precision at 5% (0.05)

2.3 Sample Collection and Sampling

A systematic random sampling technique was utilized in this study. Specimen bottles were distributed to the consenting women who participated in the study. They return the containers with urine and stool in them. Blood was however collected intravenously by laboratory technicians from women using appropriate aseptic techniques. Blood samples were examined immediately, while stool and urine samples were taken to the laboratory for microscopic examination.

Demographic information, clinical manifestations of parasites as well as information pertaining to parasitism were collected from the study subjects using well-structured questionnaires so as to enable adequate correlation of results obtained with demographic variables.

2.4 Examination of Faecal Samples

Faecal samples collected during the study were examined according to the formol ether concentration technique [18]. Using an applicator stick, an estimated 1g (pea-size) of faeces was emulsified in about 4 ml of 10% formol water contained in a screw-cap tube. A further 3–4 ml of 10% v/v formol water was added, the bottle capped, and mixed well by shaking. The emulsified faeces were sieved, collecting the sieved suspension in a beaker and the suspension was transferred to a conical (centrifuge) tube made of strong glass, and about 3–4 ml of ethyl acetate was added. The tube was stoppered and mixed for 1 minute and with a tissue or piece of cloth wrapped around the top of the tube, the stopper was loosened. The content of the tube was centrifuged immediately at 750–1000 g (approx. 3000 rpm) for 1 minute. Using a stick, the layer of faecal debris from the side of the tube was loosened and the tube inverted to discard the ether, faecal debris, and formol water while the sediment remained. The tube was returned to its upright position, and the fluid from the side of the tube was allowed to drain to the bottom. The bottom of the tube was tapped to re-suspend and mix the sediment, and the sediment was transferred to a slide, and covered with a cover glass. The preparation was examined microscopically using the 10x objective with the condenser iris closed sufficiently to give good contrast. The 40x objective was used to examine cysts and eggs. To assist in the identification of cysts, a small drop of iodine was run under the cover glass [21].

2.5 Examination of Urine Samples

Urine samples were examined microscopically. About 10 ml of well mixed urine was aseptically transferred to a labeled conical tube and centrifuged at 500–1000 g for 5 minutes. The sediment was afterwards remixed by tapping the bottom of the tube after which one drop of the *well-mixed* sediment was transferred to a slide and covered with a cover slip. The preparation was examined microscopically using the 10x and 40x objective with the condenser iris *closed sufficiently* to give good contrast [21].

2.6 Examination of Blood Samples

Blood samples were examined for malaria parasite using a Rapid Test Diagnostic Kit. The area to be pricked was cleansed with an alcohol swab after which the end of the fingertip was

squeezed gently and pierced with a sterile lancet provided in the test kit. 5µl of whole blood was collected using the pipette provided in the test kit and added into the sample well of the test device, then two drops of the assay buffer was added into the buffer well. The result was read within 20 minutes and recorded accordingly.

The result was said to be positive if two colour bands appear, one at the control line 'C' and the other at the test line 'T'. Negative result occurred if only one colour band appear, at control line 'C'. In a case where no colour band appears, or colour band appeared on the test line, the result was said to be invalid and repeated as such [21].

2.7 Molecular Analysis of Samples

Representative positive samples were sent to FOWM Biotechnology Limited, Yaba, Lagos and Molecular Laboratory, Covenant University Centre for Research, Innovation and Development (CUCRID) for DNA extraction and amplification (Polymerase Chain Reaction) so as to confirm the true identity of the parasite identified.

2.8 Data Analysis

Data obtained from parasitological analysis and questionnaire administration were entered into IBM Statistical Package for Social Sciences (SPSS) version 21.0. The percentage prevalence (%) was calculated in each case. Descriptive statistics such as proportion and percentages were computed for sociodemographic data while associations and relationship between risk factors and parasitic infections were tested using chi square (χ^2) at 95% confidence level. A p-value less than .05 ($P < .05$) was considered statistically significant.

3. RESULTS AND DISCUSSION

The overall rate of polyparasitism in this study was 3.50%. The parasites encountered include: *Entamoeba histolytica*, Malaria parasite, *Schistosoma haematobium*, Hookworm, *Ascaris lumbricoides*, and *Trichuris trichuria* with prevalence of 9.51%, 6.05%, 5.47%, 1.44%, 1.72%, 52.16%, 18.15% and 5.47% respectively.

In terms of parasite combinations, coinfection with malaria parasite and *Schistosoma haematobium* recorded the highest prevalence (36.11%). Also, Idelle community recorded the highest prevalence of polyparasitism while Oju centre recorded the least prevalence (Fig. 2).

In terms of risk factors of infection, excretion and water sources were observed to have a significant association with rate of polyparasitism ($P < 0.05$) while meat consumption and sanitation were not statistically significant. Results obtained showed that women who utilize pit toilet recording the highest prevalence of infection (7.10%) while those who utilize water system recorded the least prevalence (2.00%).

Women who utilize rainwater recorded highest prevalence rates (10.60%) while the least prevalence was recorded among women who use well other sources of water (1.70%). For meat consumption, women who consume pock the most recorded the least prevalence of infection (2.10%) while those who consume goat meat recorded the highest prevalence of infection (4.60%). Also, women who use other means of sanitary disposal recorded least prevalence of infection with polyparasites while those who utilize garbage pits recorded the highest prevalence of infection (4.80%).

Table 1. Distribution patterns of polyparasites among women of igede land, Benue State, Nigeria.

Parasite Combination	Frequency	Percentage
E.H+MP+S.H	4	11.11
HW+A.L+MP	3	8.33
MP+SH+AL	3	8.33
HW+EH+SH	3	8.33
MP+SH	13	36.11
MP+TT	6	16.67
MP+TT+HW	4	11.11
Total	36	100.00

E.H= Entamoeba histolytica, MP= Malaria parasite, SH= Schistosoma haematobium, HW: Hookworm, AL= Ascaris lumbricoides, TT= Trichuris trichuria




Fig. 2. Polyparasitism in the selected communities surveyed

Findings from the study revealed a 33.50% prevalence of parasitic infection among the study population with 3.50% of them suffering from polyparasitism. This prevalence of polyparasitism recorded among women of Oju and Obi LGAs, Benue State is comparatively low compared to what was reported in past in Naka area of Benue State where a cumulative prevalence of 9.8% was reported [20]. It was also lower than the reported prevalence of 13.86% for polyparasitism in Oju Local Government Area of Benue State [22]. Differences between the findings of this study and that of Omudu et al. [20] might be attributed in part to the differences in the geographical conditions of the various study location since Naka regions are dominated mainly by the Tiv people whose practices and beliefs are quite different from that of the Iggede people. A possible reason for the variation experienced between the findings of this study and that of Akor et al. [22] [202 may be due to the gender bias of this study which was limited to women alone unlike what was obtainable in the study of [12] where the study population encompassed both males and females.

The prevalence of polyparasitism among women of Oju and Obi LGAs as observed in this study is lower than that of Cheesbrough M [21], who

reported a prevalence of 10.3% in Okpokwu area of Benue State [22]. It is also lower than a 12.2% prevalence reported for polyparasitism among children in Nassarawa State, Nigeria [23]. A prevalence of 3.50% is also lower than an 8.66% prevalence reported for polyparasitism in Kwasa village in Makurdi Nigeria [11]. Differences in the prevalence rates between the retrospective studies highlighted above and that of this study might be attributed in part to the study population examined, the methods of diagnosis and the geographical location which is a predisposing factor to the prevalence of parasitic infection in Nigeria.

Associated factors of infection with parasites was shown to play a significant role in determining the prevalence of polyparasitism among women of Iggede land, Benue State. Findings from the study showed that Excretion place and source of drinking water had a significant association with prevalence of polyparasitism among the study population ($P < 0.05$). Higher prevalence of polyparasitism were recorded for people who utilize pit toilet than those who utilize other forms of toilet. This finding is in agreement with the report of Onah IE et al. [11] who reported a higher prevalence of polyparasitism among people who utilize pit toilet than other forms of

Table 2. Parasite distribution in women of igede land in relation to location

Location	Communities	Number examined	Number Positive (%)	Parasites (%)							
				<i>E. histolytica</i>	<i>A. lumbricoides</i>	Hookworm	<i>S. stercoralis</i>	<i>Taenia</i> spp.	<i>Plasmodium</i> spp.	<i>S. haematobium</i>	<i>Trichuris trichura</i>
Oju	Uje	140	43 (30.71)	3 (6.97)	5 (11.62)	2 (4.65)	0 (0.00)	0 (0.00)	20 (46.51)	13 (30.23)	0 (0.00)
	Uwokwu	30	12 (40.00)	0 (0.00)	0 (0.00)	0 (0.00)	4 (33.33)	2 (16.66)	3 (25.00)	3 (25.00)	0 (0.00)
	Ega	50	30 (60.00)	4 (13.33)	4 (13.33)	1 (3.33)	0 (0.00)	0 (0.00)	9 (30.00)	6 (20.00)	6 (20.00)
	Oju center	34	23(67.64)	3 (13.04)	1 (4.34)	2 (8.69)	0 (0.00)	0 (0.00)	10 (43.47)	7 (30.43)	0 (0.00)
Obi	Idelle	178	58(32.60)	5 (8.62)	4 (6.89)	6 (10.34)	0 (0.00)	0 (0.00)	30 (51.72)	10 (17.24)	3 (5.17)
	Adiko	220	75 (34.54)	9 (12.00)	3 (4.00)	3 (4.00)	0 (0.00)	1 (1.33)	44 (58.66)	12 (16.00)	3 (4.00)
	Obarike	150	33 (22.00)	3 (27.27)	2 (6.06)	0 (0.00)	0 (0.00)	3 (9.09)	18 (54.54)	5 (15.15)	2 (6.06)
	Itogo	150	49 (32.70)	4 (8.16)	1 (2.04)	3 (6.12)	0 (0.00)	0 (0.00)	36 (73.46)	2 (4.08)	3 (6.12)
	Adum	85	24 (28.23)	2 (8.33)	1 (4.16)	2 (8.33)	1 (4.16)	0 (0.00)	11 (45.83)	5 (20.83)	2 (8.33)
Total		1037	347 (33.50)	33 (9.51)	21 (6.05)	19 (5.47)	5 (1.44)	6 (1.72)	181 (52.16)	63 (18.15)	19 (5.47)

$\chi^2=17.05; P=.003$

Table 3. Distribution patterns of polyparasites among women in igede land, Benue State in relation to location

Location	Communities	Number examined	Number Positive	Parasites Combinations						
				<i>E.H+MP+S.H</i>	<i>HW+A.L+MP</i>	<i>MP+SH+AL</i>	<i>HW+EH+SH</i>	<i>MP+SH</i>	<i>MP+TT.</i>	<i>MP+TT+HW</i>
Oju	Uje	140 (13.50)	4 (2.86)	0 (0.00)	1 (25.00)	0 (0.00)	1 (25.00)	2 (50.00)	0 (0.00)	0 (0.00)
	Uwokwu	30 (2.89)	2 (6.67)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)	1 (50.0)	0 (0.00)	1 (50.00)
	Ega	50 (4.82)	3 (6.00)	0 (0.00)	0 (0.00)	1 (33.30)	2 (6.60)	0 (0.00)	0 (0.00)	0 (0.00)
	Oju center	34 (3.28)	1 (2.94)	0 (0.00)	1 (100.0)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)
Obi	Idelle	178 (17.16)	7 (3.93)	2 (28.60)	0 (0.00)	1 (14.30)	0 (0.00)	3 (42.90)	1 (14.30)	0 (0.00)
	Adiko	220 (21.22)	6 (2.72)	2 (33.33)	0 (0.00)	0 (0.00)	0 (0.00)	2 (33.33)	2 (33.33)	0 (0.00)
	Obarike	150 (14.46)	5 (3.33)	1 (20.00)	0 (0.00)	0 (0.00)	0 (0.00)	1 (20.00)	1 (20.00)	2 (40.00)
	Itogo	150 (14.46)	5 (3.33)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)	2 (40.00)	2 (0.00)	1 (20.00)
	Adum	85 (8.20)	3 (3.53)	0 (0.00)	1 (33.33)	1 (33.33)	0 (0.00)	1 (33.33)	0 (0.00)	0 (0.00)
Total		1037(100.00)	36(3.50)	4(11.11)	3(8.33)	3(8.33)	3(8.33)	13(36.11)	6(16.67)	4(11.11)

$\chi^2=16.92 P=.02$

E.H= Entamoeba histolytica, MP= Malaria parasite, SH= Schistosoma haematobium, HW: Hookworm, AL= Ascaris lumbricoides, TT= Trichuris trichuria

Table 4. Polyparasitism in women of igede land in relation to possible risk factors

Risk factor	Number Examined (%)	Number Positive (%)	Polyparasitism (%)	χ^2	P-value
Excretion					
Water System	488	124 (25.40)	10 (2.00)		
Pit Toilet	239	116 (48.50)	17 (7.10)		
Bucket System	222	74 (33.30)	7 (3.20)		
Open defecation	88	33 (37.50)	2 (2.30)	12.848	.005
Water Source					
Rain	85	49 (57.60)	9 (10.60)		
Borehole	468	149 (31.80)	14 (3.00)		
Tap	295	102 (34.60)	10 (3.40)		
Well	119	38 (31.90)	2 (1.70)		
Others	70	9 (12.90)	1 (1.40)	15.185	.004
Meat Consumption					
Beef	353	103 (29.20)	9 (2.50)		
Chicken	168	53 (31.50)	5 (3.00)		
Goat	194	73 (37.60)	9 (4.60)		
Pork	48	28 (58.30)	1 (2.10)		
Fish	270	90(32.80)	12 (4.40)	2.758	.599
Sanitation					
Garbage pit	84	25 (29.80)	4 (4.80)		
Outside the compound	384	141 (36.70)	13 (3.40)		
Waste bin	547	176 (32.20)	19 (3.50)		
Others	22	5 (22.70)	0 (0.00)	1.217	.749
Total	1037	347(33.50)	36 (3.50)		

toilet. People who utilize pit toilets are usually at risk of coming into direct contact with parasites associated with stool since the pits are usually exposed and parasites can easily infect the women when they attempt to defaecate. This finding is also in agreement with the findings of Akor, JO et al. [22] who reported higher polyparasitism rates among people utilizing pit toilet [22]. Higher prevalence recorded for people who utilize rain water could however be attributed to the ability of parasites to be carried in rain water and on surfaces during rain. Also, those who treat rain water recorded a higher prevalence of polyparasitism than those who don't treat water. According to Omudu EA et al. [20], the major reasons behind the high prevalence of polyparasitism can be blamed on the sanitary and environmental condition which they described as deplorable [20]. These conditions in most regions including the study location were below standard and as such might promote the proliferation of parasitic organisms.

4. CONCLUSION

Polyparasitism rate among women of Igede land was 3.50%. This rate was relatively low in comparison to previous rates by previous authors. Also, occurrence of single parasite infection was high with malaria parasite being the most dominant parasite species; an implication

for higher malaria incidence in Igede land if possible preventive measures are not sought and implemented. Also, several associated factors of infection have a significant impact on polyparasitism among women of Igede land, Benue State. These risk factors of infection include: type of toilet utilized and source of drinking water. The parasites encountered which included *Plasmodium* species, *Schistosoma haematobium*, *Ascaris lumbricoides*, *Entamoeba histolytica*, *Taenia* species, *Trichuris trichuria* and Hookworms suggests the existence of a significant public health hazard in the study area.

Based on the findings of this study, it is recommended that the women of Igede land should be enlightened on the health hazards associated with polyparasitism. Also, insecticide treated nets and other antihelminths drugs should be distributed to infected women by the government for free so as to help reduce the burden of polyparasitism in the study area.

ETHICAL APPROVAL AND CONSENT

Ethical approval was obtained from the Benue State Ministry of Health (Ref Number: MOH/STA/204/VOL1/182) prior to the commencement of the study. This was in accordance with the requirements for conducting research on human subjects. Also, informed

consent forms were distributed to the women and were verbally translated to the women in their local languages. Only women who consented to the study by signing the consent forms were recruited for the study.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Brooker S, Akhwale W, Pullan R, Estambale B, Clarke SE, Snow RW, Hotez PJ. Epidemiology of Plasmodium-helminth co-infection in Africa: populations at risk, potential impact on anemia and prospects for combining control. *The American Journal of Tropical Medicine and Hygiene*. 2007;77(6 Suppl):88.
2. Hartgers FC, Yazdanbakhsh M. Co-infection of helminths and malaria: modulation of the immune responses to malaria. *Parasite Immunology*. 2006;28(10):497-506.
3. Mwangi TW, Bethony JM, Brooker S. Malaria and helminth interactions in humans: an epidemiological viewpoint. *Annals of Tropical Medicine & Parasitology*. 2006;1:100(7):551-70.
4. Fleming FM, Brooker S, Geiger SM, Caldas IR, Correa-Oliveira R, Hotez PJ, Bethony JM. Synergistic associations between hookworm and other helminth species in a rural community in Brazil. *Tropical Medicine & International Health*. 2006;11(1):56-64.
5. Howard SC, Donnelly CA, Kabatereine NB, Ratard RC, Brooker S. Spatial and intensity-dependent variations in associations between multiple species helminth infections. *Acta tropica*. 2002;83(2):141-9.
6. Tchuem Tchuente LA, Behnke JM, Gilbert FS, Southgate VR, Vercruysse J. Polyparasitism with *Schistosoma haematobium* and soil-transmitted helminth infections among school children in Loum, Cameroon. *Tropical Medicine & International Health*. 2003;8(11):975-86.
7. Druilhe P, Tall A, Sokhna C. Worms can worsen malaria: towards a new means to roll back malaria?. *Trends in parasitology*. 2005 Aug 1;21(8):359-62.
8. Pullan RL, Brooker SJ. The global limits and population at risk of soil-transmitted helminth infections in 2010. *Parasites & Vectors*. 2012 Dec;5(1):1-4.
9. Ezeamama AE, Friedman JF, Olveda RM, Acosta LP, Kurtis JD, Mor V, McGarvey ST. Functional significance of low-intensity polyparasite helminth infections in anemia. *The Journal of Infectious Diseases*. 2005;192(12):2160-70.
10. Sorensen WC, Cappello M, Bell D, DiFedele LM, Brown MA. Poly-helminth infection in east Guatemalan school children. *Journal of Global Infectious Diseases*. 2011;3(1):25.
11. Onah IE, Omudu EA, Emmy-Egbe IO, Ekwesiana EO, Ukaga CN, Ajero CM, Umeaku CN, Achor JC, Ezeugo GE, Ofochebe AC, Ekwughe CH. Prevalence of polyparasitism in kwasa village (a sub-urban settlement) in makurdi, Benue state, Nigeria. *Global Res J Sci*. 2011;1:98-103.
12. Menendez C, Ordi J, Ismail MR, Ventura PJ, Aponte JJ, Kahigwa E, Font F, Alonso PL. The impact of placental malaria on gestational age and birth weight. *The Journal of infectious diseases*. 2000;181(5):1740-5.
13. Stephenson LS, Latham MC, Ottesen EA. Malnutrition and parasitic helminth infections. *Parasitology*. 2000;121(S1):S23-38.
14. Hotez PJ, Bethony J, Bottazzi ME, Brooker S, Buss P. Hookworm: "the great infection of mankind". *PLoS Medicine*. 2005;2(3):e67.
15. Al-Delaimy AK, Al-Mekhlafi HM, Nasr NA, Sady H, Atroosh WM, Nashiry M, Anuar TS, Moktar N, Lim YA, Mahmud R. Epidemiology of intestinal polyparasitism among Orang Asli school children in rural Malaysia. *PLoS Neglected Tropical Diseases*. 2014 Aug 21;8(8):e3074.
16. Department of Geography, Benue State University, Makurdi
17. Egwunyenga, A.O., Ajayi, J.A., Nmorsi, O.P, Duhlińska-Popova DD. Plasmodium/intestinal helminth coinfections among pregnant Nigerian women. *Memórias do Instituto Oswaldo Cruz*. 2001;96(8):1055–1059.
18. Wilson S, Vennervald BJ, Kadzo H. Hepatosplenomegaly in Kenyan schoolchildren: exacerbation by concurrent chronic exposure to malaria and *Schistosoma mansoni* infection. *Tropical Medicine and International Health*, 2007;12(12):1442–1449.

19. Beena, J. Sanjay R, Grover, PS. Prevalence of intestinal parasites among patients in Ambala city, Haryana, India. International Journal of Research and Medical Science. 2015;3(12):2753-3758.
20. Omudu EA, Onah E, Emberga-Biem M. Epidemiological studies of polyparasitism in two different communities in Benue State, Nigeria. Journal of King Abdulaziz University. Medical Sciences. 2012 Jan 4;19(1):21-34.
21. Cheesbrough M. District laboratory practice in tropical countries, part 2. Cambridge university press; 2005.
22. Akor, JO, Omudu, EA, Obisike, V.U, Imandeh, G.N. The prevalence of polyparasitism in Oju, Local Government Area of Benue State, Nigeria. International Journal of Tropical Disease and Health. 2019;39(2):1-6.
23. Alaku IA, Omudu EA, Imande NG, Akor JO. Epidemiology of intestinal polyparasitism among primary school pupils in Awe, Awe Local Government Area, Nasarawa State, Nigeria. South Asian Journal of Parasitology. 2019;11:1-6.

© 2022 Uweh and Omudu; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/89902>