

# Synthesis, Characterization and Biological Activity of New Amides Containing Azo Moieties

Elham Abdalrahem Bin Selim <sup>a</sup>, Mohammed Hadi Al-Douh <sup>a\*</sup>,  
Samia Mabrook Bin Mahram <sup>a</sup>, Wedad Mohammed Al-Haik <sup>b</sup>  
and Khalid Saeed Sonbol <sup>b,c</sup>

<sup>a</sup> Chemistry Department, Faculty of Science, Hadhramout University, Mukalla, Hadhramout, Yemen.

<sup>b</sup> Department of Biology, Faculty of Science, Hadhramout University, Mukalla, Hadhramout, Yemen.

<sup>c</sup> Alsalam Medical Specialized Center, Mukalla, Hadhramout, Yemen.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/AJACR/2022/v11i230250

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/85522>

Received 10 December 2021

Accepted 14 February 2022

Published 15 February 2022

Original Research Article

## ABSTRACT

Three new amide compounds **6**, **8** and **9** were derived from diazo dye **4**. 4-Hydroxy-3-methoxy-5-(2,6-dinitrophenyl) benzaldehyde **4** was reacted separately with compounds **5** and **7** to obtain **6** and **8**, respectively. The amide compound **9** was synthesized by two different methods. The reaction between **6** and a diacyl chloride compound **7** gave compound **9**. Also, compound **9** was produced via the reaction of compound **8** and an *N*-terminus compound **5**. All compounds were characterized by their melting points, UV-Vis, and FTIR spectra. Moreover, the mass spectrum and elemental analysis of compound **9** were determined. All three synthesized compounds were tested against *P. mirabilis*, *E. coli*, and *S. aureus* at different concentrations and showed significant results.


**Keywords:** Synthesis; amide; vanillin; azo functional group; UV-Vis.; FTIR; MS.

## 1. INTRODUCTION

One of the most important functional groups in organic chemistry and in nature is amide group.

This group is occurred in a wide variety of compounds. Proteins and enzymes have amide bonding structure. Medicinally, the amide group is one of the common groups in drug synthesis.

\*Corresponding author: Email: m.aldouh@hu.edu.ye, mhd\_douh@yahoo.com;



**Scheme 1. The new synthesized of amide 9**

About a quarter of available drugs contain at least one amide bond [1-3]. The amide compounds are thermally stable [4]. Amide bonds are also present in huge molecules. The supramolecule compounds arises from the binding of a group of molecules of a well-defined structure and grouped together through non covalent interactions such as hydrogen bonds, dipole-dipole, van der Waals forces,  $\pi$ - $\pi$ , and coordination bonds [5].

In recent years, supramolecular of aromatic systems have attracted attention of many researchers [6-9]. The diazo dye derivatives have undergone extensive studies in several fields, and exhibited an important applications such as dyes, biomedical studies, organic synthesis, biological activities including antineoplastics, antidiabetics, antiseptics, anti-inflammatory and other useful chemotherapeutics agents [10-15].

The supramolecular arrangement of azo dyes is based on non-covalent supramolecular interactions which keep packaged the whole systems. Also, it has raised particular interest in this chemical system [16]. Since the amide group, diazo dye, and supramolecular compounds are an important and have many applications. This is the motivation in our present

study to synthesis three new amides derived from diazo dye and have a high molecular weight, as well as evaluate their biological activities against three types of bacteria.

Recently, we have been synthesized and characterized a new dinitro diazo dye **4**, Scheme 1 [11]. Additionally, the dipole interaction and the intramolecular hydrogen bonds of **4** are studied [12]. Based on previous information, we were interested to synthesize a new amide **9**, which produced from the reaction between compound **4** with compounds **5** and **7** by two pathways, Scheme 1. All synthesized compounds were assigned using melting points, FTIR and UV-Vis spectroscopies. However, the microbial inhibitory of these derivatives against some pathogenic bacteria such as *Staphylococcus aureus* as gram +ve, *Escherichia coli* and *Proteus mirabilis* as gram -ve were tested.

## 2. MATERIALS AND METHODS

### 2.1 General

The melting points were determined using Electrothermal Melting Point Apparatus. FTIR spectra were performed using JASCO FTIR 4600, single beam, path Laser, by KBr disk technique in the frequency range of 4000-400

cm<sup>-1</sup>. UV–Vis spectra of the compounds were recorded with JASCO V–750 Spectrophotometer in ethanol at the wavelength range of 900–200 nm. Mass spectra were carried out on Direct Inlet part to mass analyzer in Thermo Scientific GC/MS model ISQ, and Elemental analyses were done using FLASH 2000 CHNS/O analyzer, Thermo Scientific.

## 2.2 Synthesis

Compound **4** was synthesized according to our earlier published procedure [11]. Both compounds 1,6-*N,N*-di(6-aminohexyl)hexan diamide **5** [17] and diacyl chloride **7** [18] were prepared and characterized previously.

### 2.2.1 Synthesis of compounds 6

It was synthesized according to the method that mentioned in [19-21] with some modification. An equimolar amounts from **4** (3.50 g, 0.01 mol) and **5** (3.0 g, 0.01 mol) were dissolved in 15 mL absolute ethanol EtOH containing a drop of AcOH. The reaction mixture was refluxed on a water bath for 1 hr, after that allowed to cool at room temperature. The resulting solid was filtered off, washed with a little 2% HCl solution and distilled water.

### 2.2.2 Synthesis of compounds 8

It was obtained as a method that described previously in the literature [22], with some modification. A (3.50 g, 0.01 mol) of **4**, (4.10 g, 0.01 mol) of **7**, and (7.45 g, 0.054 mol) of anhydrous K<sub>2</sub>CO<sub>3</sub> in 70 mL ethanol 96% were refluxed for 6 hrs. The resulting solution was cooled at room temperature and poured into about 300 mL of cold water. The solid material immediately was formed, filtered off, washed several times with cold water, and dried.

### 2.2.3 Synthesis of compounds 9

#### 2.2.3.1 Pathway A

About (0.40 g, 0.001 mol) of **6**, (0.30 g, 0.001 mol) of **7**, and anhydrous K<sub>2</sub>CO<sub>3</sub> (7.45 g, 0.054 mol) with 70 mL ethanol 96% were refluxed for 6 hrs, left at room temperature then poured into about 300 mL cold water. A deep green solid was formed, filtered, washed several times with cold water and dried. Yield: 21.6%, 0.14 g, and

m.p. 150°C. CHN elemental analysis: Found (calculated) for C<sub>64</sub>H<sub>82</sub>N<sub>14</sub>O<sub>18</sub>: C: 57.44(60.68), H: 6.45(6.54), N: 14.72(14.68).

#### 2.2.3.2 Pathway B

About (0.50 g, 0.0005 mol) of **8** and (0.15 g, 0.0004 mol) of **5** were dissolved in 15 mL of absolute ethanol containing a drop of AcOH. The reaction mixture was refluxed on a water bath for 1 hr, then left to cool at room temperature. The resulting solid was filtered, washed with a little 2% HCl, distilled water, and left to dry. Yield: 21.6%, 0.14 g, m.p. 150°C. CHN elemental analysis: Found (calculated) for C<sub>64</sub>H<sub>82</sub>N<sub>14</sub>O<sub>18</sub>: C: 57.38(60.68), H: 6.41(6.54), N: 14.57(14.68). The physical properties of all synthesized compounds were recorded in Table 1.

## 2.3 Methodology of Antimicrobial Activity


The bacteria strains that used in this study according to our earlier published procedure [11]. Three bacteria strains *Staphylococcus aureus*, *Escherichia coli* and *Proteus mirabilis* were isolated from different patients attending Alsalam Medical laboratory. The compounds **6**, **8**, and **9** were added to the growth medium in different concentrations, Table 3. The minimum inhibitory concentration MIC of **4**, **6**, **8** and **9** were determined by using a different dilution for those compounds, which diluted with sterile ethanol 75%. It was weighed 0.07 g from **4**, 0.21 g from **6**, 0.20 g from **8** and 0.27 g from **9**, separately, in 100 mL of ethanol 75%. The following concentration was prepared (1:2, 1:4, 1:8, 1:16 and 1:32) from the stock of each solution [23-24]. The antibiotic susceptibility tests were carried out by the Kirby–Bauer disk diffusion technique according to Clinical Laboratory Standard Institute guidelines [25-26]. Bacterial suspension for each strain was tested as described by McFarland [27]. The test medium was used Mueller–Hinton agar [28]. All prepared plates were incubated at 37°C for 24 hrs [29].

The Discs of Ciprofloxacin Cip, Amikaci Ami, Clarithromycin Cit, Cefaclor Cef, Cefepime Cep, Azithromycin Azt and Co-trimoxazole Ctm were used as a reference for evaluation of antibacterial [30]. By this manner, some of the synthesized compounds were screened *in vitro* for their antibacterial activity against *S. aureus*, *E. coli* and *P. mirabilis*.

### 3. RESULTS AND DISCUSSION

Compound **9** in Scheme 1 was synthesized by two pathways. Its physical properties such as elemental analysis, melting points, and yields etc. were tabulated in Table 1. The CHN analyses showed a good agreement with suggested molecular formula  $C_{64}H_{82}N_{14}O_{18}$ . Compound **4** was early synthesized by coupling reaction between 2,6-dinitroaniline **1** and 4-hydroxy-3-methoxybenzaldehyde **3** as shown in Scheme 2 [11].


In the first pathway, the amide **9** was synthesized by reaction of diazo dye **4** with *N*-terminus diamine **5** by refluxing in absolute ethanol using a drop of acetic acid to yield **6**. After that, compound **6** was treated with diacyl chloride **7** to obtain amide **9**, Scheme 3a. The amide **9** was obtained also by the second different pathway, in which the diazo dye **4** treated with a diacyl chloride compound **7** and refluxing in 96% ethanol to give compound **8**, then the resulted **8** was reacted with *N*-terminus diamine **5** to obtain **9**, Scheme 3b. The physical data of compounds **4**, **6**, and **9** were tabulated in Table 1.



Scheme 2. Synthesis of 4-hydroxy-3-methoxy-5-(2,6-dinitrophenylazo) benzaldehyde **4**

Table 1. Physical properties of compounds **4**, **6**, **8** and **9**

Physical properties	Compounds			
	<b>4</b>	<b>6</b>	<b>8</b>	<b>9</b>
Molecular formula	$C_{14}H_{10}O_7N_4$	$C_{46}H_{54}O_{14}N_{12}$	$C_{46}H_{48}O_{18}N_{10}$	$C_{64}H_{82}N_{14}O_{18}$
Molecular weight (g/mol)	346.25	998.99	1028.93	1335.42
Melting point ( $^{\circ}C$ )	88	191	153.4	150
Color	Light orange	Yellowish brown	Brown	Deep green
Yield %	46.2	27	6.5	21.6
CHN elemental analysis	-	-	-	<u>Pathway A</u>
Found (calculated)				C: 57.44(60.68), H: 6.45(6.54), N: 14.72(14.68)
				<u>Pathway B</u>
				C: 57.38(60.68), H: 6.41(6.54), N: 14.57(14.68)



**Scheme 3. The pathways for synthesis of 9.**

### 3.1 FTIR Spectroscopy

The compound **6** was characterized by infrared spectroscopy, Table 2. The important infrared bands of **6** compared to the FTIR of **4** were the appearance of strong band at  $1658\text{ cm}^{-1}$  which was attributed to the imine functional group  $\nu\text{ C=N}$ , Fig. 1a and 1b, accompanied by disappearance of aldehydic hydrogen stretching frequencies, as well as the two spikes of primary amine. The FTIR spectrum of **8** displayed a new band at  $1765\text{ cm}^{-1}$  corresponding to an ester functional group stretching frequency as well as disappearing of the  $\nu\text{ OH}$  band as comparing with the FTIR spectrum of diazo dye **4**, Table 2, Fig. 1c.

The comparison of the FTIR spectra of the compounds **6** and **8** as well as the amide spectrum. Table 2 gives important information about the suggested structure of **9**. The FTIR spectrum of compound **9** synthesized from the first pathway, Fig. 1d. Table 2 characterized by the observance of the band at  $1787\text{ cm}^{-1}$  indicated an ester functional group and disappearance of  $\nu\text{ OH}$  stretching frequency as

compared to the FTIR spectrum of **6**. In contrast, the FTIR spectrum of **9** produced from the second pathway characterized by the absorption band at  $1619\text{ cm}^{-1}$  assigned for imine functional group as well as the absence of two spikes of primary amine that comparing with the FTIR spectrum of **8**.

### 3.2 UV-VIS. Spectroscopy

The UV-Vis of **6** exhibited a high intense absorption peak at 390-477 nm assigned to  $\pi \rightarrow \pi^*$  electronic transition of the azo group  $-\text{N}=\text{N}-$ , the spectrum also contained a shoulder significant peak at the range of 360-340 nm which attributed to  $n \rightarrow \pi^*$  transition of the  $\text{C=N}$ -imine group [31].

The UV-Vis spectrum of **8**, showed bands at 391.2 nm and 230 nm indicated of  $\pi \rightarrow \pi^*$  electronic transitions, and 442.8 nm assigned for  $n \rightarrow \pi^*$  transitions. While the UV-Vis spectrum of **9** was recorded in ethanol and exhibited bands at 390 nm and 230 nm that assigned for  $\pi \rightarrow \pi^*$  electronic transitions, and 445 nm, 340.4 nm assigned for  $n \rightarrow \pi^*$  electronic transitions, Fig. 2.





Fig. 1. The FTIR spectra of compounds a. 4, b. 6, c. 8, and d. 9

**Table 2. FTIR spectral data of compounds 4, 6, 8 and 9 (cm<sup>-1</sup>) in KBr**

Functional Group	Compounds			
	4	6	8	9
$\nu$ -OH arom.	3277	3111	-	-
C=O aldehyde	1672	-	1720	-
$\nu$ C=C	1589	1494	1494	1549
$\nu$ CH <sub>3</sub> aliph.	as 2976 sy 2858	as 1694 sy 1590	as 2948 sy 2366	as 2925 sy 2360
$\nu$ -N=N-	as 1589 sy 1430	as 1511 sy 1465	as 1598 sy 1403	as 1604 sy 1503
$\delta$ CH <sub>3</sub>	as 1454 sy 1372	as 1428 sy 1345	as 1403 sy 1362	as 1504 sy 1403
$\nu$ -NO <sub>2</sub>	as 1508 sy 1301	as 1428 sy 1345	as 1530 sy 1463	as 1530 sy 1400
$\nu$ C-O-C aliph.	as 1266 sy 1153	as 1267 sy 1141	as 1268 sy 1113	as 1255 sy 1114
$\nu$ C-N-C	-	1027	-	1004
$\nu$ C=O amide	-	1672	1615	1681
$\nu$ C=N	-	1590	-	1619
$\nu$ C=O arom. ester	-	-	1765	1787

**Fig. 2. The UV-Vis spectra of compounds a. 4, b. 6, c. 8, and d. 9**

### 3.3 Mass Spectrum

The molecular weight of the suggested structure compound  $m/z$  1335.42 was outside the range of the used MS spectrometer. The mass spectrum of **9** showed several characteristic peaks that confirmed the suggested structure, Fig. 3. The cleavage of **9** gave the charged fragment **10** corresponded to the  $m/z$  343.94 in the spectrum and free radical fragment **11**, Fig. 4a. Also, the peak at  $m/z$  850.72 (calculated 851.43) corresponded to the side of the broken bonds of cation fragment **11**, which indicated the loss of C<sub>7</sub>H<sub>13</sub>NO and methylene group free radical fragments from it, Fig. 4b. However, the mass spectrum of **9** exhibited peak at  $m/z$  807.36 of **13**

(calculated 807.37) corresponded to the loss of two hydrogen and butyl group free radicals from charged fragment **12** as shown in Fig. 4c.

### 3.4 Microbiology Activity

The synthesized compounds **6**, **8**, and **9** are showed most sensitive to many kinds of bacteria due to the type of the functional groups of these compounds, which had an increasing effect on the levels of toxicity of bacterial species [11]. In fact, the imine compounds are very important in industrial application and in biological activities. They have found a good effect to the pharmacological activities such as antimicrobial [32].




Fig. 3. The Mass Spectrum of new synthesized amide 9




Fig. 4. The fragments a. 10 and 11, b. 12, and c. 13 of compound 9



The results of antibacterial activity of these compounds against *Staphylococcus aureus*, *Escherichia coli* and *Proteus mirabilis* are showed in Table 3. Compound **6** showed active of *P. mirabilis* at the concentrations 2.0 mM [+ = IZ 7–13 mm], 1.0 mM [+ = IZ 7–13 mm], 0.25 mM [+ = IZ 7–13 mm], 0.125 mM [+ = IZ 7–13 mm] and 0.0625 mM [+ = IZ 7–13 mm].

On other hand, compound **8** was effective against two types of pathogenic bacteria, *E. coli* at concentration 0.25 mM [+ = IZ 7–13 mm] and 0.125 mM [+ = IZ 7–13 mm]. Also, *P. mirabilis* at concentrations 2.0 mM [+ = IZ 7–13 mm], 1.0 mM [+ = IZ 7–13 mm], 0.25 mM [+ = IZ 7–13 mm], 0.125 mM [+ = IZ 7–13 mm] and 0.0625 mM [+ = IZ 7–13 mm].

**Table 3. Inhibitory growth activity of compounds 6, 8 and 9 against pathogenic bacteria**

Compound	Conc. mM	microorganisms		
		gram +ve <i>S. aureus</i>	gram -ve <i>E. coli</i>	<i>P. mirabilis</i>
<b>6</b>	2.0	–	–	+
	1.0	–	–	+
	0.5	–	–	+
	0.25	–	–	–
	0.125	–	–	+
	0.0625	–	–	+
<b>8</b>	2.0	–	–	+
	1.0	–	–	+
	0.5	–	–	–
	0.25	–	+	+
	0.125	–	+	+
	0.0625	–	–	+
<b>9</b>	2.0	–	–	–
	1.0	–	–	–
	0.5	–	+	–
	0.25	+	–	–
	0.125	–	–	–
	0.0625	–	–	–
Ethanol 75%	–	–	–	–
Cip	5*	+++	–	–
Ami	30*	+++	–	–
Cit	15*	+++	–	–
Cef	30*	–	++	+++
Cep	30*	–	++	++
Azt	15*	–	+	–
Ctm	25*	–	+++	+++

Key to symbols: \*ug, Cip: Ciprofloxacin, Ami: Amikaci, Cit: Clarithromycin, Cef: Cefaclor, Cep: Cefepime, Azt: Azithromycin and Ctm: Co-trimoxazole as antibacterial, Disc diameter = 6 mm, Highly active: +++ [IZ > 19 mm], Moderately active: ++ [IZ 13-19 mm], Slightly active: + [IZ 7-13 mm], Inactive: – [IZ < 7 mm]

The amide **9** showed no activated on *P. mirabilis* with all concentrations. It exhibited the most sensitive for *S. aureus* at concentration 0.25 mM [+ = IZ 7–13 mm] and *E. coli* at concentration 0.5 mM [+ = IZ 7–13 mm]. However, amide **9** is inactive of other concentrations that indicated all bacteria tested is resistance of this compound, as shown in Table 3. A similar result is previously reported by Al-Shara'ey *et al.* They have been reported that most bacteria tested are resistance to some synthesized polycyclic compounds [33-34].

Seven antibiotics were used in this study that the highest percentage of antibiotics active was to Ciprofloxacin, Amikaci and Clarithromycin to *S. aureus*, while inactive to both *E. coli* and *P. mirabilis*. Each of Cefaclor, Cefepime, Azithromycin and Co-trimoxazole were shown more active to *E. coli* and *P. mirabilis* except *S. aureus*. The studies indicated to better understand virulence and multidrug resistance, revealed that most of the antibiotic resistance genes and many of the virulence genes reside on mobile genetic elements, such as plasmids,

transposons, and prophages, indicating substantial horizontal gene transfer from other bacteria [35]. We believe that the functional groups of these compounds are an increasing effect on the levels of toxicity of bacterial species.

#### 4. CONCLUSION

In this paper, we have synthesized three new amide compounds **6**, **8**, and **9**. Compound **6** has been synthesized *via* the condensation reaction between **4** and **5**. Compound **8** has been obtained through the nucleophilic substitution reaction between **4** and **7**. Both compounds **6** and **8** can be used to synthesize **9** through two different pathways. Compound **9** can be obtained either by the reaction of **6** and **7**, or by the reaction of **8** with **5**. All synthesized compounds have been characterized by spectroscopic UV-Vis, and FTIR technique, as well as elemental analysis CHN and mass spectrometry of **9**. The antibacterial activity has been evaluated by using minimum inhibitory concentration MIC method. The results showed that **6** exhibited a high sensitivity for *P. mirabilis* at the most concentrations, while **8** had positive results for *E. coli* and *P. mirabilis* at many concentrations. Compound **9** displayed antibacterial activity against *S. aureus* at the concentration 0.25 mM, and against *E. coli* at 0.5 mM.

#### ACKNOWLEDGMENT

We thank Mrs. Gehan Abdullah Balala, Dr. Ali Abdullah Saeed, and Mr. Naef Abdulrahman Al-Kalali for assistance in analyses.

#### DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- Seavill PW, Wilden JD. The preparation and applications of amides using electrosynthesis. *Green Chem.* 2020;22:7737-7759. DOI: 10.1039/D0GC02976A
- Roughley SD, Jordan AM. The medicinal chemist's toolbox: An analysis of reactions used in the pursuit of drug candidates. *J Med Chem.* 2011;54(10):3451-3479. DOI: 10.1021/jm200187y
- Wang X. Challenges and outlook for catalytic direct amidation reactions. *Nat Catal.* 2019;2:98-102. DOI: 10.1038/s41929-018-0215-1
- Rostami E, Bagherzadeh M, Alinassab T, Mohammadpour M, Zangoeei M, Feraidooni M, Tavazo F, Keshavarz Z. Synthesis and characterization of new organo soluble and thermally stable poly(thioether-amide)s bearing pyridine subunit in the main chain. *Polymer Sci Intern Scholarly Res Notices.* 2014;2014(531754):1-7. DOI: 10.1155/2014/531754
- Biedermann F, Schneider HJ. Experimental binding energies in supramolecular complexes. *Chem Rev.* 2016;116(9):5216-5300. DOI: 10.1021/acs.chemrev.5b00583
- Breneman CM, Wiberg KB. Determining atom-centered monopoles from molecular electrostatic potentials. The need for high sampling density in formamide conformational analysis. *J Comput Chem.* 1990;11(3):361-373. DOI: 10.1002/jcc.540110311
- Kennedy AR, Hughes MP, Monaghan ML, Staunton E, Teat SJ, Smith WE. Supramolecular Motifs in *s*-Block Metal Bound Sulfonated Monoazo Dyes. *J Chem Soc Dalton Trans.* 2001;14:2199-2205. DOI: 10.1039/b102106k
- Desiraju GR. Perspectives in supramolecular chemistry. *Crystal Design: Structure and Function.* vol. 7, John Wiley & Sons, Ltd., England; 2003, 418. DOI: 10.1002/0470868015
- Hoeben FJM, Jonkheijm P, Meijer EW, Schenning APH. About supramolecular assemblies of  $\pi$ -conjugated systems. *J Chem Rev.* 2000;105(4):1491-1546. DOI: 10.1021/cr030070z
- Pagariya SK, Pathade RM, Bodkhe PS. Synthesis, Characterization and

- Antimicrobial Screening of Some Azo Compounds Derived from Ethyl Vanillin. *Res J Chem Sci.*2015;5(7):20-28. Available:<http://www.isca.in/rjcs/Archives/v5/i7/4.ISCA-RJCS-2015-088.pdf>
11. Al-Douh MH, Selim EAB, Al-Haik WM, Mahram SMB, Al-Bakri AK, Al-Nohey DS, Sonbol KS. Synthesis and antimicrobial activity of some compounds derived from diazo dyes. *Intern J Chem Stud.* 2019;3(4):6-12. ISSN: 2581-348X. Available:<http://www.chemistryjournal.in/archives/2019/vol3/issue4/3-3-16>
  12. Selim EAB, Al-Douh MH. Unusual effects of intramolecular hydrogen bonds and dipole interaction on FTIR and NMR of some imines. *Asian J Appl Chem Res.* 2019;3(3):1-6. DOI: 10.9734/AJACR/2019/v3i330096
  13. Al-Douh MH, Selim EAB, Tahir EM, Esmail SAA, Naji YA, Abdullah HH. Biophysical study for the interactions of some diazo dyes with human serum albumin. *Asian J Appl Chem Res.* 2019;4(3):1-10. DOI: 10.9734/AJACR/2019/v4i330113
  14. Selim EAB, Al-Douh MH, Al-Nohey DS, Al-Haik WM, Sonbol KS. Synthesis, Characterization, and Antimicrobial Activity Studies of New Schiff Base Tetradentate Ligands and Their Manganese (II) Complexes. *Asian J Chem Sci.* 2020;7(4):35-46. DOI: 10.9734/AJOCS/2020/v7i419030
  15. Al-Douh MH, Selim EAB, Abdullah HH, Abdullah HY, Al-Bakri AK, Al-Nohey DS, Mahram SMB, Yusr, KM, Hawiel, FAB. Molecular docking study of some nitro diazo dye derivatives as antiviral candidates of COVID-19. *Asian J Appl Chem Res.* 2021;8(8):8-15. DOI: 10.9734/AJACR/2021/v8i230187
  16. Garcia HC, Ferreira GR, de Oliveira LFC. An interesting coordination complex formed between the azo dye sudan red g and cobalt ion. *J Mol Struct.* 2014;1061:41-46. DOI: 10.1016/j.molstruc.2013.12.078
  17. Brydson JA. *Plastics materials*, 7th ed. Butterworth-Heinemann: UK; 1999. ISBN: 978-0-7506-4132-6
  18. Guha PC, Sankaran DK. *Muconic Acid. Org Synth.* 1964;26:57-61 DOI: 10.15227/orgasyn.026.0057
  19. Al-Douh MH, Al-Fatlawy AA, Abid OH. Synthesis and characterization of some 2-(*N*-Benzoyl-*N*-pyrid-4-yl aminobenzyl)-aminobarbituric Acids *via* Schiff's Bases. *Hadh Stud Res.* 2003;4(2):37-49. ISBN-National Library Aden 18-2004
  20. Al-Douh MH, Al-Fatlawy AA, Abid OH. Synthesis and characterization of 2-(*N*-Benzoyl-*N*-pyrid-3-yl aminobenzyl)-amino barbituric acids *via N*-Benzylidene Pyridine-3-amines. *Fac Sci Bull.* 2003;16(3):83-94. ISSN: 1684-100X
  21. Al-Douh MH, Al-Fatlawy AA, Abid OH. Synthesis and characterization of some 2-(*N*-Benzoyl-*N*-pyrid-2-yl aminobenzyl)-aminobarbituric Acids *via N*-Benzylidene Pyridine-2-amines. *Univ Aden J Nat Appl Sci.* 2004;8(1):181-194. ISSN: 1606-8947
  22. Samad MK, Chawishli LH, Hussein AJ. Synthesis and spectroscopic characterization of some diazodibenzyloxy pyrazolines from some diazodibenzyloxy chalcones. *Zanco J Pure Appl Sci.* 2015;27(2):53-60. DOI: 10.21271/zjpas.v27i2.148
  23. Othman AS. Antibacterial activity of bee and yemeni sidr honey against some pathogenic bacterial species. *Inter J Curr Microbiol Appl Sci.* 2014;3(10):1015-1025. ISSN: 2319-7706
  24. Andrews JM. Determination of minimum inhibitory concentrations. *J Anitmicrob Chemother.* 2001;48(S1):5-16. DOI: 10.1093/jac/48.suppl\_1.5
  25. Piddock LJV. Techniques used for the determination of antimicrobial resistance and sensitivity in bacteria. *J Appl Bacteriology.* 1990;68(4):307-318. DOI: 10.1111/j.1365-2672.1990.tb02880.x
  26. CLSI M02-A10. Performance Standards for Antimicrobial Disk Susceptibility Tests; Approved Standard. 10th Ed. Clinical and Laboratory Standards Institute: Wayne, Pennsylvania, USA; 2010;29(3). ISBN 1-56238-716-2
  27. McFarland J. The Nephelometer: An instrument for estimating the number of bacteria in suspensions used for calculating the opsonic index and for vaccines. *J Am Med Assoc.* 1907;49(14):1176-1178. DOI:10.1001/jama.1907.25320140022001f
  28. Rajput A, Prajapati B, Chauhan B, Shah A, Trivedi T, Kadam M. Prevalence of metallo-beta lactamases (MBL) producing *Pseudomonas aeruginosa* in a tertiary care

- hospital. Indian J Basic & Appl Med Res. 2012;1(4):304-308.  
Available:<https://www.ijbamr.com/assets/images/issues/pdf/September%202012%20%20304-308%20%20ZZ.pdf>
29. Joshi S, Pawar V, Uma V. Synthesis, characterization and biological studies of schiff bases metal complexes Co (II), Zn (II), Ni (II), and Mn (II) derived from amoxicillin trihydrate with various aldehydes. Intern J Pharma and Bio Sciences. 2011;2(1):170-178  
Available:<https://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.294.1231&rep=rep1&type=pdf>
30. Lundstrom J. Simple Isoquinoline Alkaloids. In: Brossi, A, editor. The Alkaloids: Chemistry and pharmacology. Academic Press, Inc.: New York, USA; 1983;21:255.  
DOI: 10.1016/S0099-9598(08)60052-8
31. Unver H. Synthesis and spectroscopic studies in some new schiff bases. Spectrosc Lett. 2001;34(6):783-791.  
DOI: 10.1081/SL-100107900
32. Kratky M, Sova JV, Volkova M, Buchta V, Trejtnar FS, Rikova J. Antimicrobial activity of sulfonamides containing 5-Chloro-2-hydroxybenzaldehyde and 5-Chloro-2-hydroxybenzoic acid scaffold. Eur J Med Chem. 2012;50(4):433-440.  
DOI: 10.1016/j.ejmech.2012.01.060
33. Al-Shara'ey AA, Al-Douh MH, Gahnem YS. Synthesis and antimicrobial study of some polycyclic compounds by [3+2] cycloaddition reactions of aldazines with N-Aryl maleimides. TOJSAT. 2019;9(1):80-90.  
Available in:  
<http://www.tojsat.net/journals/tojsat/articles/v09i01/v09i01-15.pdf>
34. Al-Shara'ey AA, Al-Douh MH, Thabet EA, Gahnem YS. Synthesis of new heterocyclic compounds from reaction of N-aryl maleimides with cinnamal aryl hydrazones and their antimicrobial activity. Inter J Chem Stud. 2019;3(3):36-45.  
Available:<https://www.chemistryjournal.in/download/100/3-3-12-158.pdf>
35. Carboneau C, Bengé E, Jaco MT, Robinson M. A lean six sigma team increases hand hygiene compliance and reduces hospital-acquired MRSA infections by 51%. J Health Quality. 2010;32(4):61-70.  
DOI: 10.1111/j.1945-1474.2009.00074.x

© 2022 Selim et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/85522>