



## **Study of Low Back Pain in Obese Patients**

**Rahul Agola<sup>a</sup> and Nareshkumar Dhaniwala<sup>a</sup>**

<sup>a</sup> Department of Orthopedics, Jawaharlal Nehru Medical College, Sawangi (Meghe) and Acharya Vinoba Bhave Rural Hospital Wardha, India.

### **Authors' contributions**

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

### **Article Information**

DOI: 10.9734/JPRI/2022/v34i22A35855

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/82504>

**Study Protocol**

**Received 06 December 2021**  
**Accepted 11 February 2022**  
**Published 11 March 2022**

## **ABSTRACT**

Low back pain, a very common presenting complaint in orthopaedic Outpatient department is caused by various factors such as strenuous physical work, Poor Posture, Trauma, Age related degeneration of vertebral joints and various diseases of ligaments and bones. Overweight and Obesity cause changes in Posture and biomechanics of lumbar spine function and weight transmission. Relationship between low back pain and Obesity requires study to learn about its causation and its effect. This study is an attempt in this direction by measuring the parameters of Body mass Index and its effects on Lumbosacral angles.

**Keywords:** Low back pain; overweight; obesity; lumbosacral angles.

## **1. INTRODUCTION**

Overweight and obesity is a serious community health problem that contributes to the global trouble of chronic disease as a pandemic condition [1]. Hypertension, Diabetes and Obesity have been considered as predisposing factors to diseases like Gastro-oesophageal reflux, Obstructive sleep apnoea (OSA), certain

types of cancers like breast cancer(post-menopausal), endometrial cancer, Gastro intestinal track related cancer etc [2].

Obesity causes changes in normal function and biomechanics of the vertebral column and big joints of limbs [3,4]. Study of curvature and angles of Lumbar and sacral spine helps to understand transmission of upper body weight to

<sup>≡</sup> M.B.B.S., Junior resident in M.S. Orthopedics;

<sup>°</sup> M.B.B.S., M.S. Orthopedics, Professor;

\*Corresponding author: E-mail: rahul.agola.ra@gmail.com, drrahulagola@gmail.com;

the trunk and lower limbs. Lumbosacral angles could be affected by various factors such as race, age, posture, diseases of spine and surgery on the spine [5,6]. Due to Obesity the lumbosacral spine undergoes changes in its alignment causing chronic back pain [7,8]. Complaint of low back pain is one of the most common complaints in the Outpatient department in Orthopaedics.

Various studies have been done to correlate obesity to chronic nonspecific back pain in adult population [7]. The objectives of the present study, titled "Study of Low back pain in Obese patients", being done in tertiary care rural hospital, are to compare the lumbosacral angles in overweight and obese people with non-obese persons and to analyse the relation of Body Mass Index on Lumbosacral angles.

## 2. MATERIALS AND METHODS

The present Study type is Prospective Cross-sectional study. The Study period is from November 2020 to October 2022. Study sample size is 100 each for case and control, as calculated by Sample Size formula with desired error of margin:

$$n = \frac{2x^2/2 \cdot P \cdot (1 - P)}{d^2}$$

Where,

$2x^2$  is the level of significance at 5% level of significance i.e. 95% confidence interval = 1.96

P = Prevalence of low back pain

D = Desired error of margin

The inclusion criteria are patients between 20 to 60 years of age who complain of nonspecific chronic low back pain (LBP) without any radiation to lower limbs and without any history of neurologic, respiratory, musculoskeletal or lower limb pathology.

Patients having spinal deformity such as scoliosis or kyphosis and or suffering from Ankylosing spondylitis, severe Osteoporosis, Progressive Neurologic disease or Vertebral infection are the exclusion criteria for this study.

### 2.1 Methods

Adult patients of non-specific chronic LBP (case group) consenting to participate in the study will be enrolled. The demographic details of the patients and the details of complaints will be

noted. Physical examination including a detailed examination of spine in terms of inspection, palpation, movement and alignment will be done. X ray lumbosacral spine AP and Lateral view will be taken.

Body mass index (BMI) will be measured from weight and height of the patient. The various angle in lumbosacral spine will be measured as shown below. Corelation between BMI and lumbosacral angles will be found out.

Similar study on normal adults (control group) without any complaints of LBP will be carried out simultaneously. Medical students, Post-graduate residents and staff of the hospital will be encouraged to participate as control subjects.

Comparison between patients of LBP and control group will be done and suitable conclusions drawn.

BMI is measured by weight of person in Kg divided by height of patient in meter square [9,10]. A person is labelled underweight if BMI is below 18.5. BMI of 18.5 to 24.9 is said to be the normal range. Person is called Overweight if BMI between 25.0 to 29.9. Obesity is divided in three classes with Class I (BMI – 30.0 to 34.9), Class II (BMI – 35.0 to 39.9) and Class III (If BMI is more than 40).

The following angles will be measured to asses relation between obesity and lumbosacral angles:

1. lumbosacral angle (LSA): A
2. sacral inclination angle (SIA): B
3. lumbosacral disc angle (LSDA): C
4. lumbar lordosis angle (LLA): D

Age, posture, race, disease, and surgery can affect these angles between the lumbar spine and sacrum.

Lumbosacral angle (LSA) is the angle measured between upper border of 1<sup>st</sup> sacral vertebra and the horizontal plane. Sacral inclination angle (SIA) is measured between Vertical line and a line along the posterior border of S1 vertebra. Lumbosacral disc angle (LSDA) is the angle measured between superior end plate of S1 and the inferior end plat of L5 intervertebral disc. Lumbar lordosis angle (LLA) is angle formed by intersection of two perpendiculars on the lines drawn through superior end plate of L1 and inferior end plat of L5.




Fig. 1. A, B, C and D are showing lumbo-sacral spine's geometric angles and how to measure it

### 3. CONCLUSION

The study will compare the above angles in relation to BMI in case and control subjects and suitable conclusion will be drawn. The Statistical analysis will be done using SPSS version 17.0.

### CONSENT AND ETHICAL APPROVAL

As per university standard guideline, participant consent and ethical approval will be collected and preserved by the authors.

### COMPETING INTERESTS

Authors have declared that no competing interests exist.

### REFERENCES

1. Haslam DW, James WP. Obesity. *Lancet*. 2005;366(9492):1197–1209.
2. Pi-Sunyer FX. Comorbidities of overweight and obesity: current evidence and research issues. *Med Sci Sports Exerc*. 1999;31(11 suppl):S602–S608.
3. Lord MJ, Small JM, Dinsay JM, Watkins RG. Lumbar lordosis. Effects of sitting and standing. *Spine*. 1997;22:2571–2574.
4. Fernand R, Fox DE. Evaluation of Lumbar lordosis: A prospective and retrospective study. *Spine*. 1985;10:799–803.
5. Gelb DE, Lanke LG, Bridwell KH, Blanke K, McEneaney KW. An analysis of sagittal spine alignment in 100 asymptomatic middle aged and older volunteers. *Spine*. 1995;20:1351–1358.
6. Amonoo-Kuofi HS. Changes in the lumbo-sacral angle, sacral inclination and lumbar spine during ageing. *Acta Anatomica*. 1992;145(4):373–377.
7. Sarikaya S, Ozdolap S, Gumustas S, Koc U. Low back pain and lumbar angles in Turkish coal miners. *Am J Ind Med*. 2007;50:92–96.
8. Caglayan M, Tacar O, Demirant A, et al. Effects of lumbo-sacral angles on development of low back pain. *J Musculoskelet Pain*. 2014;12(3):251–255.
9. Leboeuf-Yde C. Body weight and low back pain: A systematic literature review of 56 journal articles reporting on 65 epidemiologic studies. *Spine*. 2000;25:226–237.
10. Akpinar E, Bashan I, Bozdemir N, Saatci E. Which is the best anthropometric technique to identify obesity: Body mass index, waist circumference or waist-hip ratio. *Coll Antropol*. 2007;31(2):387–393.

© 2022 Agola and Dhaniwala; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/82504>