



Hinge Total Domination on Some Graph Families

Leocint F. Consistente ^{a*} and Isagani S. Cabahug, Jr. ^a

^aDepartment of Mathematics, College of Arts and Sciences, Central Mindanao University, Philippines.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: 10.9734/ARJOM/2022/v18i930404

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/88580>

Received: 20 April 2022

Accepted: 27 June 2022

Published: 28 June 2022

Original Research Article

Abstract

set S of vertices in a graph $G = (V(G), E(G))$ is a hinge dominating set if every vertex $u \in V \setminus S$ is adjacent to some vertex $v \in S$ and a vertex $w \in V \setminus S$ such that (v, w) is not an edge in $E(G)$. The hinge domination number $\gamma_h(G)$ is the minimum size of a hinged dominating set. A set S is called a total dominating set of G if for every vertex in V , including those in S is adjacent to at least one vertex in S . The cardinality of a minimum total dominating set in G is called the total domination number of G and denoted as $\gamma_t(G)$. In this study, a new parameter called hinged total dominating set was introduced and defined as, a hinge total dominating set of a graph G is a set S of vertices of G such that S is both a hinge dominating set and total dominating set. The hinge total domination number, $\gamma_{ht}(G)$, is the minimum cardinality of a hinge total dominating set of G . We initiate a study of hinge total dominating set and present its characterization. In addition, we also determine the exact values of hinge total domination number on some graph families.

Keywords: Hinge domination number; total domination number; hinge total domination number.

2010 Mathematics Subject Classification: 05C35.

*Corresponding author: E-mail: s.consistente.leocint@cmu.edu.ph;

1 Introduction

Total domination in graphs was introduced by Cockayne, Dawes and Hedetniemi [1]. The literature on this subject has been surveyed and detailed in the two excellent domination books by Haynes, Hedetniemi, and Slater who did an outstanding job of unifying results scattered through some 1200 domination papers. Much interest in total domination in graphs has recently arisen from a computer program Graffiti.pc that has generated several hundred conjectures on total domination [2].

On one hand, hinge domination in graphs is a new parameter for domination that was recently accepted and published last 2018. It was introduced by Kavitha B.N. and Indrani Kelkar and is still open for many possible studies [3].

In this paper, we combine the hinge domination and total domination of graphs to form a new variation called hinge total domination of graphs [6, 7, 8]. Moreover, this paper investigates how the hinge total dominating set behaves in different families of graphs [9, 10, 11, 12, 13, 14]. Specifically, we intend to accomplish the following:

- (i) Characterize the hinge total dominating set of some families of graphs namely; path, cycle, star graph, complete bipartite graphs, and complete graphs; and
- (ii) Find the exact values of the hinge total domination number of some families of graphs namely; path, cycle, star graph, complete bipartite graphs, complete graphs, and wheel graphs.

All graphs under considered here are nontrivial, simple, undirected, and finite. For graph theoretic terminologies not specifically defined nor described in this study, please refer to [4].

2 Preliminary Notes

Definition 2.1. [4] A vertex of degree 0 is referred to as an isolated vertex and a vertex of degree 1 is an end-vertex or a leaf.

Definition 2.2. [5] A support vertex is the neighbor of a leaf. If a vertex v is adjacent to two or more leaves, v is said to be a strong support vertex

Definition 2.3. [15] A graph H is a subgraph of a graph G , denoted by $H \subseteq G$, if the vertex set $V(H)$ of H is contained in the vertex set $V(G)$ of G and all edges of H are edges in G , i.e, $V(H) \subseteq V(G)$ and $E(H) \subseteq E(G)$. For any vertex subset $S \subseteq V(G)$, the induced subgraph by S denoted by $\langle S \rangle_G$ contains all the edges of $E(G)$ whose extremities belong to S .

Definition 2.4. [4] A set S of vertices of G is a dominating set if every vertex in $V(G) \setminus S$ is adjacent to at least one vertex in S . The minimum cardinality among the dominating sets of G is called the domination number of G and is denoted by $\gamma(G)$. A dominating set of cardinality $\gamma(G)$ is then referred to as a minimum dominating set.

Definition 2.5. [16] A set S of vertices in a graph $G = (V(G), E(G))$ is a hinge dominating set if every vertex $u \in V \setminus S$ is adjacent to some vertex $v \in S$ and a vertex $w \in V \setminus S$ such that (v, w) is not an edge in $E(G)$. The hinge domination number $\gamma_h(G)$ is the minimum size of a hinged dominating set.

Definition 2.6. [17] A set S is called a total dominating set of G if for every vertex in V , including those in S is adjacent to at least one vertex in S . The cardinality of a minimum total dominating set in G is called the total domination number of G and denoted $\gamma_t(G)$. [18] In addition, a dominating set S of a graph G is a total dominating set if the induced subgraph $\langle S \rangle$ has no isolated vertices.

Example 2.1. In Figure 1, notice that the $\gamma_t(G) = 4$ and $\gamma_h(G) = 2$. Hence, in C_6 , $\gamma_t(G) > \gamma_h(G)$.




Fig. 1. Total domination and hinge domination on C_6

Example 2.2. In Figure 2, notice that the $\gamma_h(G) = 3$ and $\gamma_t(G) = 2$. Hence, in C_3 , $\gamma_h(G) > \gamma_t(G)$.




Fig. 2. Hinge domination and total domination on C_3

Example 2.3. Consider the path P_6 below. By the definition of a hinge dominating set and total dominating set, it is visible that the set $S = \{v_0, v_1, v_4, v_5\}$ is a hinge dominating set and at the same time a total dominating set. Hence, S is a hinge total dominating set of P_6 . This implies that $\gamma_{ht}(P_6) \leq 4$. But no set with three or lesser number of vertices can hinge totally dominate P_6 , meaning $\gamma_{ht}(P_6) > 3$ or $\gamma_{ht}(P_6) \geq 4$. Therefore, we have $\gamma_{ht}(P_6) = 4$.




Fig. 3. Hinge total domination on P_6

3 Main Results

Definition 3.1. A hinge total dominating set of a graph G is a set S of vertices of G such that S is both a hinge dominating set and total dominating set. The hinge total domination number, $\gamma_{ht}(G)$, is the minimum cardinality of a hinge total dominating set of G .

3.1 Some realization results

Theorem 3.1. Let $G = (V(G), E(G))$ be a nontrivial connected graph with leaf vertices v . If $S \subseteq V(G)$ is a nonempty hinge total dominating set of G , then S contains all the leaf vertices of G .

Proof. Let $v \in V(G)$ be a leaf vertex of G . On a contrary, assume that $v \notin S$, then $v \in V(G) \setminus S$. Since a leaf vertex is only adjacent to one support vertex, then vertex v cannot be adjacent to a vertex in S and a vertex in $V(G) \setminus S$. This means that S is not a hinge dominating set, a contradiction to the assumption that S is a hinge dominating set of G . Thus, $v \in S$. \square

Corollary 3.2. *If S is a nonempty hinge total dominating set of G with leaf vertices, then S contains all support vertices of G .*

Proof. Let $y \in V(G)$ be the support vertices of G . By Theorem 3.1, all leaf vertices are in S . Suppose $y \notin S$, then all leaf vertices are not adjacent to any vertex that is in S . This means that S is not a total dominating set, a contradiction to the assumption that S is a hinge total dominating set of G . Thus, $y \in S$. \square

Remark 3.1. For any nontrivial, connected graph $G = (V(G), E(G))$, the set $V(G)$ is a hinge total dominating set.

3.2 Path graphs

Theorem 3.3. *Let P_n be a path graph, $n \geq 6$, with vertex set $V(P_n)$. Then $S \subset V(P_n)$ is a nonempty hinge total dominating set if and only if the following holds:*

- (i) *The leaf and support vertices of P_n are in S ;*
- (ii) *The induced subgraph, $\langle V(P_n) \setminus S \rangle$, form a class of P_2 ;*
- (iii) *The induced subgraph, $\langle S \rangle$, has no trivial graph.*

Proof. Since P_n , $n \geq 6$, is a nontrivial connected graph, by Theorem 3.1 and Corollary 3.2, the leaf and support vertices of P_n are in S . This proves (i).

Let $v \in \langle V(P_n) \setminus S \rangle$ and S be a nonempty hinge total dominating set of P_n . Then, $|N(v) \cap \langle V(P_n) \setminus S \rangle| = 1$ and $|N(v) \cap S| = 1$. This implies $\deg(v) = 1$. Since v is arbitrary, thus for all $v \in \langle V(P_n) \setminus S \rangle$, $\deg(v) = 1$. So, the induced subgraph, $\langle V(P_n) \setminus S \rangle$, form a class of P_2 . This proves (ii).

Suppose the induced subgraph, $\langle S \rangle$ has an isolated vertex. This implies that there exist a vertex a in the vertex set of $\langle S \rangle$ that is not adjacent to some $b \in S$. This means that S is not a total dominating set, a contradiction since S is a hinge total dominating set. Hence, the $\langle S \rangle$ has no trivial graph. This proves (iii).

The converse immediately follows. \square

Remark 3.2. For a path P_n of order $2 \leq n \leq 5$, $V(P_n)$ is the hinge total dominating set.

Theorem 3.4. *If P_n is a path graph of order $n \geq 6$, then*

$$\gamma_{ht}(P_n) = \begin{cases} \frac{n+4}{2} & \text{if } n \equiv 0 \pmod{4}; \\ \frac{n+5}{2} & \text{if } n \equiv 1 \pmod{4}; \\ \frac{n+2}{2} & \text{if } n \equiv 2 \pmod{4}; \\ \frac{n+3}{2} & \text{if } n \equiv 3 \pmod{4}. \end{cases}$$

Proof. Let P_n be a path graph of order $n \geq 6$ and $V(P_n) = \{v_0, v_1, \dots, v_{n-1}\}$. Assume $S \subset V(P_n)$ is nonempty set hinge total dominating set of P_n . Since P_n is a connected graph, by Theorem 3.1 and Corollary 3.2, $\{v_0, v_1, v_{n-2}, v_{n-1}\}$ is in S .

If $n \equiv 0 \pmod{4}$, then S contains vertex sets V_{4i} and V_{4i+1} where $i = \{0, 1, 2, \dots, \frac{n-4}{4}\}$. Hence, $V_{4i} = \{v_0, v_4, v_8, \dots, v_{n-4}\}$ and $V_{4i+1} = \{v_1, v_5, v_9, \dots, v_{n-3}\}$. Notice that $|V_{4i}| = \frac{n-4}{4} + 1 = |V_{4i+1}|$. The remaining vertices that is not covered by V_{4i} and V_{4i+1} are v_{n-2} and v_{n-1} . By Theorem 3.1 and Corollary 3.2, $\{v_{n-2}, v_{n-1}\}$ are in S . Thus,

$$\begin{aligned} |S| &= |V_{4i}| + |V_{4i+1}| + |\{v_{n-2}, v_{n-1}\}| = \frac{n-4}{4} + 1 + \frac{n-4}{4} + 1 + 2 \\ &= \frac{n-4}{4} + 1 + \frac{n-4}{4} + 1 + 2 = \frac{n-4+4+n-4+4+8}{4} \\ &= \frac{2n+8}{4} = \frac{n+4}{2}. \end{aligned}$$

If $n \equiv 1 \pmod{4}$, then S contains vertex sets V_{4i} and V_{4i+1} where $i = \{0, 1, 2, \dots, \frac{n-5}{4}\}$. Hence, $V_{4i} = \{v_0, v_4, v_8, \dots, v_{n-5}\}$ and $V_{4i+1} = \{v_1, v_5, v_9, \dots, v_{n-4}\}$. Notice that $|V_{4i}| = \frac{n-5}{4} + 1 = |V_{4i+1}|$. The remaining vertices that is not covered by vertex sets V_{4i} and V_{4i+1} are v_{n-3} , v_{n-2} and v_{n-1} . By Theorem 3.1 and Corollary 3.2, $\{v_{n-2}, v_{n-1}\}$ are in S . Now, suppose $v_{n-3} \notin S$. Then, $v_{n-3} \in V(P_n) \setminus S$ is not adjacent to all vertices of $V(P_n) \setminus S$, which is a contradiction since S is a hinge total dominating set. Hence, $v_{n-3} \in S$. Therefore,

$$\begin{aligned} |S| &= |V_{4i}| + |V_{4i+1}| + |\{v_{n-3}, v_{n-2}, v_{n-1}\}| = \frac{n-5}{4} + 1 + \frac{n-5}{4} + 1 + 3 \\ &= \frac{n-5+4+n-5+4+12}{4} = \frac{2n+10}{4} \\ &= \frac{n+5}{2}. \end{aligned}$$

If $n \equiv 2 \pmod{4}$, then S contains vertex sets V_{4i} and V_{4i+1} where $i = \{0, 1, 2, \dots, \frac{n-2}{4}\}$. Hence, $V_{4i} = \{v_0, v_4, v_8, \dots, v_{n-2}\}$ and $V_{4i+1} = \{v_1, v_5, v_9, \dots, v_{n-1}\}$. Notice that $|V_{4i}| = \frac{n-2}{4} + 1 = |V_{4i+1}|$. All vertices are covered by vertex sets V_{4i} and V_{4i+1} . Therefore,

$$\begin{aligned} |S| &= |V_{4i}| + |V_{4i+1}| = \frac{n-2}{4} + 1 + \frac{n-2}{4} + 1 \\ &= \frac{n-2+4+n-2+4}{4} = \frac{2n+4}{4} \\ &= \frac{n+2}{2}. \end{aligned}$$

If $n \equiv 3 \pmod{4}$, then S contains vertex sets V_{4i} and V_{4i+1} where $i = \{0, 1, 2, \dots, \frac{n-3}{4}\}$. Hence, $V_{4i} = \{v_0, v_4, v_8, \dots, v_{n-3}\}$ and $V_{4i+1} = \{v_1, v_5, v_9, \dots, v_{n-2}\}$. Notice that $|V_{4i}| = \frac{n-3}{4} + 1 = |V_{4i+1}|$. The remaining vertex that is not covered by remark 3 is v_{n-1} . By Theorem 3.1, v_{n-1} is in S . Therefore,

$$\begin{aligned} |S| &= |V_{4i}| + |V_{4i+1}| + |\{v_{n-1}\}| = \frac{n-3}{4} + 1 + \frac{n-3}{4} + 1 + 1 \\ &= \frac{n-3+4+n-3+4+4}{4} = \frac{2n+6}{4} \\ &= \frac{n+3}{2}. \end{aligned}$$

Furthermore, let x be an arbitrary element in S . Suppose we have $S \setminus \{x\}$. Then the induced subgraph $\langle V(P_n) \setminus S \rangle$ now contains either P_1 or P_3 , a contradiction to the second condition of Theorem 3.3. This means that S is not a hinge total dominating set. Hence, S is a hinge total dominating set with minimum cardinality, $|S| = \gamma_{ht}(P_n)$. Therefore, the results follow. \square

3.3 Cycle graph

Theorem 3.5. Let C_n be a cycle graph, $n \geq 4$, with vertex set $V(C_n)$. Then $S \subset V(C_n)$ is nonempty set hinge total dominating set if and only if the following holds:

- (i) The induced subgraph, $\langle V(C_n) \setminus S \rangle$, form a class of P_2 ;
- (ii) The induced subgraph, $\langle S \rangle$, has no trivial graph.

Proof. Let $v \in \langle V(C_n) \setminus S \rangle$ and S be a nonempty hinge total dominating set of C_n . Then, $|N(v) \cap \langle V(C_n) \setminus S \rangle| = 1$ and $N(v) \cap S = 1$. This implies $\deg(v) = 1$. Since v is arbitrary, thus for all $v \in \langle V(C_n) \setminus S \rangle$, $\deg(v) = 1$. So, the induced subgraph, $\langle V(C_n) \setminus S \rangle$, form a class of P_2 . This proves (i).

Suppose the induced subgraph, $\langle S \rangle$ has a trivial graph. This implies that there exist a vertex a in the vertex set of $\langle S \rangle$ that is not adjacent to some $b \in S$. This means that S is not a total dominating set, a contraction since S is a hinge total dominating set. Hence, the $\langle S \rangle$ has no trivial graph. This proves (ii).

The converse immediately follows. □

Remark 3.3. For a cycle graph C_3 , $\gamma_{ht}(C_3) = 3$.

Theorem 3.6. If C_n is a cycle graph of order $n \geq 4$, then

$$\gamma_{ht}(C_n) = \begin{cases} \frac{n}{2} & \text{if } n \equiv 0 \pmod{4}; \\ \frac{n+1}{2} & \text{if } n \equiv 1 \pmod{4}; \\ \frac{n+2}{2} & \text{if } n \equiv 2 \pmod{4}; \\ \frac{n+3}{2} & \text{if } n \equiv 3 \pmod{4}. \end{cases}$$

Proof. Let C_n be a path graph of order $n \geq 4$ and $V(C_n) = \{v_0, v_1, \dots, v_{n-1}\}$. Assume $S \subset V(C_n)$ to be nonempty hinge total dominating set of C_n . If $n \equiv 0 \pmod{4}$, then S contains vertex sets V_{4i} and V_{4i+1} where $i = \{0, 1, 2, \dots, \frac{n}{4}\}$. Hence, $V_{4i} = \{v_0, v_4, v_8, \dots, v_n\}$ and $V_{4i+1} = \{v_1, v_5, v_9, \dots, v_{n+1}\}$. Notice that $|V_{4i}| = \frac{n}{4} + 1 = |V_{4i+1}|$. Since $v_0 = v_n$ and $v_1 = v_{n+1}$, it necessary to subtract 2 in the cardinality of S to avoid counting the same vertex twice. Therefore,

$$\begin{aligned} |S| &= |V_{4i}| + |V_{4i+1}| - 2 = \frac{n}{4} + 1 + \frac{n}{4} + 1 - 2 \\ &= \frac{n + 4 + n + 4 - 8}{4} = \frac{2n}{4} \\ &= \frac{n}{2}. \end{aligned}$$

If $n \equiv 1 \pmod{4}$, then S contains vertex sets V_{4i} and V_{4i+1} where $i = \{0, 1, 2, \dots, \frac{n-1}{4}\}$. Hence, $V_{4i} = \{v_0, v_4, v_8, \dots, v_{n-1}\}$ and $V_{4i+1} = \{v_1, v_5, v_9, \dots, v_n\}$. Notice that $|V_{4i}| = \frac{n-1}{4} + 1 = |V_{4i+1}|$. Since $v_0 = v_n$, it necessary to subtract 1 in the cardinality of S to avoid counting the same vertex twice. Therefore,

$$\begin{aligned} |S| &= |V_{4i}| + |V_{4i+1}| - 1 = \frac{n-1}{4} + 1 + \frac{n-1}{4} + 1 - 1 \\ &= \frac{n-1 + 4 + n-1 + 4 - 4}{4} = \frac{2n+2}{4} \\ &= \frac{n+1}{2}. \end{aligned}$$

If $n \equiv 2 \pmod{4}$, then S contains vertex sets V_{4i} and V_{4i+1} where $i = \{0, 1, 2, \dots, \frac{n-2}{4}\}$. Hence, $V_{4i} = \{v_0, v_4, v_8, \dots, v_{n-2}\}$ and $V_{4i+1} = \{v_1, v_5, v_9, \dots, v_{n-1}\}$. Notice that $|V_{4i}| = \frac{n-2}{4} + 1 = |V_{4i+1}|$. Since no vertex is repeated, therefore,

$$\begin{aligned} |S| &= |V_{4i}| + |V_{4i+1}| = \frac{n-2}{4} + 1 + \frac{n-2}{4} + 1 \\ &= \frac{n-2+4+n-2+4}{4} = \frac{2n+4}{4} \\ &= \frac{n+2}{2}. \end{aligned}$$

If $n \equiv 3 \pmod{4}$, then S contains vertex sets V_{4i} and V_{4i+1} where $i = \{0, 1, 2, \dots, \frac{n-3}{4}\}$. Hence, $V_{4i} = \{v_0, v_4, v_8, \dots, v_{n-3}\}$ and $V_{4i+1} = \{v_1, v_5, v_9, \dots, v_{n-2}\}$. Notice that $|V_{4i}| = \frac{n-3}{4} + 2$ and $|V_{4i+1}| = \frac{n-3}{4} + 1$. In addition, the vertex v_{n-1} is also in S since $S \setminus \{v\}$ would make S not a hinge dominating set, a contradiction. Now, notice that $v_1 = v_{n+1}$, so, it necessary to subtract 1 in the cardinality of S to avoid counting the same vertex twice. Therefore,

$$\begin{aligned} |S| &= |V_{4i}| + |V_{4i+1}| + |\{v_{n-1}\}| - 1 = \frac{n-3}{4} + 2 + \frac{n-3}{4} + 1 + 1 - 1 \\ &= \frac{n-3+8+n-3+4+4-4}{4} = \frac{2n+6}{4} \\ &= \frac{n+3}{2}. \end{aligned}$$

Furthermore, let x be an arbitrary element in S . Suppose we have $S \setminus \{x\}$. Then the induced subgraph $\langle V(C_n) \setminus S \rangle$ now contains either P_1 or P_3 , a contradiction to the second condition of Theorem 3.3. This means that S is not a hinge total dominating set. Hence, S is a hinge total dominating set with minimum cardinality, $|S| = \gamma_{ht}(C_n)$. Therefore, the results follow. \square

3.4 Star graph

Theorem 3.7. For any star graph $K_{1,n}$, $n \geq 1$, $\gamma_{ht}(K_{1,n}) = n + 1$.

Proof. Let $K_{1,n}$ be a star graph with $V(K_{1,n}) = \{x, v_0, v_1, \dots, v_{n-1}\}$. Since $K_{1,n}$ are composed of leaf vertices $\{v_0, v_1, \dots, v_{n-1}\}$ and support vertex $\{x\}$, by Theorem 3.1 and Corollary 3.2, $V(K_{1,n})$ must be in the hinge total dominating set S . Therefore, $\gamma_{ht}(K_{1,n}) = |S| = n + 1$. \square

3.5 Complete bipartite graph

Theorem 3.8. Let $K_{m,n}$ be a complete bipartite graph, $m, n \geq 2$ with partite sets A and B such that $|A| = m$ and $|B| = n$. Then, $S \subset V(K_{m,n})$ is nonempty hinge total dominating set if and only if the following holds.

- (i) $1 \leq |S \cap A|$ and $1 \leq |S \cap B|$;
- (ii) $|S \cap A| \leq m - 1$ and $|S \cap B| \leq n - 1$.

Proof. Without loss of generality, suppose all vertices from partite set A are not in S . Then, all vertices in B that is in S are not adjacent to vertices in S . This means that S is not a total dominating set of $K_{m,n}$, a contradiction to the assumption that S is a hinge total dominating set. Hence, atleast one vertex from A must be in S . Therefore, $1 \leq |S \cap A|$. The same is true for partite set B , so, $1 \leq |S \cap B|$. This proves (i). Now, suppose all vertices from partite set A are in S . Then, all vertices in B that is not in S are not adjacent to vertices in $K_{m,n} \setminus S$. This means that S is not a hinge dominating set of $K_{m,n}$, a contradiction to the assumption that S is a hinge total dominating set. Hence, at least one vertex from A must not be in S . Therefore, $|S \cap A| \leq m - 1$.

The same is true for partite set B , so, $|S \cap B| \leq n - 1$. This proves (ii).

The converse immediately follows. □

Theorem 3.9. *If $K_{m,n}$ is a complete bipartite graph with $m, n \geq 2$, then $\gamma_{ht}(K_{m,n}) = 2$.*

Proof. Let $K_{m,n}$ be a graph with partite sets A and B such that $m, n \geq 2$. Assume S to be a hinge total dominating set of $K_{m,n}$.

Suppose $\gamma_{ht}(K_{m,n}) = 1$ where $a \in A$ is the only element of S . Then, vertices $A \setminus \{a\}$ is not adjacent to any vertex in S . This implies that S is not a hinge dominating set. In addition, vertex $a \in S$ is not adjacent to any vertices in S which implies that S is not a total dominating set. This is a contradiction since S is a hinge total dominating set. Hence, $\gamma_{ht}(K_{m,n}) \neq 1$ or $\gamma_{ht}(K_{m,n}) \geq 2$. Since we can find vertices $a \in A$ and $b \in B$ that makes the set $S = \{a, b\}$ a hinge total dominating set, hence, $\gamma_{ht}(K_{m,n}) \leq 2$. Therefore, $\gamma_{ht}(K_{m,n}) = 2$. □

3.6 Complete graph

Theorem 3.10. *For any complete graph K_n , $n \geq 2$, $\gamma_{ht}(K_n) = n$.*

Proof. Assume S to be a hinge total dominating set of K_n . Suppose $\gamma_{ht}(K_n) < n$. Then there exists a vertex $x \in V(K_n) \setminus S$. Since all the vertices in K_n are adjacent, then no vertex x can make S a hinge dominating set. This is a contradiction since S is a hinge dominating set. Hence, $\gamma_{ht}(K_n) \geq n$. By Remark 3.1, all vertices of K_n can make S a hinge total dominating set. Hence, $\gamma_{ht}(K_n) \leq n$. Therefore, $\gamma_{ht}(K_n) = n$. □

3.7 Wheel graph

A wheel graph W_3 is the same as a complete graph K_4 . By Theorem 3.10, $\gamma_{ht}(K_4) = 4$. Hence, by transitive property of equality, $\gamma_{ht}(W_3) = 4$.

Remark 3.4. For a wheel graph W_3 , $\gamma_{ht}(W_3) = 4$.

The next result presents the parameter of the wheel graph which is equivalent to the hinge total domination of cycle graphs.

Theorem 3.11. *For any wheel graph W_n , $n \geq 4$, $\gamma_{ht}(W_n) = \gamma_{ht}(C_n)$.*

Proof. By definition, wheel graph is a join of a cycle graph C_n and a trivial graph K_1 . Assume $S \subseteq V(W_n)$ is a nonempty hinge total dominating set of W_n . Now, by Theorem 3.6, the $\gamma_{ht}(C_n)$ was determined. Suppose $x \in K_1$ is not in S , then no condition was defined in the assumption of S . The same result holds if $x \in K_1$ is in S . With this, the least number of vertices in W_n is the same as the least number of vertices in C_n . Therefore, $\gamma_{ht}(W_n) = \gamma_{ht}(C_n)$. □

4 Conclusion and Recommendation

Hinge domination has been used to create better network communication [16] while Total dominations has been used to represent network design in minimizing trade-off between resource allocation and redundancy [17]. In this study, we combined these types of domination and introduced Hinge Total domination. We present some characterizations and exact values for a hinge total domination number on some graph families namely; path graphs, cycle graphs, star graphs, complete bipartite graphs, complete graphs, and wheel graphs. Moreover, some vertices that is always present in any hinge total dominating set was also identified.

It is recommended to investigate further the hinge total domination in the binary operations of graphs.

Competing Interests

Authors have declared that no competing interests exist.

References

- [1] Cockayne EJ, Dawes RM, Hedetniemi ST. Total domination in graphs. *Networks*. 1980;10:211–219.
- [2] Henning MA. A survey of selected recent results on total domination in graphs. *Discrete Mathematics*. 2009;309(1):32–63. Available: <https://doi.org/10.1016/j.disc.2007.12.044>.
- [3] Kavitha BN, Indrani Pramod Kelkar "Hinge Domination Number of a Graph " *International Journal of Engineering Research and Applications (IJERA)*. 2018;8(7):70-7.
- [4] Chartrand G, Lesniak L, Zhang P. *Graphs digraphs*. London: Chapman Hall. 1996;22.
- [5] Desormeaux WJ, Haynes TW, Henning MA. Total domination changing and stable graphs upon vertex removal. *Discrete applied mathematics*. 2011;159(15):1548-1554.
- [6] Acosta HR, Eballe RG, Downhill domination in the tensor product of graphs. *International Journal of Mathematics Analysis*. 2019;13(12):555-569.
- [7] Bresar et al. On the 2-rainbow domination in graphs. *Discrete Applied Mathematics*. 2010;155:2394-2400.
- [8] Cabahug IS, Canoy SR. Connected dr. power dominating sets in graphs. *Advances and Applications in Discrete Mathematics*. 2018;19(3):171-182.
- [9] Cabahug IS, Canoy SR. Dr. power dominating sets in graphs. *International Journal of Mathematics Analysis*. 2016;10(3):139-149. 14
- [10] Cabahug IS, Canoy SR. Independent Dr. power dominating sets in graphs. *Applied Mathematical Sciences*. 2016;10(8):377-387.
- [11] Christopher G. (2011-2020). *Graph Theory*. Penn State Math 485 Lecture Notes, Version 1.5.
- [12] Cohen N, Agneda R, Fujita S, Legay S, Manoussakis Y, et al. Safe Sets in Graphs: Graph Classes and Structural Parameters. *COCOA 2016 - 10th International Conference Combinatorial Optimization and Applications*. 2016, Hong Kong, China;241-253. hal-0162455.
- [13] Harary F. *Graph Theory*. United States of America: Addison-Wesley Publishing Company, Inc; 1969.
- [14] White AT. Chapter 2- A brief introduction to graph theory *North Holland Mathematics Studies*. North-Holland. 2001;188:5-12.
- [15] Beggas F. *Decomposition and Domination of Some Graphs*. *Data Structures and Algorithms [cs.DS]*. University Claude Bernard Lyon 1; 2017. English. fftel-02168197f.
- [16] Kumar VC, Mekala A, Murali R. On hinge domination in graphs. *Journal of Mathematical and Computational Science*. Published; 2021. Available: <https://doi.org/10.28919/jmcs/6089>.
- [17] Henning MA, Naicker V. Graphs with large disjunctive total domination number. *Discrete Mathematics & Theoretical Computer Science*. 2015;17 no. 1(Graph Theory). <https://doi.org/10.46298/dmtcs.2104>.

- [18] Nicholas T, Sheeba Helen T. Global Cototal Domination in Graphs. International Journal of Mathematics and Its Applications. 2018;1051–1055.

© 2022 *Consistente and Cabahug*; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here (Please copy paste the total link in your browser address bar)

<https://www.sdiarticle5.com/review-history/88580>