



Grain Yield and Economic Importance of Bio Fertilizer Rates of Soybean at Begi and Asossa Districts Western Ethiopia

Bakala Anbessa^{a*}, Dessie Almaw^a and Zerihun Getachew^a

^a Assosa Agricultural Research Center, Ethiopian Institute of Agriculture Research, Assosa, Ethiopia, Africa.

Authors' contributions

This work was carried out in collaboration among all authors. Author BA designed the study, performed the statistical analysis, wrote the protocol and wrote the first draft of the manuscript. Authors ZG and DA were implemented the experiment on the ground, collected data and managed the analyses of the study.

Article Information

DOI: 10.9734/IJPSS/2022/v34i2031162

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/63810>

Original Research Article

Received 03 April 2022
Accepted 07 June 2022
Published 21 June 2022

ABSTRACT

A field experiment was conducted on Nitisols of Asossa Agricultural Research Centre during 2017/18 to 2018/2019 cropping season to investigate the response of yield partial budget analysis to biofertilizer rates. The N fertilizer treatments considered in the study consisted of six levels (rates) of biofertilizer, one nationally recommended inorganic N and negative control of N. The treatments consist of: 125 g ha⁻¹ (T1), 250 g ha⁻¹ (T2), 500 g ha⁻¹ (T3), 625 g ha⁻¹ (T4), 750 g ha⁻¹ (T5), 900 g ha⁻¹ (T6), negative control (T7) and 18 kg N ha⁻¹. The treatments were laid out as a Randomized Complete Block Design with three replications. The analysis of variance revealed that bio-fertilizer rates significantly ($P \leq 0.01$) affected grain yield at Begi district, while non-significant ($p < 0.05$) at Asossa district. The maximum (3483.7 kg ha⁻¹ and 1750.9 kg ha⁻¹) grain yield was recorded from 500 g ha⁻¹ bio-fertilizer at Begi and Asossa district respectively. The application of 500 g ha⁻¹ bio-fertilizer had the highest net-benefit of 20850.8 ETB, followed by 125 g ha⁻¹ bio-fertilizer which also had a total of 20196.8 ETB net benefit at Asossa district, while the application of 500 g ha⁻¹ bio-fertilizer had the highest net-benefit of 41644.4 ETB, followed by 125 g ha⁻¹ bio-fertilizer which also had a total of 38315.6 ETB net benefit at Begi district. The application of 500 g ha⁻¹ bio-fertilizer

*Corresponding author: E-mail: anbessa2004@gmail.com;

ha⁻¹ had highest net benefit. Therefore, we recommended the treatment 500 g ha⁻¹ since it produced high marginal rate of return, high net benefit and relatively small total cost of production, for soybean production in Begi and Asossa area.

Keywords: Soybean; bio-fertilizer; net benefit; partial budget analysis.

1. INTRODUCTION

Soybean (*Glycine max* L.) is one of the grain legumes native to East Asia, perhaps domesticated in North and Central China [1,2]. Soybean is grown for its edible bean around the world, which is an important source of inexpensive and high-quality protein (40%) and important source of oil (20%). Soybean is grown on 10⁴ million hectares of land on five continents with annual total production of 241 million tons and productivity of 2.0 ton ha⁻¹ [3].

Soybean is widely produced in the west and southwestern parts of Ethiopia as food crop. Its production is highly concentrated in high rainfall areas, like Asossa, Wollega and Jimma areas. It was recently integrated into the cropping systems of smallholder farmers in Asossa areas and serves as a cash crop for farmers [3]. In Ethiopia, the area allocated for soybean and the corresponding total annual production has been 31,876 ha and 63,653 tons, respectively, with the productivity less than 2 ton ha⁻¹ [4], while the potential soybean yield has been estimated to be in the range of 6 – 8 tons ha⁻¹ in USA [5].

About 50 countries worldwide grow soybean [6]. The United States of America (USA) accounted for 40 to 45% of the world's total soybean production in 2003 [6]. Egypt, the largest producer of soybean in Africa, produces about 180,000 tons annually [7]. Soybean improves soil fertility by fixing nitrogen from the atmosphere [8]. Soybeans are sensitive to low pH. In acid soils, liming is essential to raise the pH to 6.0 or 6.5 for optimum yield production. Soils in Africa are typically highly variable in fertility and how they respond to application of inputs [9]. Soil nutrient depletion, nutrient mining and degradation have been considered serious threats to agricultural productivity and have been identified as major causes of decreased crop yields and per capita food production in sub-Saharan Africa [10].

Biological nitrogen fixation (BNF) and nitrogen fertilizers are the main sources for meeting the N requirement of high-yielding soybean varieties. BNF is an effective and efficient source of N

supply to plants under favorable atmospheric and environmental conditions [11]. BNF is very useful for small holder farmers as it is cost effective, environmentally friendly, meets the N requirement of the legumes and reduces the N demand of the succeeding crops. Inoculation with compatible and effective rhizobia may be necessary to optimize the nitrogen fixation and hence legumes grain yields, where a low population of native rhizobial strains predominates [12]. Therefore, evaluation and identification of appropriate and effective rhizobial strains are crucial to enhance nitrogen fixation and yield of soybean [13-16]. In the present investigation, the influences of inoculation rate in soil with low rhizobia population associated with soybean is thoroughly examined under field conditions during the main rain season. Therefore, the specific objective of this piece of research work was to evaluate the rates of inoculant that are used as soybean rhizobial biofertilizer and to determine the optimum rate that effectively improve BNF of soybean at Assossa and Begi Districts.

2. MATERIALS AND METHODS

2.1 Treatments and Experimental Design

A field experiment was conducted under rain fed conditions during the main rainy season (June to October) of 2017 and 2018 to investigate the effects of biofertilizer application rates on grain yield of soybean at Asossa and Begi district. The N fertilizer treatments considered in the study consisted of six levels (rates) of biofertilizer, one nationally recommended inorganic N and negative control of N. All plots uniformly received phosphorous as basal application (20% P from triple super phosphate) during planting. The experiment was then laid in a randomized complete block design with three replications consisting of a total of 8 treatments. A plot size of 3.6*3m (10.8 m²) was used as an experimental unit. The blocks were separated by a 1.5 m wide open space where as the plots within a block were separated by a 0.75 m wide space. Soil bunds were constructed around each plot and around the entire experimental field to minimize nutrient, water movement, and cross

contamination from plot to plot. Weed control was achieved manual by hand picking. Crop growth was then monitored until harvest.

2.2 Statistical Data Analysis

Analyses of variances for the data were conducted using the SAS GLM procedure ([17]. Least significant difference (LSD) test at 5% probability used for mean separation when the analyses of variance indicate the presence of significant differences.

2.3 Economic Analysis

Mean grain yield of the selected treatments were used in partial budget analysis [18]. Economic analysis was performed to investigate the economic feasibility of the treatments (fertilizer rates). A partial budget, dominance and marginal analysis were conducted. The average open market price (Birr kg^{-1}) for maize and the official prices of blended, Urea and TSP fertilizers were used for the analysis. The dominance analysis procedure as detailed in [18] was used to select potentially profitable treatments from the range that were tested. The selected and discarded treatments using this technique are referred to as 'undominated' and 'Dominated' treatments, respectively. The undominated treatments were ranked from the lowest (the farmers' practice) to the highest cost treatment. For each pair of ranked treatments, a % marginal rate of return (MRR) was calculated. The % MRR between any pair of undominated treatments denotes the return per unit of investment in fertiliser expressed as a percentage.

3. RESULT AND DISCUSSION

Analysis of the grain yield data showed non-significant difference among the treatments at Asossa district. Though there was no observable statistical difference among treatment at Asossa area, still T3 increased the grain yield of soybean by 37.4% over the control plot, that can be taken as acceptable yield increment for the small holder farmers. In addition, compared to the recommended N and P treatment, T3 increased the mean value of soybean grain yield by 20.5%. On the other hand, the analysis of variance of soybean grain yield showed significant difference among rates of inoculation and fertilizer application at Begi district (Table 1). Accordingly, the biofertilizer rate, T3, resulted in 60.7% and 56.5% yield increase compared to the negative control N (T7) and the recommended

inorganic (N) application, respectively. Hence, the normal recommendation rate (500 g/ha) of the National Soil Testing Center is optimal rate specifically for Begi and similar areas, while it still remains comparable to Asossa area.

3.1 Partial Budget Analysis at Begi

The application of 500 g/ ha biofertilizer had the highest net-benefit of 41644.4 ETB, followed by 125 g ha^{-1} biofertilizer which also had a total of 38315.6 ETB net benefit. The lowest net benefit was obtained by the application of the negative control with total of 26011.2 ETB followed by 900 g ha^{-1} biofertilizer with net benefit of 24861.2 ETB. The increased production of the crop due to the application of inputs might not be beneficiary to farmers [18]. Therefore, partial budget analysis [18] was employed to estimate the net benefit, dominance analysis and marginal rate of return that could be obtained from various alternative treatments [18]. The profitability of the study showed that application of 500 g ha^{-1} biofertilizer which provided the relatively high net benefit (41644.4 ETB), was the peak to apply fertilizers.

Economic analysis revealed that maximum marginal rate of return was recorded with application of 125 g/ha biofertilizer (307.61%), followed by 500 ha^{-1} biofertilizer (110.2%). The marginal rates of those treatments were well above the 100% minimum [18]. According to [18] experience and empirical evidence, for the majority of situations indicated that the minimum rate of return acceptable to farmers would be between 50 and 100%. In the present study the treatments that had above 100% marginal rate return was recommended for the farmers, with treatments that had small number of variable cost. This treatment was 125 g/ha and 500 biofertilizer g ha^{-1} .

The % MRR between any pair of undominated treatments denotes the return per unit of investment in biofertilizer expressed as a percentage. The results of undominated treatments indicated that for each one birr invested in purchase or production of fertilizers that was possible to recover one birr plus an extra of 3.08 birr ha^{-1} and 1.1 birr ha^{-1} as the fertilizer application changed from unfertilized plot to 125 g ha^{-1} biofertilizer g/ha and 500 g/ha biofertilizer respectively. Passing from the first treatment that had the lowest costs that vary to the end treatment which had the highest cost that vary, the marginal rate of return obtained was

above the minimum acceptable marginal rate of return. Accordingly, the study revealed that application of 500 g ha⁻¹ biofertilizer was considered as the best for recommendation. The best recommendation for treatments subjected to marginal rate of return is not necessarily based

on the highest marginal rate of return, rather based on the minimum acceptable marginal rate of return and the treatment with the highest net benefit, relatively low variable cost together with an acceptable MRR becomes the tentative recommendation [18].

Table 1. Effect of biofertilizer rate on seed yield of soybean

Treatment	Grain yield (kg/ha)	
	Begi	Asossa
125 g ha ⁻¹ (T1)	3196.3ab	1686.4
250 g ha ⁻¹ (T2)	2742.6bc	1430.8
500 g ha ⁻¹ (T3)	3483.7 a	1750.9
625 g ha ⁻¹ (T4)	2328.5dc	1261.2
750 g ha ⁻¹ (T5)	2477.0cd	1454.9
900 g ha ⁻¹ (T6)	2105.1cd	1329.1
No (T7)	2167.6cd	1274.5
18kg Nha ⁻¹ (T8)	2226.2cd	1451.9
LSD	608.97	-
CV %	13.42	29.53

Table 2. Partial Budget analysis of biofertilizer rate at Begi

Treatment	VC (ETB ha ⁻¹)	TGR (ETBha ⁻¹)	NB (ETBha ⁻¹)	MRR%
No biofertilizer	0	26011.2	26011.2	0
125 g ha ⁻¹ biofertilizer	40	38355.6	38315.6	307.61
250 g ha ⁻¹ biofertilizer	80	32911.2	32831.2	D
500 g ha ⁻¹ biofertilizer	160	41804.4	41644.4	110.2
625 g ha ⁻¹ biofertilizer	240	27942	27702	D
750 g ha ⁻¹ biofertilizer	320	29724	29404	D
18kg N ha ⁻¹	322.4	26714.4	26392	D
900 g ha ⁻¹ biofertilizer	400	25261.2	24861.2	D

N. B: Prices - Urea= 8.24 birr kg⁻¹, TSP=12.75 birr kg⁻¹, Price of soybean=12 birr kg⁻¹, Seed=10 birr kg⁻¹ & Labor cost =30 birr/ person/day for 8 hours, D= dominated, TC=Total cost, Gross return (Return from Grain) =price /kg yield in kg and Net return = gross return – Total cost, VC = variable cost, GR= growth return, TGR = total growth return from grain, NB = net benefit*

Table 3. Partial Budget analysis of biofertilizer rate at Asossa

Treatment	VC (ETB ha ⁻¹)	TGR (ETBha ⁻¹)	NB (ETBha ⁻¹)	MRR%
No biofertilizer	0	15294	15294	0
125 g ha ⁻¹ biofertilizer	40	20236.8	20196.8	122.57
250 g ha ⁻¹ biofertilizer	80	17169.6	17089.6	D
500 g ha ⁻¹ biofertilizer	160	21010.8	20850.8	47.1
625 g ha ⁻¹ biofertilizer	240	15134.4	14894.4	D
750 g ha ⁻¹ biofertilizer	320	17458.8	17138.8	D
18kg N ha ⁻¹	322.4	17422.8	17100.4	D
900 g ha ⁻¹ biofertilizer	400	15949.2	15549.2	D

N. B.Prices: - Urea= 8.24 birr kg⁻¹, TSP=12.75 birr kg⁻¹, Price of soybean=12 birr kg⁻¹, Seed=10 birr kg⁻¹ & Labor cost =30 birr/ person/day for 8 hours, TC=Total cost, Gross return (Return from Grain) =price /kg yield in kg and Net return = gross return – Total cost, VC = variable cost, GR= growth return, TGR = total growth return from grain, NB = net benefit*

3.2 Partial Budget Analysis at Asossa

The application of 500 g/ ha biofertilizer had the highest net-benefit of 20850.8 ETB, followed by 125 g/ha biofertilizer which also had a total of 20196.8 ETB net benefit. The lowest net benefit was obtained by the application of the 625 g/ ha biofertilizer with total of 14894.4 ETB followed by negative N control with net benefit of 15294 ETB. The profitability of the study showed that application of 500 g ha⁻¹ biofertilizer which provided the relatively high net benefit (20850.8 ETB), was the peak to apply fertilizers. Economic analysis revealed that maximum marginal rate of return was recorded with application of 125 g/ha biofertilizer (122.57%), followed by 500 g/ha biofertilizer (47.1%). The marginal rate of 125 g/ha biofertilizer treatments were well above the 100% minimum [18]. On the other hand the marginal rate of return of 500 g/ha biofertilizer treatments were below the 50 - 100% minimum [18]. According to [18] experience and empirical evidence, for the majority of situations indicated that the minimum rate of return acceptable to farmers would be between 50 and 100%. In the present study the treatments that had above 50% marginal rate return was recommended for the farmers, with treatments that had small number of variable cost. This treatment was 125 g ha⁻¹ biofertilizer g/ha for Asossa district, while based on net benefit and relatively lower cost production the farmers of Asossa area can use 500 g/ha biofertilizer.

The % MRR between any pair of undominated treatments denotes the return per unit of investment in biofertilizer expressed as a percentage. The results of undominated treatments indicated that for each one birr invested in purchase or production of fertilizers that was possible to recover one birr plus an extra of 1.23 birr ha⁻¹ and 0.47 ha⁻¹ as the fertilizer application changed from unfertilized plot to 125 g ha⁻¹ biofertilizer g ha⁻¹ and 500 g/ha biofertilizer respectively. Accordingly, the study revealed that application of 125 g ha⁻¹ biofertilizer was considered as the best for recommendation, while based on net benefit and relatively lower cost production the farmers of Asossa area can use 500 g ha⁻¹ biofertilizer. The best recommendation for treatments subjected to marginal rate of return is not necessarily based on the highest marginal rate of return, rather based on the minimum acceptable marginal rate of return and the treatment with the highest net benefit, relatively low variable cost together with an acceptable MRR becomes the tentative recommendation [18].

4. SUMMARY AND CONCLUSIONS

The present study was conducted in Benishangul Gumuz Regional State, at Asossa Agricultural Research Center station in the 2016/2017 to 2017/2018 main cropping season under rain fed field conditions. The profitability of the study showed that application of 500 g/ha biofertilizer which provided the relatively highest net benefit (41644.4 ETB and 20850.8 ETB), was the optimum rate to apply biofertilizers for Begi and Asossa district. The best recommendation for treatments subjected to marginal rate of return is not necessarily based on the highest marginal rate of return, rather based on the minimum acceptable marginal rate of return and the treatment with the high net benefit, relatively low variable cost together with an acceptable MRR becomes the tentative recommendation. Therefore, we recommend the treatments (500 g/ha biofertilizer) that have acceptable marginal rate of return, relatively high net benefit and relatively small total cost of production for soybean production at Begi and Asossa district respectively.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Laswai HS, Mpagalile JJ, Silayo VCK, Ballegu WR. Use of soybeans in food formulation in Tanzania. In: Myaka FA, Kirenga G, Malema, B (Eds.). Proceedings of the first national soybean stakeholders workshop, 10th-11th November 2005, Morogoro, Tanzania. 2005;52-59.
2. Food and Agriculture Organization (FAO) United Nations Food and Agricultural Organization, FAOSTAT Agricultural Database; 2012.
3. Nigussie M, Girma A, Anchala C, Kirub A. Improved technologies and resource management for Ethiopian Agriculture. A Training Manual. RCBP-MoARD, Addis Ababa, Ethiopia; 2009.
4. Central Statistical Authority (CSA) Agricultural Sample Survey. Ethiopia: Addis Ababa; 2008/9 Report on area and production for major crops (Private Peasant Holdings, Main Season); 2013.
5. Specht JE, Hume DJ, Kumudini SV. Soybean yield potential— A genetic and physiological perspective. *Crop Sci.* 1999; 39:1560-1570.

6. Boerme RH, Specht JE. Soybeans: Improvement, production and uses, Third Edition, No. 16 in the series AGRONOMY, ASA, CSSA, and SSSA, Madison; 2004.
7. USDA-ERS. Foreign agriculture service-commodity intelligence report; 2009. Available:<http://www.pecad.fas.usda.gov/>.
8. Sanginga N, Dashiell K, Diels J, Vanlauwe B, Lyasse O, Carsk RJ et al. Sustainable resource management coupled to resilient germplasm to provide new intensive cereal-grain legume-livestock systems in the dry savanna. *Agriculture, Ecosystems and Environment* 2003;100:305-314.
9. [agra-progress-report-2007-2014.pdf](https://www.rockefellerfoundation.org/wp-content/uploads/agra-progress-report-2007-2014.pdf), <https://www.rockefellerfoundation.org/wp-content/uploads/agra-progress-report-2007-2014.pdf>
10. Henao J, Baanante C. Agricultural production and soil nutrient mining in Africa: implication for resource conservation and policy development. IFDC Technology Bulletin. International Fertilizer Development Center. Muscle Shoals, A. L. USA; 2006.
11. Hungria M, Vargas MAT. Environmental factors affecting N₂ fixation in grain legumes in the tropics, with an emphasis on Brazil. *Field Crops Res.* 2000;65:151–164.
12. Chianu JN, Nkonya EM, Mairura FS, Chianu JN, Akinnifesi FK. Biological nitrogen fixation and socioeconomic factors for legume production in sub-Saharan Africa: A review. *Agronomy for Sustainable Development.* 2011;31(1):139-154.
13. Amare B. Progress and future prospect in soybean research in Ethiopia. In: Proc 19th National crop improvement Conference. Addis Ababa, Ethiopia: Institute of agricultural Research; 1987;252–265.
14. Bhangoo MS, Albritton DJ. Nodulating and non-nodulating Lee soybean isolines response to applied nitrogen. *Agron J.* 1976;68:642–645.
15. Graham PH, Vance CP. Legumes: Importance and Constraints to Greater Use. *Plant Physiol.* 2003;131:872–877.
16. Ogoke IJ, Togun AO, Carsky RJ, Dashiell KE. N₂ Fixation by Soybean in the Nigerian Moist Savanna: Effects of Maturity Class and Phosphorus Fertilizer. *Tropicultura.* 2006;24(4):193–199.
17. SAS (Statistical Analysis system) Institute Inc.,1998, SAS Campus Drive, Cary, NorthCarolina 27513.
18. CIMMYT (International Maize and Wheat Improvement Center), 1988. From Agronomic Data to Farmer Recommendations: An Economics Training Manual.

© 2022 Anbessa et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/63810>