



## ***Xylopi*a *aethi*o*pica* Fruit Increased Prothrombin Time, Activated Partial Thromboplastin Time and Erythrocyte Sedimentation Rate of Wistar Rats**

Emmanuel O. Ogbuagu <sup>a</sup>, Augustine I. Airaodion <sup>b\*</sup>, Chika L. Uche <sup>c</sup>,  
Uloaku Ogbuagu <sup>b</sup>, Edmund O. Ezirim <sup>d</sup>, Prince C. Unekwe <sup>e</sup>  
and Ifeoma N. Nweke <sup>a</sup>

<sup>a</sup> Department of Pharmacology and Therapeutics, Abia State University, Uturu, Nigeria.

<sup>b</sup> Department of Biochemistry, Federal University of Technology, Owerri, Imo State, Nigeria.

<sup>c</sup> Department of Haematology, Abia State University, Uturu, Nigeria.

<sup>d</sup> Department of Obstetrics and Gynecology, Abia State University, Uturu, Nigeria.

<sup>e</sup> Department of Pharmacology and Therapeutics, Nnamdi Azikiwe University, Awka Anambra State, Nigeria.

### **Authors' contributions**

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

### **Article Information**

#### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/86463>

**Original Research Article**

**Received 27 February 2022**

**Accepted 07 May 2022**

**Published 09 May 2022**

### **ABSTRACT**

**Background:** Traditionally, the use of *Xylopi*a *aethi*o*pica* fruit for therapeutic purposes is on the increase without any consideration of its safety and toxicity.

**Aim:** This study was therefore designed to examine its effect on coagulation factors of Wistar rats.

**Methodology:** The fruits of *Xylopi*a *aethi*o*pica* were air-dried and extracted by Soxhlet extractor using ethanol as solvent. The median lethal dose (LD<sub>50</sub>) of the extract was assessed using standard method. Thirty adult female Wistar rats were divided into five groups of six rats each. Animals in groups A, B, C, and D were treated with 130, 259, 389 and 518 mg/kg body weight of *X. aethi*o*pica* fruit extract respectively, while those in group E received normal animal feeds and water only. The oral method was used to provide the medication once a day for 28 days. Standard procedures were used to determine prothrombin time, activated partial thromboplastin time, and erythrocyte sedimentation rate.

\*Corresponding author: E-mail: [augustineairaodion@yahoo.com](mailto:augustineairaodion@yahoo.com);

**Results:** The extract of *Xylopiæ aethiopicæ* fruit was observed to show a significant ( $p < 0.05$ ) increase in prothrombin time, activated partial thromboplastin time and erythrocyte sedimentation rate when compared with those of the control group.

**Conclusion:** *Xylopiæ aethiopicæ* significantly prolongs prothrombin time (PT) and activated partial thromboplastin time (aPTT) and in addition has thrombocytopenic property. These anticlotting properties have clearly elucidated and unraveled mechanisms of action which have laid credence to its use by women in flushing out remnants of product of conception post-delivery. Increase in ESR by *Xylopiæ aethiopicæ* may result from an inherent property in this fruit that is yet to be identified.

**Keywords:** Prothrombin time; activated partial thromboplastin time; erythrocyte sedimentation rate  
*Xylopiæ aethiopicæ* fruit.

## 1. INTRODUCTION

*Xylopiæ aethiopicæ* is a useful remedy used as herbal medication in traditional medicine [1]. "Virtually every aspect of the plant has medicinal value. However, its fruits have more pronounced utilization for therapeutic purposes. It has been reported to be effective against cough, biliousness, bronchitis, rheumatism, dysentery, malaria, uterine fibroid as well as amenorrhæa" [2,3]. It has also been reported to be effective against different pains and aches [4]. The fruit has been demonstrated to have excellent anthelmintic activity against *Nippostrongylus brasiliensis*, thus, its usage by human as an anthelmintic agent should be examined [5].

Different extracts of *Xylopiæ aethiopicæ* have also proved their efficacy in managing sickle cell disease [6]. "Oil extracted from the seeds served as a lotion for boils and eruptions, and as a liniment for lumbago. Herbal medical practitioners and birth attendants use a decoction of the seeds to induce placental discharge postpartum due to its abortifacient effect" [7].

The roots of the plant are used in tinctures, against toothache. Aqueous preparation of the root is given after parturition to prevent infection [8]. The powdered root is very useful in dressing of wounds and in the local treatment of cancer [1]. "The leaves and bark are used in traditional medicine to manage boils, sores, wounds and cuts" [9]. "A preparation of the leaves is useful as an anti-emetic agent. The ground leaves are administered as snuff for managing headaches" [10]. A survey carried out by Kadirî et al. [11] proved that the stem bark of *Xylopiæ aethiopicæ* when combined with other herbal remedy prepared with alcohol is effective in the management of breast infections after parturition.

*Xylopiæ aethiopicæ* has also been applied as carminative, stimulant as well as adjunct to other remedies for the management of skin infections [12]. A study by Van-Hai [13] reported that the fruits of *Xylopiæ aethiopicæ* could boost the immunity of fishes against different infections. This application of *Xylopiæ aethiopicæ* fruit is of great interest to scholars of aquaculture as these natural agents, unlike antibiotics, have little or no side effect to the fishes as well as the environment. The propensity of *Xylopiæ aethiopicæ* to have the antioxidant potential of tomatoes avails the opportunity for the assessment of its candidacy as a stabilizer in the food industry.

*Xylopiæ aethiopicæ* is characterized with numerous chemical components with various medicinal potentials [14]. The chemical components of this plant have been investigated to include saponins, sterols, carbohydrates, glycosides, mucilage, acidic compounds, tannins, balsams, cardiac glycosides, volatile aromatic oils, phenols [15,16], alkaloids, rutin and fixed oils [17,18]. "The plant has also been known to contain vitamins such as vitamin A, vitamin B, vitamin C, vitamin D, and vitamin E, and proteins as well as several minerals such as copper, manganese and zinc" [16,18]. The impact of the fruit on body weight and glucose concentration [19] as well as lipid profile [20] of animals has been reported. The fruit has also been reported to induce hepatotoxicity [21], renal toxicity [22] as well as oxidative stress [23]. Recently, Ogbuagu et al. [24] reported that the fruit extract of *Xylopiæ aethiopicæ* adversely perturbed sperm qualities in male Wistar rats. This study was therefore designed to examine its effect on coagulation factors and erythrocyte sedimentation rate of Wistar rats.

## 2. MATERIALS AND METHODS

### 2.1 Collection and Authentication of Plant Materials

The *Xylopia aethiopica* fruits were purchased from a market in Aba, Abia State. They were authenticated by Prof. (Mrs.) Margaret Bassey of Botany and Ecological Studies Department, University of Uyo. It was assigned a voucher number of UU/PH/4e and deposited in the Herbarium of the Department of Pharmacognosy and Natural Medicine, University of Uyo, Akwa-Ibom State, Nigeria.

### 2.2 Extraction of Plant Materials

Extraction of the plant was carried out based on the outlined method in Ogbuagu et al. [7]. The fruits were rinsed under flowing tap water to eliminate contaminants and air-dried. The plant material was milled by laboratory blender. The pulverized plant material was macerated in 250 mL of 99.8% ethanol (Sigma Aldrich) contained in a flask attached to a Soxhlet extractor coupled with condenser and heating mantle (Isomantle). It was then poured into the sample holder (thimble) and inserted in the apparatus. The side arm is lagged with glass wool. The mixture was heated using the heating mantle (Isomantle) at 60 °C and as the temperature rises it starts to evaporate, going via the extractor to the condenser. The condensate dropped into the reservoir keeping the thimble. As soon as the solvent gets to the siphon it emptied itself into the flask and the process repeats itself. The procedure goes on until it is exhaustively extracted. The process runs for a total of 13 hours. As soon as it was set up, it was allowed to run without interruption as long as water and power supply were not interrupted. The apparatus was switched on and off and overnight running was not allowed, and the time for the complete process split over some days. The extract was poured into 1000 mL beaker and concentrated to dryness in water bath (A3672- Graffin Student Water Bath) at 35 °C. The total weight of the marc (residue) and the concentrated extract were noted. Several days was spent on the entire process. The evaporated extract was kept in the refrigerator until when the need for it arise [25].

### 2.3 Determination of Median Lethal Dose (LD<sub>50</sub>)

The lethal dose (LD<sub>50</sub>) of the fruit extract was determined using albino mice according to the

method described by Airaodion et al. [19]. This method involves two phases:

In Phase one, five groups containing five mice each weighing between 20 g and 27g were fasted for 18 hours. They were respectively treated with 1000 mg/kg, 2000 mg/kg, 3000 mg/kg, 4000 mg/kg and 5000 mg/kg body weight via intraperitoneal (i.p) route and were monitored for visible signs of toxicity and mortality for 24 hours. A dosage of 1000 mg/kg recorded 0% mortality while 2000 mg/kg, 3000 mg/kg 4000 mg/kg and 5000 mg/kg recorded 100% mortality within 24 hours. Based on the value of phase one, phase two was conducted.

In Phase two, twenty-five albino mice weighing between 20 and 27g were grouped into 5 of 5 mice per group and were fasted for 18 hours. Each group was administered 1200 mg/kg, 1400 mg/kg 1600 mg/kg, 1800 mg/kg and 2000 mg/kg b. wt. intraperitoneally (i.p) and was observed for physical signs of toxicity and mortality within 24 hours. 1200 mg/kg recorded 0% mortality while 1400 mg/kg, 1600 mg/kg, 1800 mg/kg and 2000 mg/kg recorded 100% mortality within 24 hours. The LD<sub>50</sub> was computed as according to the formula below:

$$LD_{50} = \sqrt{ab}$$

Where a = highest dose giving 0% mortality  
b = lowest dose giving 100% death

### 2.4 Experimental Design

Thirty Wistar rats used in this study were purchased from the University of Uyo, Nigeria. They were acclimatized for seven days prior to the start of the treatment. The weights were determined and were separated into five groups of six rats each. Groups A, B, C, D served as the experimental groups, while group E served as the control. Animals in group A were exposed to 130 mg/kg b. wt. (10% of LD<sub>50</sub>) of *X. aethiopica* fruit extract, those in group B were treated with 259 mg/kg b. wt. (20% of LD<sub>50</sub>) of *X. aethiopica* fruit extract, those in group C were exposed to 389 mg/kg b. wt. (30% of LD<sub>50</sub>) of *X. aethiopica* fruit extract, those in group D were treated with 518 mg/kg b. wt. (40% of LD<sub>50</sub>) of *X. aethiopica* fruit extract, while those in group E (control) received normal animals feeds and water only. The treatment was done once daily for 28 days via oral route. After the treatment period, the animals were sacrificed under ether anaesthesia

in a desiccator after an overnight fast. Blood was taken from the rats through cardiac puncture [26].

## 2.5 Estimation of Prothrombin Time (PT)

Prothrombin time (PT) was determined according to the method described by Onyebuagu [27]. 0.1 mL of platelet poor plasma (PPP) sample from each animal was dropped into test tubes numbered 1-30. Thereafter, 0.1 mL of commercially prepared control sample was dropped into tube no. 31. All the tubes were warmed at 37 °C for 3 minutes using water bath. After that, each test tube received 0.2 mL of prothrombin-thromboplastin reagent, and the clotting time for each tube was measured.

## 2.6 Estimation of Activated Partial Thromboplastin Time (aPTT)

Activated partial thromboplastin time (aPPT) was determined according to the method described by Onyebuagu [27]. 0.1 ml of aPTT reagent (Darkez Ltd) was added to a pair of test tubes containing 0.1 ml of the test samples. The tubes were incubated at 37 °C for 5 minutes, before adding 0.1 mL of freshly prepared CaCl<sub>2</sub> solution. The clotting time was recorded using stop watch.

## 2.7 Estimation of Erythrocyte Sedimentation Rate (ESR)

Erythrocyte sedimentation rate (ESR) was determined using Westergreen method described by Onyebuagu [27]. 0.4ml of 2.8% sodium citrate was drawn into a syringe and the anticoagulated blood sample was added to the 2ml mark. The mixture was then suctioned over the zero mark in the calibrated Westergreen tube. With the index finger, the top of the tube was swiftly closed, and then carefully spun until the upper level of blood was exactly at the zero point. The clock was then started and left to run for one hour while the temperature was taken. After 1 hour, the height of the clear fluid above the upper limit of the red cells column was measured in millimeters per hour.

## 2.8 Statistical Analysis

Results are expressed as mean ± standard deviation. The levels of homogeneity among the groups were assessed using One-way Analysis

of Variance (ANOVA) followed by Tukey's test. All analyses were done using Graph Pad Prism Software Version 8.2 and P values < 0.05 were considered statistically significant.

## 3. RESULTS

### 3.1 Median Lethal Dose (LD<sub>50</sub>) Result

The physical signs of toxicity observed in the animals included excitation, paw licking, increased respiratory rate, decreased motor activity, gasping and coma which was followed by death. In the first phase of the median lethal dose determination, no mortality was recorded in the group treated with 1000 mg/kg body weight of *X. aethiopica* fruit extract. However, 100 % mortality was recorded in the groups treated with 2000, 3000, 4000, and 5000 mg/kg body weight of *X. aethiopica* fruit extract respectively. Similarly, in the second phase of medial lethal dose determination, no mortality was recorded in the group treated with 1200 mg/kg body weight of *X. aethiopica* fruit extract while 100% mortality was recorded in the groups treated with 1400, 1600, and 1800 mg/kg body weight of *X. aethiopica* fruit extract respectively as presented in Table 1.

The median lethal dose (LD<sub>50</sub>) was calculated as geometrical means of the maximum dose producing 0% (a) and the minimum dose producing 100% mortality (b).

$$LD_{50} = \sqrt{ab}$$

Where a = 1200 mg/kg

b = 1400 mg/kg

$$LD_{50} = 1296.15 \text{ mg/kg}$$

### 3.2 Impact of Ethanol Extract of *Xylopi* *aethiopica* fruit on Prothrombin time (PT), Activated Partial Thromboplastin Time (aPTT), Erythrocyte Sedimentation Rate (ESR) after 28 Days of Treatment

The extract of *Xylopi aethiopica* was observed to show a significant rise in prothrombin time, activated partial thromboplastin time and erythrocyte sedimentation rate when compared with those of the control group (p< 0.05), as presented in Table 2.

**Table 1. The Median lethal dose (LD<sub>50</sub>) of *Xylopiya aethiopic* fruit extract**

Study Phase/ (Animal)	Dosage of Extract (mg/kg) b.w	No of Mice per Group	No. of Death Recorded	% Mortality
<b>PHASE ONE</b>				
I	1000	5	0	0
II	2000	5	5	100
III	3000	5	5	100
IV	4000	5	5	100
V	5000	5	5	100
<b>PHASE TWO</b>				
I	1200	5	0	0
II	1400	5	5	100
III	1600	5	5	100
IV	1800	5	5	100
V	2000	5	5	100

$LD_{50} = 1296.15 \text{ mg/kg}$

**Table 2. Effect of ethanol extract of *Xylopiya aethiopic* fruit on Prothrombin time (PT), activated partial thromboplastin time (aPTT), Erythrocyte Sedimentation Rate (ESR) after 28 days of treatment**

Group	A	B	C	D	E	P Value
<b>Dose of extract (mg/kg)</b>	<b>130</b>	<b>259</b>	<b>389</b>	<b>518</b>	<b>Control</b>	
PT (Sec)	15.34±2.92*	17.15±3.47*	20.01±3.27*	22.56±2.35*	12.99±2.66	0.03
aPTT (Sec)	36.83±3.37*	38.56±3.93*	39.63±4.27*	42.87±3.24*	32.67±2.78	0.05
ESR (mm/hr)	5.04±0.94	6.97±0.23*	7.56±0.45*	9.67±1.04*	4.67±0.34	0.02

Values are presented as Mean±S.D, where n = 6. Values with \* are statistically significant at p value ≤ 0.05 when compared with the control group

Legend: aPTT = Activated Partial Thromboplastin Time, PT = Prothrombin Time, ESR = Erythrocyte Sedimentation Rate

#### 4. DISCUSSION

The acute toxicity study of the plant extracts recorded 100% mortality at a dose of 1400 mg/kg bodyweight and above (table 1). This shows that the fruit of *Xylopiya aethiopic* might be highly toxic. Excitation, paw licking, increased respiration rate, decreased motor activity, gasping, and coma were among the physical symptoms of toxicity detected in the animals, which were followed by death.

In this study, administration of ethanol extract of *Xylopiya aethiopic* fruit produced a significant increase in prothrombin time (PT) and activated partial thromboplastin time (aPTT) when compared with those in the control group. This increase was most significant at the highest dose. This significant increase appears to be

related to the decrease in the platelet count reported by Ogbuagu et al. [28] when they exposed animals to *Xylopiya aethiopic* fruit for 28 days. This claim is based on the fact that a decrease in platelet number is linked to hemostatic mechanisms problems. "The findings on the effect of dietary *Xylopiya aethiopic* on coagulation factors observed" in this study is similar to the report by Obembe et al. [29] who reported a significant increase in clotting time when animals were treated with 200 mg/kg body weight of *Xylopiya aethiopic*, but contradicted the report by Nwafor [14], who observed "a slight decrease in PT and aPTT in albino rats treated with methanolic extract of fruits of *Xylopiya aethiopic*". The difference in these results might be due to the different solvents used in the extraction of *Xylopiya aethiopic* fruit.

In this study, a dose-dependent increase was observed in the erythrocyte sedimentation rate (ESR) as presented in table 2. This result contradicted the findings of Onyebugu et al., [27] who investigated “the effects of dietary *Xylopiya aethiopicica* on hematological parameters and plasma lipids in male Wistar rats”. Saha et al. [30] reported that “increase in ESR could result from inflammation, pregnancy, anemia, autoimmune disorders (such as rheumatoid arthritis and lupus), infections, some kidney diseases and some cancers (such as lymphoma and multiple myeloma)”. This corroborated the adverse effect of *Xylopiya aethiopicica* fruit on liver [21] and kidney [22] indices recently reported.

## 5. CONCLUSION

*Xylopiya aethiopicica* significantly prolongs prothrombin time (PT) and activated partial thromboplastin time (aPTT) and in addition has thrombocytopenic property. These anticlotting properties have clearly elucidated and unraveled mechanisms of action which have laid credence to its use by women in flushing out remnants of product of conception post-delivery. Increase in ESR by *Xylopiya aethiopicica* may result from an inherent property in this fruit that is yet to be identified.

## DISCLAIMER

The products used for this research are commonly and predominantly use products in our area of research and country. There is absolutely no conflict of interest between the authors and producers of the products because we do not intend to use these products as an avenue for any litigation but for the advancement of knowledge. Also, the research was not funded by the producing company rather it was funded by personal efforts of the authors.

## ETHICAL APPROVAL

Animal Ethic committee approval has been taken to carry out this study.

## ACKNOWLEDGEMENT

Authors are grateful to Mr. Nsikan Malachy for his assistance in the practical aspect of this work.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Oloyede AM, Aduramigbe-Modupe AO. Antimicrobial activities of crude ethanolic extract of *Xylopiya aethiopicica*. International journal of current Reseach. 2011;3(10): 005-007.
2. Woode E, Ameyaw EO, Boakye-Gyasi E, Abotsi WK. Analgesic effects of an ethanol extract of the fruits of *Xylopiya aethiopicica* (Dunal) A. Rich (Annonaceae) and the major constituent, xylopic acid in murine models. Journal of Pharmacy & Bioallied Sciences. 2012;4(4):291-301.
3. Soh D, Nkwengoua E, Ngantchou I, Nyasse B, Denier C, Hannaert V. Xylopioxyde and other bioactive kaurenediterpenes from *Xylopiya aethiopicica* Dunal (Annonaceae). 2013
4. Igwe SA, Afonne, JC, Ghasi SI. Ocular dynamics of systemic aqueous extracts of *Xylopiya aethiopicica* (African guinea pepper) seeds on visually active volunteers. Journal of Ethnopharmacology. 2003;86(2-3):139-142.
5. Suleiman MM, Mamman M, Aliu YO, Ajanusi JO. Anthelmintic activity of the crude methanol extract of *Xylopiya aethiopicica* against *Nippostrongylus brasiliensis* in rats. Veterinarski arhiv. 2005;75(6):487.
6. Uwakwe A. *In vitro* antisickling effects of *Xylopiya aethiopicica* and *Monodora myristica*. Journal of Medicinal Plants Research. 2013;2(6):119-24.
7. Woode E, Alhassan A, Abaidoo CS. Effect of xylopic acid on sex hormones and spermatogenesis in male rats. Al Ameen j Med Sci. 2012;5(3):28-29.
8. Nwangwa EK. Antifertility Effects of Ethanolic Extract of *Xylopiya aethiopicica* on Male Reproductive Organ of Wistar Rats. American Journal of Medicine and Medical Sciences. 2012;2(1):12-15.
9. Busia K. Ghana herbal pharmacopoeia. Science and Technology Policy Research Institute, Council for Scientific and Industrial Research; 2007.
10. Yapi TA, Boti JB, Ahibo CA, Bighelli A, Castola V, Casanova J. Chemical variability of the leaf essential oil of *Xylopiya aethiopicica* (Dunal) A. Rich.from

- côte d'ivoire. Chemistry & Biodiversity. 2012;9(12):2802-2809.
11. Kadiri M, Ojewumi A, Onatade T. Indigenous Uses and phytochemical contents of plants used in the treatment of menstrual disorders and after-child birth problems In Abeokuta South Local Government area of Ogun State, Nigeria. Journal of Drug Delivery and Therapeutics. 2015;5(3):33-42.
  12. Konning GH, Agyare C, Ennison B. Antimicrobial activity of some medicinal plants from Ghana. Fitoterapia. 2004;75(1):65-7.
  13. Van-Hai N. The use of medicinal plants as immunostimulants in aquaculture: A review. Aquaculture. 2015;446:88-96.
  14. Nwafor A, Adienbo MO, Egwurugwu JN. *In vivo* effects of *Xylopi aethiopia* on hemorheological parameters in Guinea pigs. African. Journal of Applied Zoology/ Environmental Biology. 2009;11:79-81.
  15. Esekhiagbe M, Agatemor MMU, Agatemor C. Phenolic content and antimicrobial potentials of *Xylopi aethiopia* and *Myristica argentea*. Macedonian Journal of Chemistry and Chemical Engineering. 2009;28(2):159-162.
  16. John-Dewole O, Agunbiade S, Alao O, Arojojoye O. Phytochemical and antimicrobial studies of extract of the fruit of *Xylopi aethiopia* for medicinal importance. Journal of Biotechnology and Pharmaceutical Research. 2012;29(6): 118-122.
  17. Asekun O, Kunle O. The chemical constituents of the fruit essential oil of *Xylopi aethiopia* (Dunal) A. Rich from Nigeria. Journal of Essential Oil Bearing Plants. 2004;7(2):186-189.
  18. Nwaichi E, Igbino baro O. Effects of some selected spices on some biochemical profile of Wistar albino rats. American Journal of Environmental Engineering. 2012;2(1):8-11.
  19. Ogbuagu EO, Nweke IN, Uneke PC, Airaodion AI, Ogbuagu U. Weight gain reduction and hypoglycemic effects of *Xylopi aethiopia* fruit extract on Wistar rats. International Journal of Research and Reports in Hematology. 2020;5(3):1-8.
  20. Ogbuagu EO, Uneke PC, Airaodion AI, Nweke IN, Ogbuagu U. Hypolipidemic propensity of ethanolic extract of *Xylopi aethiopia* fruit in Wistar rats. Asian Journal of Research in Cardiovascular Diseases. 2020;3(5):1-11.
  21. Ogbuagu EO, Uneke PC, Airaodion AI, Nweke IN, Ogbuagu U. Hepatotoxic effect of *Xylopi aethiopia* fruit in Wistar rats. International Research Journal of Gastroenterology and Hepatology. 2021; 4(1):1-16
  22. Ogbuagu EO, Airaodion AI, Ogbuagu U, Nweke IN, Uneke PC. Nephrotoxicity of ethanol extract of *Xylopi aethiopia* fruit in Wistar rats. International Journal of Advances in Nephrology Research. 2021;4(1):1-16.
  23. Ogbuagu EO, Ogbuagu U, Airaodion AI, Uche CL, Ezirim EO, Nweke IN, Uneke PC. Effect of Ethanol Extract of *Xylopi aethiopia* Fruit on Oxidative Stress Indices of Wistar Rats. Asian Journal of Immunology. 2022;6(1):40-51.
  24. Ogbuagu EO, Airaodion AI, Ogbuagu U, Ezirim EO, Nweke IN, Uneke PC. Ethanol Extract of *Xylopi aethiopia* (African Negro Pepper) Fruit adversely perturbed Semen Qualities in Male Wistar Rats. International Journal of Research and Report in Gynaecology. 2022;7(2):43-52
  25. Ogbuagu EO, Ogbuagu U, Uneke PC, Nweke IN, Airaodion AI. Qualitative determination of the phytochemical composition of ethanolic extract of *Xylopi aethiopia* fruit. Asian Journal of Medical Principles and Clinical Practice. 2020; 4(3):1-5
  26. Airaodion AI, Ekenjoku JA, Ogbuagu EO, Okoroukwu VN, Ogbuagu U. *Carica papaya* leaves might cause miscarriage. Asian Research Journal of Gynaecology and Obstetrics. 2019;2(2):1-9.
  27. Onyebuagu PC, Pughikumo DT, Aloamaka CP. Effects of dietary *Xylopi aethiopia* on hematological parameters and plasma lipids in male Wistar rats. International Journal of Basic, Applied and Innovative Research. 2014;3(1):29 – 34.
  28. Ogbuagu EO, Airaodion AI, Ogbuagu U, Uche CL, Ezirim EO, Uneke PC, Nweke IN. *Xylopi aethiopia* fruit extract elevated red blood cell parameters but reduced white blood cell parameters in Wistar rats. International Journal of Medical and Alternative Medicine. 2022;4(6):58-65
  29. Obembe AO, Ofutet EO, Okpo-ene AI, Okon VE, Eyong EE. The effect of aqueous extract of *Xylopi aethiopia* (Nigro pepper) on some haematological parameters in albino rats. World Journal of

- Pharmaceutical Research. 2015;4(10): 2576-2583.
30. Saha AK, Schmidt BR, Wilhelmy J, Nguyen V, Do J, Suja VC, Nemat-Gorgani M, Ramasubramanian AK, Davis RW. Erythrocyte deformability as a potential biomarker for chronic fatigue syndrome. Blood. 2018;132(1):4874.

© 2022 Ogbuagu et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*

*<https://www.sdiarticle5.com/review-history/86463>*