



Unification of Extragalactic Radio Sources and Re-Acceleration of Radio Jets

Ezeugo Jeremiah Chukwuemerie ^{a*}

^a Department of Physics and Industrial Physics, NnamdiAzikiwe University, Awka, Anambra State, Nigeria.

Author's contribution

The sole author designed, analysed, interpreted and prepared the manuscript.

Article Information

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/83815>

Original Research Article

Received 20 January 2022
Accepted 30 March 2022
Published 02 April 2022

ABSTRACT

We use statistical methods of analysis to show that Large extended extragalactic radio sources (large EGRS) and compact steep spectrum (CSS) sources are similar sources evolving in different environments. Because of the diversity of their ambient media, one suffers higher magnitude of jet re-acceleration than the other. Linear regression analysis of observed source linear sizes, \mathcal{D} , of large extended extragalactic radio quasars against their individual observed redshifts, z , was carried out. With appreciable correlation coefficient, result shows that \mathcal{D} relates with redshift(z) according to the equation, $\mathcal{D} = \mathcal{K}(1 + z)^{-1.5}$; where \mathcal{K} is a constant. The relation shows that projected source size is smaller at earlier epoch in the evolution of the universe. In addition, if we take \mathcal{D} to be distance between any two points in space, then the relation may mean that the diameter of the universe is smaller at earlier epoch of its evolution. Moreover, with some plausible assumption, we find that jet velocity may be written as, $v_j \sim [\mathcal{P}_{BH}(1 + z)^3]^{0.5}$. This indicates that mechanism of jet production derives from the source central core (presumed to house a super massive blackhole), as well as, from cosmic evolution. Since cosmic evolution is propelled by dark energy, the relation simply shows that dark energy enhances jet expansion more than blackhole power. Dark energy creates more empty spaces. It is anti-gravity in nature, and is believed to be the driving force behind cosmic dilation. Moreover, we estimate the magnitude of the effect of dark energy on the jet expansion to be three times the effect caused by blackhole power. Therefore, this result suggests that in addition to other factors, a culprit for re-acceleration mechanism for radio jets is dark energy. In order to check the authenticity of the assertion, we carry out linear regression analyses of

*Corresponding author: E-mail: chuksemerie@yahoo.com;

observed luminosities, (\mathcal{P}), against observed redshifts for both the extended extragalactic radio quasars and CSS quasars. With good correlations in both cases, we obtain $\mathcal{P} \sim (1+z)^a$; where $a = 0.04$ and 4.33 for the extended quasars and CSS quasars respectively. The results show strong dependence of source luminosity on redshift. Moreover, the staggering difference in their indices suggestively implies that if we match the two samples at similar redshifts, luminosities of CSS quasars will be higher than those of the more extended sources. Hence, the results show that at similar epoch, luminosities of the more extended quasars are more extinguished (or more attenuated) than those of the CSS quasars. The jets and lobes of the larger/extended quasars are located in the intergalactic medium (IGM). This medium is very much rarefied and dark energy should be expected to exhibit more effect there than in the interstellar medium (ISM). On the other hand, the components of the CSS sources are sub-galactic. This means that they are buried in the dense ISM. In this medium, there is lesser manifestation of dark energy. Therefore, the disparity in the two indices (0.04 and 4.33) simply indicates manifestation of dark energy. Due to more rapid expansion of the IGM in which the jets and lobes of the extended sources are located, their luminosities are diluted more by more spaces created by dark energy in that medium. This shows that Large EGRS and CSS sources are similar sources evolving in different ambient media. Therefore, because of the diversity of their ambient media, large EGRS suffers higher magnitude of jet re-acceleration than the CSS source. In addition to this, we may state that dark energy is a suspect in enhancement of re-acceleration of the extragalactic jets. In conclusion, our results suggest that in addition to being similar sources evolving in different ambient media, a factor which fuels re-acceleration of extragalactic radio jets is dark energy.

Keywords: Cosmic evolution; linear size; astronomical distance; luminosity; radio sources; quasars; dynamical evolution; re-acceleration; dark energy.

1. INTRODUCTION

Extragalactic radio sources (EGRS) emit large amount of radio radiation. These are those sources with high ratio of radio to optical emission. This ratio is generally defined by the quotient of the two flux densities given by $S_{5\text{ GHz}}/S_{6 \times 10^5\text{ GHz}} > 10$ [1–5]. They are found outside the Milky Way galaxy. They are mainly made up of the following sub-classes: radio galaxies and radio-loud quasars [1–12]. Radio emission from these sources commonly takes the morphology of two opposite sided relativistic jets emanating from the central core presumed to harbour a super massive blackhole. The jets terminate with strong shocks forming lobes. The two radio-emitting lobes straddling the central component which is more or less coincident with the nucleus of the host galaxy is irregular in shape [1,4,7]. In some sources, the lobes contain hotspots believed to be the termination points of the radio jets [1,4,7].

The more extended EGRS have linear sizes, D , given by $D > 30\text{ Kpc}$ assuming Hubble constant, $H_0 = 75\text{ kms}^{-1}\text{Mpc}^{-1}$. Since 30 Kpc is roughly the diameter of a typical galaxy, then it simply means that the jets and lobes of these sources are located in the intergalactic media. Their radio luminosity is in excess of 10^{26} W at 5 GHz and

overall luminosities are found to be $\geq 10^{37}\text{ W}$ [4–11].

However, the compact steep spectrum (CSS) sources are scaled-down versions of the more extended sources with projected linear sizes well below 30 Kpc . They comprise radio galaxies and radio-loud quasars but on sub-galactic dimensions. They show steep spectra ($\alpha \geq 0.5, S_\nu \sim \nu_p^\alpha$; where S_ν is flux density, α , spectral index, and ν_p , spectral turnover) at high frequencies (e.g. $\geq 0.02\text{ GHz}$) from the entire radio morphological structures; hence, their name. Their radio luminosities at 5 GHz are above 10^{21} W ; while their bolometric luminosities are higher than 10^{37} W [12–18].

A number of authors have wondered on the relationship between CSS sources and larger EGRS. Sequel to this, some authors have come up with some models for the evolution of these sources. These models include: Youth Scenario (i.e. young evolving sources), Frustration Scenario (i.e. sources confined by dense gases in their respective interstellar media), and Relativistic Beaming and Orientation Effects (i.e. the source sizes are foreshortened by orientation and projection effects) [12–18].

It has been pointed out by some authors that presence of jets in radio sources generally

depicts presence of gaseous ambient media [12–18]. Some hydrodynamic simulations of jet propagations in dense media have been carried out to investigate their properties and nature [5–6]. Some of the results in literature have shown that the jets are made up of fast moving particles (presumably electrons) [12–13,19].

In addition to the foregoing, many authors have been puzzled on the mechanism behind the un-attenuated nature of the speeds with which the jet particles propagate through the immense distances from the central core to the radio-emitting lobes. Some authors have suggested that there must be some re-acceleration mechanisms. This shows that the particles' kinetic energies may be boosted by some mechanisms while on their way to the lobes [4]. It is generally believed that if this is true, then layers of magnetic fields present in the intergalactic media should be a suspect [4]. However, this has been a presumption and there is no concrete evidence for the assertion. Therefore, in this paper, we show that dark energy is a culprit behind this re-acceleration phenomenon; and that Large extended extragalactic radio sources and compact steep spectrum sources are similar sources evolving in different environments. Because of the diversity of their ambient medium densities, one suffers higher magnitude of jet re-acceleration than the other.

This is done by carrying out some regression analyses using samples of some powerful extragalactic radio sources. The first sample

contains large extended extragalactic radio-loud quasars obtained from [20]. Their projected linear sizes are well above 30 Kpc, and are 170 in number. The second sample contains 27 CSS quasars obtained from O'Dea (1998)[15].

2. SIZE/REDSHIFT (\mathcal{D}/z) RELATION

In this section, we carry out linear regression analysis of source projected linear sizes of quasars and observed redshifts (see Fig. 1).

Results from the $\mathcal{D} - z$ plane shows that projected linear size and source redshift are connected according to the following relation:

$$\text{Log}\mathcal{D} = 2.562 - 1.522\text{Log}(1 + z) \quad (1)$$

If we assume that the correlation is appreciable (with correlation coefficient given as 0.41), we may rewrite equation (1) to be

$$\mathcal{D} = \mathcal{K}(1 + z)^{-1.5} \quad (2)$$

where, \mathcal{K} is a constant.

Or evaluating further, we obtain

$$z \sim \mathcal{D}^{-2/3} - 1 \quad (3)$$

Equation (3) shows that projected source size is smaller at earlier epoch. In addition to that, if we take \mathcal{D} to be distance between any two points in space, then equation (3) may mean that the diameter of the universe is smaller at earlier epoch.




Fig. 1. The scatter plot of source observed linear sizes against observed redshifts for extended quasars

Assuming that the radio jet is confined by ram-pressure balance with the ambient medium, we have [16,20, 21]

$$\mathcal{P}_j \approx \eta m_h v_j^2 \quad (4)$$

where \mathcal{P}_j is jet internal pressure, η is particle number density of the source ambient medium, m_h is hydrogen mass, and v_j is jet velocity. From (4), we derive blackhole power, \mathcal{P}_{BH} , (i.e. power supplied by the blackhole to the jet). This is given by

$$\mathcal{P}_{BH} \approx \eta m_h c v_j^2 \Gamma \mathcal{D}^2 \quad (5)$$

Evaluating for jet velocity, we obtain

$$v_j \approx \left(\frac{\mathcal{P}_{BH}}{\eta m_h c \Gamma \mathcal{D}^2} \right)^{\frac{1}{2}} \quad (6)$$

Combining equations (2) and (6), we get

$$v_j \approx \left(\frac{\mathcal{P}_{BH}}{\eta m_h c \Gamma [\mathcal{K}(1+z)^{-1.5}]^2} \right)^{\frac{1}{2}} \quad (7)$$

This implies

$$v_j \sim [\mathcal{P}_{BH}(1+z)^3]^{\frac{1}{2}} \quad (8)$$

This therefore indicates that jet expansion is powered by both blackhole and cosmic evolution. Since cosmic evolution is propelled by dark energy (i.e., the intrinsic tendency of vacuum/free space to expand in volume), equation (8) simply states that dark energy enhances jet expansion

more than blackhole power. Dark energy creates more empty spaces. It is anti-gravity, and is believed to be the driving force behind the evolution (or expansion) of the universe [22]. Using the indices in the equation, we estimate the magnitude of the effect of dark energy on the jet expansion to be three times the effect caused by blackhole power. Therefore, we suggest that in addition to other factors, a culprit for reacceleration mechanism for radio jets may be dark energy.

3. LUMINOSITY/REDSHIFT (\mathcal{P}/z)

3.1 Relation

We also carry out linear regression analyses of observed luminosities, \mathcal{P} , against observed redshifts for both the extended extragalactic radio quasars (Fig. 2) and CSS quasars (Fig. 3).

From $\mathcal{P} - z$ data (Fig. 2), we find the relation,

$$\text{Log } \mathcal{P} = 1.64 + 0.04 \text{Log } z \quad (9)$$

With correlation coefficient given as $r = 0.8$, we see that source luminosity shows strong correlation with redshift. Transforming the equation, we obtain

$$\mathcal{P} \sim (1+z)^{0.04} \quad (10)$$

Moreover, from $\mathcal{P} - z$ plane for CSS quasars (Fig. 3), we obtain similar relation. This is given by

$$\text{Log } \mathcal{P} = 26.46 + 4.33 \text{Log } z \quad (11)$$




Fig. 2. The scatter plot of source observed luminosities against observed redshifts for the extended quasars




Fig. 3. The scatter plot of source observed luminosities against observed redshifts for the CSS quasars

It shows good correlation too with $r \approx 0.8$. rewriting the equation, we have

$$P \sim (1+z)^{4.33} \tag{12}$$

Generally, the results (10) and (12) obtained from the analyses of the two samples simply indicate strong dependence of source luminosity on redshift. Moreover, the staggering difference in their indices suggestively implies that if we match the two samples at similar redshifts, luminosities of CSS quasars will be higher than those of the more extended sources just as is roughly shown in Fig. 4.




Fig. 4. Schematic diagram of variation of luminosities of CSS quasars and larger EGR quasars when matched at similar redshifts (Solid line represents CSS quasars' luminosities while broken line shows luminosities of the more extended EGRs).

This result shows that at similar epoch, luminosities of the more extended sources are

more extinguished/attenuated than those of the CSS sources. It is good to note that components (jets and lobes) of the larger sources are located in the intergalactic medium (IGM). This medium is very much rarefied and dark energy shows more effect here. However, the components of the CSS sources are sub-galactic. This means that they are encapsulate by the dense gases in the interstellar media (ISM). In this medium, there is lesser manifestation of dark energy. Therefore, the disparity in the two indices simply indicates manifestation of dark energy. Due to more rapid expansion of the IGM in which the jets and lobes of the larger sources are located, their luminosities are diluted more by more spaces created by dark energy in that medium. This also shows that dark energy is a suspect in enhancement of re-acceleration of the extragalactic jets.

4. DISCUSSION

We use statistical methods of analysis to show that Large extended extragalactic radio sources and compact steep spectrum (CSS) sources are similar sources evolving in different environments. Because of the diversity of their ambient medium densities, larger sources suffer higher magnitude of jet re-acceleration than the CSS sources.

We have proved this by carrying out linear regression analysis of observed source linear sizes, \mathcal{D} , of the more extended extragalactic quasars against their individual observed redshifts, z , (see Fig. 1). Result shows that \mathcal{D} relates with redshift (z) according to the equation, $\mathcal{D} = \mathcal{K}(1+z)^{-1.5}$;

where \mathcal{K} is a constant, $r = 0.4$ is the correlation coefficient. Even though the correlation is marginal, we consider it good enough for observed data in the field of astronomy. The relation shows that projected source size is smaller at earlier epoch in the evolution of the universe. Moreover, if we take \mathcal{D} to be distance between any two points in space, then the relation may mean that the diameter of the universe is smaller at earlier epoch of its evolution.

If we assume jet confinement by ram-pressure balance with the particles of the medium through which it propagates [16,20,21], then jet velocity may be given by, $v_j \sim [\mathcal{P}_{BH}(1+z)^3]^{0.5}$. This indicates that jet velocity is powered by blackhole and cosmic evolution. Since cosmic evolution is propelled by dark energy, equation (8) simply states that dark energy enhances jet expansion more than blackhole power. Dark energy creates more empty spaces. It is anti-gravity in nature, and is believed to be the driving force behind cosmic dilation [22]. Using the indices in the equation, we estimate the magnitude of the effect of dark energy on the jet expansion to be three times the effect caused by blackhole power. Therefore, this result suggests that in addition to other factors, a culprit for reacceleration mechanism for radio jets may be dark energy.

In addition, we carry out linear regression analyses of observed luminosities, (\mathcal{P}), against observed redshifts for both the extended extragalactic radio quasars (Fig. 2) and CSS quasars (Fig. 3).

From $\mathcal{P} - z$ data (Fig. 2) for the more extended sources, we find the relation, $\mathcal{P} \sim (1+z)^{0.04}$, with correlation coefficient given as 0.8. This shows that source luminosity shows strong correlation with redshift. Moreover, from $\mathcal{P} - z$ plane for the CSS quasars (Fig. 3), we obtain similar relation, $\mathcal{P} \sim (1+z)^{4.33}$, with 0.8 as the correlation coefficient.

The results (10) and (12) obtained from the analyses of the two samples simply shows strong dependence of source luminosity on redshift. Moreover, the staggering difference in their indices suggestively implies that if we match the two samples at similar redshifts, luminosities of CSS quasars will be higher than those of the more extended sources (see Fig. 4). This result shows that at similar epoch, luminosities of the more extended quasars are more extinguished

(or more attenuated) than those of the CSS quasars.

The components (i.e., jets and lobes) of the larger sources are located in the intergalactic medium (IGM). This medium is very much rarefied and dark energy should be expected to exhibit more effect there than in the interstellar medium (ISM). On the other hand, the components of the CSS sources are sub-galactic. This means that they are buried in the dense ISM. In this medium, there is lesser manifestation of dark energy. Therefore, the disparity in the two indices in equations (10) and (12) simply indicates manifestation of dark energy. Due to more rapid expansion of the IGM in which the jets and lobes of the larger sources are located, their luminosities are diluted more by more spaces created by dark energy in that medium. This also shows that dark energy is a suspect in enhancement of re-acceleration of the extragalactic jets.

5. CONCLUSION

In conclusion, our results suggest that dark energy is a suspect behind re-acceleration phenomenon; and that Large extended extragalactic radio sources and compact steep spectrum sources are similar sources evolving in media of different densities. Because of this difference in their ambient medium densities, larger sources are subjected more to higher magnitudes of jet re-acceleration than the CSS sources.

CONSENT AND ETHICAL APPROVAL

It is not applicable.

COMPETING INTERESTS

Author has declared that no competing interests exist.

REFERENCES

1. Ezeugo JC. Jet in the more extended radio sources and unification with compact steep spectrum sources. The Pacific Journal of Science and Technology. 2021;22:14–19.
2. Ubah OL, Ezeugo JC. Relativistic jet propagation: Its evolution and linear size cosmic dilation. International Astronomy and Astrophysics Research Journal. 2021;3(3):1–6.

3. Urry CM. AGN Unification: An update. *Astronomical Society of the Pacific conference series* 1; 2004.
4. Readhead AC. Evolution of powerful extragalactic radio sources. In *proc. Colloquium on Quasars and Active Galactic Nuclei*, ed. Kohen, M., and Kellermann, K. (USA: National Academy of Sciences, Berkman Center, Irvine). 1995;92:11447–11450.
5. Robson I. *Active galactic nuclei*, John Wiley and Sons Ltd, England; 1996.
6. Jackson JC. Radio source evolution and unified schemes. *Publications of Astronomical Society of the Pacific*. 1999;16:124–129.
7. Kawakatu N, Kino M. The velocity of large-scale jets in a declining density medium. In *Serie de Conferencias. Triggering Relativistic Jets*, ed. W.H. Lee and E. Ramirez-Ruiz. 2007;27:192–197.
8. Ezeugo JC. On the intergalactic media densities, dynamical ages of some powerful radio sources and implications. *Journal of Physical Sciences and Application*. 2021;11(1):29–34.
9. Mahatma VH, Hardcastle MJ, Williams WL. LoTSS DR1: Double-double Radio Galaxies in the HETDEX Field. *Astronomy and Astrophysics*. 2019;622:A13.
10. Mingo B, Croston JH, Hardcastle MJ. Revisiting the Fanaroff-Riley Dichotomy and Radio Galaxy Morphology with the LOFAR Two-Meter Sky Survey (LoTSS). *Monthly Notices of the Royal Astronomical Society*. 2019;488:2701-2721.
11. Hardcastle MJ, Williams WL, Best PN. Radio-loud AGN in the First LoTSS Data Release — The Lifetimes and Environmental Impact of Jet-Driven Sources. *Astronomy and Astrophysics*. 2019;622:A12.
12. Dabhade P, Gaiwad M, Bagchi J. Discovery of giant radio galaxies from NVSS: Radio and infrared properties. *Monthly Notices of the Royal Astronomical Society*. 2017;469(3):2886–2906.
13. Ezeugo JC. Compact steep-spectrum radio sources and ambient medium density. *International Journal of Astrophysics and Space Science*. 2015;3(1):1–6.
14. Ezeugo JC. On the dependence of spectral turnover on linear size of compact steep-spectrum radio sources. *International Journal of Astrophysics and Space Science*. 2015;3(2):20–24.
15. Ezeugo JC, Ubachukwu AA. The spectral turnover–linear size relation and the dynamical evolution of compact steep spectrum sources. *Monthly Notices of the Royal Astronomical Society*. 2010;408:2256–2260.
16. O’Dea CP. The Compact Steep Spectrum and Gigahertz peaked spectrum radio sources. *Publications of the Astronomical Society of the Pacific*. 1998;110:493–532.
17. Fanti C, Fanti R, Dallacasa D, Schillizzi RT, Spencer RE, Stanghellini C. Are compact steep spectrum sources young? *Astronomy and Astrophysics*. 1995;302:317–326.
18. Ezeugo JC. Compact spectrum source size and cosmological implication. *Journal of Research in Applied Mathematics*. 2021;7(2):1–4.
19. Nilsson K. Kinematical models of double radio sources and unified scheme. *Monthly Notices of the Royal Astronomical Society*. 1998;132:31–37.
20. Ezeugo JC. On the components of large extended extragalactic radio sources. *The Pacific Journal of Science and Technology*. 2021;22(1):20–23.
21. Cavalho JC. The evolution of GHZ-peaked spectrum radio sources. *Astronomy and Astrophysics*. 1998;329:845–852.
22. Friedman JA, Turner MS, Huterer D. Dark energy and the accelerating universe *annual review of astronomy*. 2008;46:385–432.

© 2022 Chukwuemerie; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/83815>