



# Field Screening of Garlic Genotypes for Identification of Resistant Sources against Purple Blotch Disease

K. N. Vijaykumar <sup>a\*</sup>, Shripad Kulkarni <sup>b</sup>, S. M. Hiremath <sup>c</sup>,  
T. R. Shashidhar <sup>c</sup>, D. N. Kambrekar <sup>d</sup>, Vijay Mahajan <sup>e</sup>,  
Anil Khar <sup>f</sup> and P. V. Patil <sup>b</sup>

<sup>a</sup> ICAR-IARI, Regional Research Centre, Dharwad-580 005, Karnataka, India.

<sup>b</sup> Department of Plant Pathology, University of Agricultural Sciences, Dharwad-580 005, Karnataka, India.

<sup>c</sup> Department of Horticulture, University of Agricultural Sciences, Dharwad-580 005, Karnataka, India.

<sup>d</sup> Department of Agricultural Entomology, University of Agricultural Sciences, Dharwad-580 005, Karnataka, India.

<sup>e</sup> ICAR-Directorate of Onion and Garlic, Rajgurunagar, Pune, India.

<sup>f</sup> ICAR-Indian Agricultural Research Institute, New Delhi, India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJPSS/2023/v35i143059

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/100352>

Original Research Article

Received: 22/03/2023

Accepted: 26/05/2023

Published: 07/06/2023

## ABSTRACT

Garlic (*Allium sativum* L.) is one of the oldest cultivated plants in the world and belongs to the family Amaryllidaceae. Purple blotch caused by *Alternaria porri* (Ellis) Ciferri is one such disease which causes a great deal of loss for farmers. The best method to manage purple blotch is to adopt resistant genotypes for cultivation, which is highly necessary in areas with a high incidence. In order to identify resistant sources, 37 genotypes were field screened under artificial epiphytotic

\*Corresponding author: E-mail: vijaykn549@gmail.com;

conditions at AINRPOG (All India Network Research Project on Onion and Garlic), MARS (Main Agricultural Research Station), Dharwad during Rabi, 2019 and 2020. Results from two years showed that no genotype was immune to purple blotch. Among the genotypes tested, five genotypes viz., G50, Bhima Purple, DOGR426, DOGR543 and DOGR569 were resistant to infection (1-10%), 17 genotypes moderately resistant to infection (11-20%), while 4 genotypes were moderately susceptible to infection (21-40%). On a scale of 0 to 5, only one genotype (HRG Local) showed susceptible (41-75%) in reaction. Among the remaining genotypes, disease reactions differed between years. The genotypes that are resistant can be utilized by breeders to produce high yielding purple blotch resistant varieties.

**Keywords:** Purple blotch; artificial screening; per cent disease index; disease reaction.

## 1. INTRODUCTION

Garlic (*Allium sativum* L.) is one of the most popular and the oldest cultivated plants in the world belong to the family Amaryllidaceae. Garlic bulb comprises of water (84.09%), organic matter (13.38%) and inorganic matter (1.53%) along with 0.1 per cent volatile oil [1]. The bulbs contain a colourless, odourless and water soluble compound called allicin. On crushing or chewing, enzyme allinase present in the raw garlic bulbs breaks down into allin to produce allicin which cannot be observed in cooked garlic. Diallyl disulfide possesses the true garlic odour [2]. Nevertheless, the crop is attacked by many diseases over its lifetime, resulting in low yields, resulting in economic losses. In recent years, purple blotch caused by *Alternaria porri* (Ellis) Ciferri has become one of the most serious diseases affecting garlic [3]. This foliar pathogen alters metabolic processes by reducing the photosynthetic activity. It has been reported that foliar infection can reach 90% in susceptible garlic cultivars [4]. The bulb shelf life is reduced as a result of leaf damage and blackening of scales at bulb maturity. The drying of leaves can significantly reduce bulb yield in garlic (25-60%).

A number of states, including Gujarat, Maharashtra, Andhra Pradesh, Haryana, Karnataka, Odisha, and Himachal Pradesh, have reported purple blotch in garlic as an economically important disease [5-10]. Initially, purple blotch appears as a multitude of tiny white spots that are irregular or circular in shape. Spots gradually grow larger, become oval-shaped or irregular, and eventually change from white to violet color. Later stages of development show the central portion of the spots changing to purple. This is surrounded by a pale yellow orange to salmon band beyond which is a pale

green zone. The dark purple colour is the most distinctive disease symptom. A distinct yellowing usually extends from both ends of the spots, often reaching the tips and bases of the leaves [11]. Host plant resistance is an effective, economical and environmentally safe component of an integrated approach. Therefore, an attempt was made to identify resistant sources to mitigating loss in farmers' field.

## 2. MATERIALS AND METHODS

Totally thirty seven garlic genotypes collected from ICAR-DOGR (Directorate of Onion and Garlic), Rajgurunagar, Pune and ICAR-IARI, New Delhi along with local genotypes (DWG1, HRG Local and Gadag Local) as commercial check were subjected for field screening during Rabi 2019 and 2020 against purple blotch at AINRPOG (All India Network Research Project on Onion and Garlic), MARS (Main Agricultural Research Station), Dharwad. All the genotypes were sown in two rows of two meter length with a plant spacing of 30 cm × 10 cm (row to row and plant to plant) in non-replicated augmented design. All the recommended agronomic practices were followed, except disease management practices. Inoculum of *Alternaria porri* ( $2 \times 10^6$  spores/ml) was uniformly sprayed during evening hours at 30 days after sowing to create uniform disease pressure throughout the field. The disease severity was recorded at 90 days after sowing using a scale of 0-5 [8] by randomly selecting ten plants in each genotype. Per cent disease index was calculated by using formula given by Wheeler [12]. Based on their disease severity, genotypes were categorized into immune (0%), resistant (1-10%), moderately resistant (11-20%), moderately susceptible (21-40%), susceptible (41-75%) and highly susceptible (75-100%).

**Chart 1. Disease scoring scale (0-5 grade) for purple blotch of garlic [8]**

Disease score	Per cent leaf area infected	Disease reaction
0	No disease symptoms	Immune
1	A few spots towards tip covering 10 per cent leaf area	Resistant
2	Several dark purplish brown patches covering 11 to 20 per cent leaf area	Moderately resistant
3	Several patches with pale outer zone covering 21 to 40 per cent leaf area	Moderately susceptible
4	Yellow streaks covering 41 to 75 per cent leaf area	Susceptible
5	Complete drying of the leaves or breaking of leaves from centre	Highly susceptible

### 3. RESULTS AND DISCUSSION

Totally 37 garlic genotypes were screened during Rabi 2019 and 2020 against purple blotch under artificial epiphytotic condition in the field to identify the resistant sources as described in "Material and methods" and data are presented in Table 1.

During Rabi 2019, none of the genotypes showed immune or highly susceptible reaction to purple blotch. Among 37 genotypes, seven genotypes (G50, Bhima Purple, DOGR81W, DOGR389, DOGR426, DOGR543 and DOGR569) showed resistant reaction, whereas, 23 genotypes (G282, GG2, GG4, Bhima Omkar, Godavari, Phule Baswant, DWG1, DOGR51, DOGR102, DOGR113, DOGR119, DOGR150, DOGR181, DOGR185, DOGR325, DOGR329, DOGR353, DOGR440, DOGR517, DOGR534, DOGR548, DOGR604 and DOGR744) showed moderately resistant reaction and six genotypes (G41, Gadag Local, DOGR75, DOGR228, DOGR409 and DOGR756) showed moderately susceptible reaction while remaining one genotype (HRG Local) was found susceptible to purple blotch of garlic.

During Rabi 2020, none of the genotypes showed immune or highly susceptible reaction to purple blotch. Among 37 genotypes, eight genotypes (G50, Bhima Purple, DOGR119, DOGR150, DOGR329, DOGR426, DOGR543 and DOGR569) showed resistant reaction whereas, 21 genotypes (G41, G282, GG2, Bhima Omkar, Phule Baswant, DWG1, DOGR51, DOGR75, DOGR81W, DOGR102, DOGR113, DOGR181, DOGR325, DOGR353, DOGR389, DOGR440, DOGR517, DOGR534, DOGR548, DOGR604 and DOGR744) showed moderately resistant reaction and seven genotypes (GG4, Godavari, Gadag Local, DOGR185, DOGR228, DOGR409 and DOGR756) showed moderately susceptible reaction while, remaining one

genotype (HRG Local) was found susceptible in reaction.

Thirty seven genotypes grouped into different reaction types based on the results obtained from two years. It was found that none of the genotypes were immune to purple blotch (Table 2). Five genotypes viz., G50, Bhima Purple, DOGR426, DOGR543 and DOGR569 showed resistant reaction over the seasons whereas, seventeen genotypes viz., G282, GG2, Bhima Omkar, Phule Baswant, DWG1, DOGR51, DOGR75, DOGR81W, DOGR102, DOGR113, DOGR181, DOGR325, DOGR353, DOGR389, DOGR440, DOGR517, DOGR534, DOGR548, DOGR604 and DOGR744 showed moderately resistant in reaction and four genotypes (Gadag Local, DOGR228, DOGR409 and DOGR756) showed moderately susceptible reaction while, only one genotype (HRG Local) showed susceptible reaction to purple blotch of garlic. However, no genotype was found to be highly susceptible to purple blotch.

Host plant resistance is one of the key strategies for reducing disease losses. Farmers can easily use this method of management, and it is relatively affordable and environmentally friendly. Moreover, resistant cultivars conserve natural resources and are more cost-effective, time-saving, and energy-efficient than other methods of disease control [13].

The results are consistent with previous reports showing that, although a few lines seem immune to purple blotch, the majority are resistant under natural infestation in open fields [14]. A study conducted by Sugha et al. [15] evaluated 94 onion genotypes under natural conditions and found that only two varieties, IC39178 and IC49371, were resistant to purple blotch. Furthermore, Behera et al. [16] observed VG-18 to be a resistant cultivar and another 12 lines to be moderately resistant to purple blotch. It is

**Table 1. The response of garlic genotypes to purple blotch caused by *Alternaria porri* during Rabi, 2019 and 2020**

Sl. No.	Genotypes	2019		2020		Sl. No.	Genotypes	2019		2020	
		Per cent disease index	Disease grade	Per cent disease index	Disease grade			Per cent disease index	Disease grade	Per cent disease index	Disease grade
1	G41	23.14	3	15.33	2	20	DOGR181	13.74	2	17.32	2
2	G50	9.65	1	8.43	1	21	DOGR185	18.97	2	26.56	3
3	G282	12.33	2	13.87	2	22	DOGR228	32.63	3	29.97	3
4	GG2	13.54	2	14.21	2	23	DOGR325	16.74	2	18.72	2
5	GG4	14.21	2	25.52	3	24	DOGR329	16.78	2	9.87	1
6	Bhima Omkar	16.38	2	15.96	2	25	DOGR353	14.23	2	15.39	2
7	Bhima Purple	8.65	1	8.23	1	26	DOGR389	9.54	1	15.59	2
8	Godavari	16.25	2	22.46	3	27	DOGR409	38.65	3	35.43	3
9	Phule Baswant	13.92	2	14.21	2	28	DOGR426	8.92	1	7.71	1
10	Gadag Local	26.53	3	31.22	3	29	DOGR440	18.28	2	19.92	2
11	HRG Local	45.23	4	46.76	4	30	DOGR517	12.21	2	13.36	2
12	DWG1	19.54	2	18.79	2	31	DOGR534	15.37	2	14.71	2
13	DOGR51	16.43	2	15.32	2	32	DOGR543	6.77	1	7.61	1
14	DOGR75	32.21	3	35.68	2	33	DOGR548	13.22	2	15.46	2
15	DOGR81W	8.97	1	15.33	2	34	DOGR569	9.34	1	8.65	1
16	DOGR102	16.47	2	15.55	2	35	DOGR604	14.68	2	15.78	2
17	DOGR113	13.23	2	14.90	2	36	DOGR744	15.44	2	16.70	2
18	DOGR119	12.23	2	9.87	1	37	DOGR756	32.11	3	34.36	3
19	DOGR150	15.38	2	9.54	1						

**Table 2. Grouping of garlic genotypes based on reaction to purple blotch caused by *Alternaria porri***

Disease grade	Per cent leaf infection	Genotypes	Disease reaction	No. of Genotypes
0	0	-	Immune	0
1	1-10	G50, Bhima Purple, DOGR426, DOGR543 and DOGR569	Resistant	5
2	10-20	G282, GG2, Bhima Omkar, Phule Baswant, DWG1, DOGR51, DOGR102, DOGR113, DOGR181, DOGR325, DOGR353, DOGR440, DOGR517, DOGR534, DOGR548, DOGR604 and DOGR744	Moderately Resistant	17
3	21-40	Gadag Local, DOGR228, DOGR409 and DOGR756	Moderately susceptible	4
4	40-75	HRG Local	Susceptible	1
5	More than 75	-	Highly susceptible	Nil

consistent with the findings of Agarwal and Tiwari [17], Nandini et al. [18], Kowser et al. [19] and Vijaykumar et al. [2,20] who evaluated different garlic genotypes for purple blotch and found that the per cent disease index was negatively correlated with bulb yields and storage quality. Although field screening showed variable responses to purple blotch, it is necessary to evaluate their efficiency as pre-breeding lines by evaluating them in multiple hotspot areas for purple blotch.

#### 4. CONCLUSION

A total of 37 garlic genotypes were screened against purple blotch under artificial epiphytic conditions and the results revealed that five genotypes showed resistant while seventeen genotypes showed moderately resistant in reaction. In contrast, four of the genotypes showed moderately susceptible reaction and one genotype showed susceptible reaction while none of the genotypes showed highly susceptible reaction to purple blotch. Farmers can cultivate these resistant or moderately resistant genotypes instead of local cultivars to get higher bulb yields when combating purple blotch disease.

#### DISCLAIMER

Some part of this manuscript was previously presented in the conference: 3rd International Conference IAAHAS-2023 "Innovative Approaches in Agriculture, Horticulture & Allied Sciences" on March 29-31, 2023 in SGT University, Gurugram, India. Web Link of the proceeding: <https://wikifarmer.com/event/iaahas-2023-innovative-approaches-in-agriculture-horticulture-allied-sciences/>

#### ACKNOWLEDGEMENTS

The authors are grateful to the members of AINRPOG and AICRP on Potato, University of Agricultural Sciences, Dharwad for providing the necessary facilities and timely assistance in the conduct of experiment.

#### COMPETING INTERESTS

Authors have declared that no competing interests exist.

#### REFERENCES

- Chalson M, McFerren M. Garlic: What we know and what we don't know arch. Intern. Med. 2007;167(4):325-326.
- Vijaykumar KN, Shripad Kulkarni, Patil PV, Kambrekar DN, Shashidhar TR. Eco-friendly management of purple blotch of garlic caused by *Alternaria porri* (Ellis) Ciferri. Biol. Forum Int. J. 2022a;14(2):1-7.
- Vijaykumar KN, Shripad Kulkarni, Patil PV, Kambrekar DN, Shashidhar TR. *In vitro* evaluation of new molecules of fungicides against purple blotch *Alternaria porri* (Ellis) Ciferri of garlic (*Allium sativum* L.). Pharma Innov. J. 2021;10(12):1048-1054.
- Bisht IS, Agarwal RC. Susceptibility to purple blotch (*Alternaria porri*) in garlic. Ann. App. Biol. 1993;122:31-38.
- Raju KS. Studies on purple blotch of onion (*Allium cepa* L.) caused by *Alternaria porri*. M. Sc. (Agri.) Thesis, Saurashtra Univ. Junagadh, Gujarat; 1970.
- Bhangale GT, Joi MB. Role of thrips in development of purple blotch of onion. J.

- Maharashtra Agric. Univ. 1983;8(3):299-300.
7. Gupta RP, Srivastava PK, Pandey UB. Control of purple blotch disease of onion seed crop. Indian Phytopath. 1986;39(1):303-304.
  8. Sharma SR. Effect of fungicidal sprays on purple blotch and bulb yield of onion. Indian Phytopath. 1986;39:78-82.
  9. Mishra D, Mahanta IC, Chhotaray PK. Chemical control of purple blotch of onion in Odisha. J. Agric. Res. 1989;2(1):25-28.
  10. Sugha SK, Tyagi PD. Effectiveness of metalaxyl + mancozeb against purple blotch (*Alternaria porri*) of onion (*Allium cepa* L.) in Himachal Pradesh. Indian J. Agric. Sci. 1994;64(2):135-136.
  11. Aveling TAS. Purple blotch (*Alternaria porri*) of onion. Recent Res. Dev. Plant Pathol. 1998;2:63-76.
  12. Wheeler BEJ. An introduction to plant disease. John Wiley and Sons Ltd, London. 1969;301.
  13. Roopadevi, Patil PV. *In vitro* bioassay of fungicides, bioagents, botanicals and itk's against *Pyricularia grisea* (Cooke) Sacc. incitant of pearl millet blast. Int. J. Pure App. Biosci. 2017;5(5):1457-1463.
  14. Pathak DP, Singh A, Despande A, Sridar TT. Source of resistance to purple blotch in onion. Veg. Sci. 1986;13:300-303.
  15. Sugha SK, Develash RK, Tyagi PD. Performance of onion genotypes against purple blotch pathogen. South Indian Hort. 1992;40:297.
  16. Behera S, Santra P, Chattopadhyay S, Das S, Maity TK. Variation in onion varieties for reaction to natural infection of *Alternaria porri* (Ellis) Cif. and *Stemphylium vesicarium* (Wallr.). Bioscan. 2013;8(3):759-761.
  17. Agarwal A, Tiwari RS. Evaluation of garlic (*Allium sativum*) genotypes for yield and susceptibility to purple blotch. J. Hortic. For. 2013;5(4):48-52.
  18. Nandini KS, Umamaheswarappa P, Srinivasa V, Abhishek KN, Sindhu K, Lavanya KS. Performance of garlic (*Allium sativum* L.) genotypes for yield and quality attributes under central dry zone of Karnataka. J. Pharmacogn. Phytochem. 2018;3:329-332.
  19. Kowser AR, Amarananjundeswara R, Doddabasappa H, Aravinda Kumar B, Pannure A. Reaction of garlic genotypes to thrips and purple blotch in South India. J. Entomol. Zool. Stud. 2019;7(1):63-66.
  20. Vijaykumar KN, Shripad Kulkarni, Shashidhar TR, Hiremath SM, Vijay Mahajan, Anil Khar, Kambrekar DN, Patil PV. Int. J. Environ. Clim. Change. 2022b;12(12):229-233.

© 2023 Vijaykumar et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:  
<https://www.sdiarticle5.com/review-history/100352>