

Article

Effect of Fertilization on Phenolics of Rapeseeds and Their Antioxidant Potential

Ryszard Amarowicz ^{1,*}, Bożena Cwalina-Ambroziak ², Michał Adam Janiak ¹, Marta Damszel ², Arkadiusz Stępień ³, Katarzyna Sulewska ¹, Magdalena Karamać ¹ and Kamila Penkacik ¹

¹ Institute of Animal Reproduction and Food Research, Polish Academy of Sciences, 10 Tuwima Street, 10-748 Olsztyn, Poland; m.janiak@pan.olsztyn.pl (M.A.J.); k.sulewska@pan.olsztyn.pl (K.S.); m.karamac@pan.olsztyn.pl (M.K.); k.penkacik@pan.olsztyn.pl (K.P.)

² Department of Entomology, Phytopathology and Molecular Diagnostics, Faculty of Agriculture and Forestry, University of Warmia and Masury, 11-041 Olsztyn, Poland; bambr@uwm.edu.pl (B.C.-A.); marta.damszel@uwm.edu.pl (M.D.)

³ Department of Agroecosystems and Horticulture, Faculty of Agriculture and Forestry, University of Warmia and Mazury, 10-721 Olsztyn, Poland; arkadiusz.stepien@uwm.edu.pl

* Correspondence: r.amarowicz@pan.olsztyn.pl

Citation: Amarowicz, R.; Cwalina-Ambroziak, B.; Janiak, M.A.; Damszel, M.; Stępień, A.; Sulewska, K.; Karamać, M.; Penkacik, K. Effect of Fertilization on Phenolics of Rapeseeds and Their Antioxidant Potential. *Foods* **2024**, *13*, 561. <https://doi.org/10.3390/foods13040561>

Academic Editors: Antonello Santini, Antonella Pasqualone and Jaroslawa Rutkowska

Received: 19 December 2023

Revised: 1 February 2024

Accepted: 9 February 2024

Published: 12 February 2024



Copyright: © 2024 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: Three varieties of rapeseed (Castilla, California, and Nelson F1) were cultivated using medium-intensive (control), intensive, and economical (spare) technologies with different nitrogen and sulfur fertilization techniques. The antioxidant potential of rapeseeds was investigated using ABTS, FRAP, and DPPH assays. The content of total phenolic compounds was determined using the Folin–Ciocalteu phenol reagent. The profile of phenolic compounds was determined using high-performance liquid chromatography (HPLC). Diversifying fertilization in various ways influenced the content of phenolic compounds in extracts of rapeseed. In extracts from the Nelson F1 rapeseeds, intensive cultivation resulted in a lower content of phenolic compounds compared to the control group. Economic fertilization reduced the content of phenolic compounds in seeds from the California and Castilla) and six (Nelson F1) main phenolic compounds. Two compounds were identified as sinapine and sinapic acid; others were classified as derivatives of sinapic acid. The effect of fertilization on the antioxidant activity of the seeds and their extracts varied depending on the plant variety and antioxidant assay. For the Castilla and California varieties, no differences were found in the results of the ABTS assay. The antiradical activity against ABTS^{•+} of extracts from the Nelson F1 intensive and spare cultivated seeds was higher than that of extracts from control seeds. The FRAP values of extracts/seeds from the Castilla variety cultivated using different methods did not differ significantly. The results of the DPPH assay were not affected by fertilization in the case of extracts from the California and Castilla varieties. However, the extracts from spare cultivated seeds of Nelson F1 exhibited stronger antiradical activity against DPPH[•]. These findings highlight the complex relationship between fertilization practices, phenolic compound accumulation, and antioxidant activity in rapeseed. Integrating varietal traits and cultivation practices is crucial for optimizing the nutritional benefits of rapeseed.

Keywords: rapeseed; fertilization; phenolic compounds; antioxidant activity; HPLC

1. Introduction

Phenolic compounds are widely distributed in the plant world. They differ in their chemical structure, which in turn affects their chemical properties and biological activity. They appear in all parts of plants, usually accumulated in cell vacuoles, paratomatal, epidermal, subepidermal leaf cells, shoots, and hairs [1], and take part in many plant life activities such as morphogenesis, sex determination, photosynthesis, respiration,

regulation of gene expression, and regulation of the synthesis of growth hormones. One of the defense mechanisms of plants is the production of phenolics, which protect against herbivore attacks but also against pathogens and many abiotic stress factors. Many insects are sensitive to the effects of phenolic acids, flavonoids, tannins, and lignans [2,3]. Phenolics exist in various forms, such as soluble-free, soluble-bound, including esterified, etherified, or glycosylated, and insoluble-bound forms, based on their association with food matrices in plants [4].

The diverse biological properties of phenolics have resulted in a sharp increase in interest in them as bioactive food ingredients, which, as nutraceuticals, can contribute to the therapy of various disorders such as neurological diseases and cancer. Most phenolic compounds exhibit strong antioxidant properties. The antioxidant activity of phenolic compounds depends on their chemical structure and molecular weight and may occur through various mechanisms of action, including electron or hydrogen atom transfer, chelation of the pro-oxidant metal ions, and inhibition of enzymes that catalyze oxidation reactions [5].

Phenolic compounds with the ability to scavenge free radicals may participate in the non-enzymatic antioxidative protection system of the human body. Free radicals can be formed between others under the influence of xenobiotics, UV radiation, and ionizing radiation. Mutual interaction of free radicals with cellular biomolecules such as nucleic acids, proteins, lipids, and carbohydrates leads to various structural damages, for example in the DNA strand, which in turn results in mutations. The resulting changes in DNA may be a signal to initiate pathological cell proliferation and the cancer process. Free oxygen species and free radicals can also contribute to the development of many civilization diseases, for example, atherosclerosis, Parkinson's and Alzheimer's diseases, obesity, and chronic inflammation [6].

Rapeseed (*Brassica napus* L.) seeds are a rich source of oil, protein, and phenolic compounds [7–10]. Sinapine, the choline ester of sinapic acid, is the dominant phenolic compound of rapeseed. Sinapic acid in rapeseed also occurs as free phenolic acids or glycosides [11]. In rapeseed, small amounts of other phenolic acids, such as ferulic acid, caffeic acid, *p*-coumaric acid, vanillic acid, syringic acid, *p*-hydroxybenzoic acid, gentisic acid, and protocatechuic acid, have been reported [9]. The major phenolic compound of rapeseed oil is canolol [12], which is produced by the decarboxylation of sinapic acid during rapeseed thermal processing [13]. The presence of condensed tannins in rapeseed and canola hulls was reported by Naczek et al. [14] and Amarowicz et al. [15].

The antioxidant activity of phenolic compounds from rapeseed or canola was reported by several authors and determined through different methods showing free radical scavenging activity and reducing abilities of these compounds [15–20]. Extracts of phenolic compounds obtained from rapeseed were stronger than those of legumes and cereals [5].

The important accumulation of phenolic compounds in rapeseed results from climatic and soil conditions [21]. Utilizing appropriate production technologies to cultivate plants is an important element in modern, sustainable agriculture, and there are several goals to it, including optimizing mineral fertilization and protecting plants against pests. Nitrogen [22] and sulfur [23] fertilization is very important for many species, not only Brassicaceae plants; however, excess fertilization is harmful and impacts the cost of growing crops [24]. Together with the smaller yields of *Brassica rapa* L. chinensis, the ecological system showed a significantly higher level of phenolic compounds in plants compared to plants cultivated using the conventional system. According to Zhao et al. [25], this may result from the unavailability of nutrients.

The effect of fertilization on the plant phenolic compounds is not easily predictable, and the explanation of the different behaviors is not univocal or understandable on the basis of a biochemical pathway. The experiments of the effect of fertilization on the content of phenolic compounds in fruits (apple, black current, blueberry, grapes, marionberry, orange, peach, strawberry, yellow plum) and vegetables (cabbage, carrot, chicory,

eggplant, lettuce, onion, pepper, potato, red beat, spinach, tomato) were finished with conclusions of “No differences between organic and conventional production”, “Higher in biodynamic than in conventional production”, and “Higher in organic than in conventional production”. The carbon/nitrogen balance (CNB) pathways can be taken into account when studying the effect of nitrogen fertilization on the content and profile of plant phenolic compounds [26–28].

This research aimed to determine the effect of agro-technology (different fertilization) on the antioxidant potential of rapeseeds and their phenolic profile. There are no publications on this important topic in the Scopus and Web of Science databases.

The aim of this research was related to the role of rapeseed as an oil plant and the possibility of using the material after oil extraction as a product to obtain natural oxidants.

2. Materials and Methods

2.1. Plant Material

Winter rape was cultivated in a 4-year monoculture in a close-field experiment in a random block system in triplicate. The experiment was performed in the Agricultural Production Plant in Bałcyny near Olsztyn in Poland (53°78' N, 20°48' E) on soil of a granulometric composition of loamy sand with a high content of phosphorus and an average amount of potassium and magnesium.

The following varieties of winter oilseed rape were included: two population varieties, California (tolerant of fungal diseases, with very good winter hardiness, moderately early in ripening) and Castilla (tolerant of fungal diseases, with very good winter hardiness, with even maturation of siliques), and the Nelson F1 hybrid variety (moderately resistant to fungal diseases, good winter hardiness, medium-early in ripening). Based on previous research [29,30], the following cultivation technologies were applied: medium-intensive (control), intensive, and economical (spare), differing in nitrogen and sulfur fertilization. The agrotechnical conditions of the experiment are shown in Table 1. Seeds were collected at the technical maturity stage.

Table 1. Schema of fertilization (kg/ha) during the field experiment.

| Fertilization | Experimental Group | | |
|---------------------|--------------------|-----------|-------|
| | Control | Intensive | Spare |
| Phosphorus: Autumn | 60 | 80 | 40 |
| Potassium: Autumn | 120 | 150 | 60 |
| Nitrogen: Autumn | 30 | 30 | 30 |
| Nitrogen: Spring I | 120 | 120 | 120 |
| Nitrogen: Spring II | 60 | 80 | 40 |
| Sulphur: Spring | 45 | 60 | - |

2.2. Chemicals

All solvents used were of analytical grade. Methanol, *n*-hexane, ferrous chloride, sinapic acid, sinapine, the Folin–Ciocalteu phenol reagent, 2,2-diphenyl-1-picrylhydrazyl (DPPH) radical, 2,2'-azinobis-(3-ethylbenzothiazoline-6-sulphonic acid) (ABTS), 2,4,6-tri(2-pyridyl)-*s*-triazine (TPTZ), and 6-hydroxy-2,5,7,8-tetramethyl-chroman -2-carboxylic acid (Trolox) were obtained from Sigma (St. Louis, MO, USA).

2.3. Extraction Procedure

Seeds were ground and defatted using a Soxhlet extractor and *n*-hexane as a solvent. Phenolic compounds present in defatted plant material were extracted with a methanol and water mixture (8:2; *v/v*) for 30 min at a temperature of 50 °C. The ratio of plant material to solvent was 1:10 (*w/v*) [31]. Extraction was carried out in a shaking water bath (Elpan 357, Wrocław, Poland). The extracts were filtrated, and the solid residue was extracted

twice more. For the evaporation of methanol from the combined filtrates, a Büchi R-200 rotary evaporator (Büchi, Flawil, Switzerland) was used. The water residue was removed using a freeze dryer.

2.4. Determination of Total Phenolics

The total phenolic content in the rapeseed extracts was estimated using a colorimetric method with the Folin–Ciocalteu phenol reagent [32]. Briefly, a 0.25 mL aliquot of extract dissolved in methanol was added to a test tube containing 4 mL of distilled water. Then, 0.25 mL of the Folin–Ciocalteu phenol reagent and 0.25 mL of a saturated Na_2CO_3 solution were added. After 30 min, the absorbance was recorded at 725 nm using a Beckman DU 7500 diode array spectrophotometer (Beckman Coulter, Inc., Brea, CA, USA), and sinapic acid was used to prepare the standard curve. The results were expressed as mg of sinapic acid equivalents (SAE) per g of extract or g of seed fresh matter (FM).

2.5. Trolox Equivalent Antioxidant Capacity Determination

The ABTS assay described by Re et al. [33] was used to measure the Trolox equivalent antioxidant capacity (TEAC). The solution of $\text{ABTS}^{•+}$ was prepared by mixing an ABTS solution in water with 2.45 mM sodium persulfate. After shaking for 12 h at room temperature, the obtained $\text{ABTS}^{•+}$ stock solution was diluted with methanol to the desired concentration, giving an absorbance of 0.720 at 734 nm. For the colorimetric determination, the absorbance of the sample containing 2 mL of the $\text{ABTS}^{•+}$ solution and 20 μL of rapeseed extract was read after 10 min at 734 nm. The calibration curve was plotted using the Trolox standard. The results were expressed as mmol of Trolox equivalents (TE) per g of extract or g of seed FM.

2.6. Ferric-Reducing Antioxidant Power Determination

A ferric-reducing antioxidant power (FRAP) assay was carried out according to the procedures of Benzie and Strain [34] and Amobonye et al. [35]. For the preparation of the working FRAP reagent, 300 mM acetate buffer (pH 3.6) was mixed with 10 mM TPTZ (in 40 mM HCl) and 20 mM $\text{FeCl}_3 \times 6 \text{H}_2\text{O}$ in a ratio of 10:1:1 (*v/v/v*). For the colorimetric determination, the absorbance of the sample containing 2.25 mL of the FRAP reagent, 75 μL of methanolic solution of rapeseed extract, and 225 μL of deionized water was read at 593 nm after 30 min of incubation. The FRAP values were calculated using the calibration curve for FeSO_4 . The results were expressed as mmol Fe^{2+} equivalents per g of extract or g of seed FM.

2.7. Determination of DPPH Radical Scavenging Activity

The antiradical activity of the extracts against DPPH^{\bullet} was determined according to the Brand-Williams et al. [36] method, with modifications proposed by Sulewska et al. [37]. A 2 mL of deionized water was mixed with a 0.1 mL methanolic solution containing between 0.47 and 2.35 mg of rapeseed extract. Then, 0.25 mL of 1 mM DPPH^{\bullet} solution and 2 mL of deionized water were added. After 20 min, the absorbance of the solution was read at 517 nm. The curves of absorbance vs. extract concentration in the reaction mixture were plotted. The results were expressed as the half maximal scavenging concentration (SC_{50})—the concentration of antioxidants that scavenges half of the radicals.

2.8. HPLC Analysis

Rapeseed extract methanolic solution (5 mg/mL) was filtered (0.45 μm cellulose acetate filter; Millipore, Burlington, MA, USA), and phenolic compounds were analyzed using a Shimadzu HPLC system (Shimadzu Corp., Kyoto, Japan). This system consisted of two LC-10AD pumps, an SPD-M 10A photodiode array detector, and an SCL 10A system controller. Phenolic compounds were separated on a prepacked Luna C_{18} column (4 \times 250 mm, 5 μm ; Phenomenex, Torrance, CA, USA) for 50 min in a gradient system of 5–40%

(*v/v*) acetonitrile in water adjusted to pH 2.5 with trifluoroacetic acid, at a flow rate of 1 mL/min [38]. The separation was monitored at 320 nm, and the injection volume was 20 μ L. The content of sinapine and sinapic acid was calculated using the external standard method. The content of other phenolic compounds showing UV-DAD spectra typical for sinapic acid was expressed as equivalents of this acid.

2.9. Statistical Analysis

In this study, three samples of each rapeseed variety from each field variant were analyzed. Moreover, chemical determinations were triplicated. Results were reported as the mean and standard deviation. For evaluation of the significance of differences among means, analyses of variance (ANOVA) and the Duncan's post hoc test were performed at a level of $p < 0.05$ (GraphPad Prism version 6.04; GraphPad Software, San Diego, CA, USA). The principal component analysis (PCA) and Pearson correlation analysis were carried out using the Statistica 13.1 software (StatSoft Corp., Kraków, Poland).

3. Results

The content of total phenolic compounds in the rapeseed extracts ranged from 48.3 mg SAE/g extract (Castilla, intensive) to 60.9 mg SAE/g extract (California, control) (Table 2). The results expressed in relation to seeds ranged from 4.47 mg SAE/g seed FM (Castillo, spare) to 6.04 mg/g seed FM (Nelson F1, control). The results obtained for the three rapeseed varieties (control) varied as follows: California > Nelson F1 > Castilla for extracts, and Nelson F1 > California > Castilla for seed FM. Diversified fertilization in various ways influenced the total phenolic content of extracts/seeds from different cultivars. For extracts from the Nelson F1 rapeseeds, intensive fertilization resulted in a lower total phenolic content compared to the control group. Economic fertilization reduced the content of total phenolic compounds in extracts of the California variety. The total phenolic content in the California seeds was higher for the intensive group than for the spare group (Table 2).

Table 2. Effect of fertilization on the content of total phenolics and antioxidant potential determined as Trolox equivalent antioxidant capacity (TEAC) and ferric-reducing antioxidant potential (FRAP) of rapeseed fresh matter (FM) and extracts.

| Assay | Unit | Cultivar | Experimental Group | | |
|-----------------|----------------------------------|------------|--------------------|-----------------|-----------------|
| | | | Control | Intensive | Spare |
| Total Phenolics | mg SAE/g extract | California | 60.9 ± 2.6 ab | 59.5 ± 0.7 a | 57.1 ± 0.8 b |
| | | Castilla | 48.9 ± 1.9 ab | 48.3 ± 0.4 a | 49.4 ± 0.5 b |
| | | Nelson F1 | 53.9 ± 0.4 a | 49.8 ± 1.2 b | 53.2 ± 0.3 a |
| | mg SAE/g seed FM | California | 5.21 ± 0.22 a | 5.16 ± 0.06 a | 4.60 ± 0.06 b |
| | | Castilla | 4.49 ± 0.17 a | 4.52 ± 0.12 a | 4.47 ± 0.05 a |
| | | Nelson F1 | 6.04 ± 0.04 a | 5.18 ± 0.12 b | 5.26 ± 0.03 b |
| TEAC | mmol TE/g extract | California | 0.468 ± 0.039 a | 0.417 ± 0.035 a | 0.469 ± 0.035 a |
| | | Castilla | 0.366 ± 0.033 a | 0.360 ± 0.033 a | 0.365 ± 0.032 a |
| | | Nelson F1 | 0.411 ± 0.015 a | 0.418 ± 0.020 a | 0.432 ± 0.015 a |
| | mmol TE/g seed FM | California | 0.040 ± 0.003 a | 0.036 ± 0.003 a | 0.038 ± 0.003 a |
| | | Castilla | 0.034 ± 0.003 a | 0.034 ± 0.003 a | 0.033 ± 0.003 a |
| | | Nelson F1 | 0.033 ± 0.003 b | 0.043 ± 0.002 a | 0.043 ± 0.002 a |
| FRAP | mmol Fe ²⁺ /g extract | California | 1.61 ± 0.03 a | 1.66 ± 0.05 a | 1.43 ± 0.05 b |
| | | Castilla | 1.34 ± 0.0 a | 1.30 ± 0.06 a | 1.30 ± 0.08 a |
| | | Nelson F1 | 1.35 ± 0.11 a | 1.15 ± 0.05 b | 1.31 ± 0.04 a |
| | mmol Fe ²⁺ /g seed FM | California | 0.138 ± 0.003 a | 0.144 ± 0.004 a | 0.115 ± 0.004 b |
| | | Castilla | 0.123 ± 0.007 a | 0.122 ± 0.006 a | 0.122 ± 0.007 a |
| | | Nelson F1 | 0.151 ± 0.012 a | 0.120 ± 0.005 b | 0.129 ± 0.004 b |

Means with different letters in the same row are significantly different ($p < 0.05$). SAE—sinapic acid equivalent; TE—Trolox equivalent.

The antioxidant activity determined using ABTS and FRAP assays also depended on the rapeseed variety (Table 2). The Castilla samples showed lower TEAC compared to the seeds and extracts of the other varieties. FRAP values were higher for the California extracts. When considering the effect of fertilization on TEAC, no differences were found between extracts for individual varieties and between seeds of different groups for the California and Castilla varieties. In the case of the Nelson F1 variety, the TEAC of seeds from the intensive and spare groups (0.043 mmol TE/g seed FM) was higher than that of seeds from the control group (0.033 mmol TE/g seed FM). The obtained FRAP values for extracts of the Castilla variety did not differ from each other. In the California variant, the FRAP values for the spare group (1.43 Fe²⁺/g extract; 0.115 mmol Fe²⁺/g seed FM) were lower than the FRAP values for the control group (1.61 mmol Fe²⁺/g extract; 0.138 mmol Fe²⁺/g seed FM) and the intensive group (1.66 mmol Fe²⁺/g extract; 0.144 mmol Fe²⁺/g seed FM).

The results of the DPPH assay are depicted in Figures 1 and 2. For all extracts, dose-related antiradical activity against DPPH• was observed. Comparing SC₅₀ values, the effect of fertilization was not observed for extracts of the California and Castilla varieties. In turn, the extracts of the seeds from the Nelson F1 spare group exhibited stronger antiradical activity against DPPH• than the extracts from the intensive and control groups.




Figure 1. Antiradical activity of rapeseed extracts against DPPH radicals.




Figure 2. Antiradical activity of rapeseed extracts against DPPH radicals expressed as SC_{50} . Means with different letters are significantly different ($p < 0.05$).

The HPLC chromatograms of the extracts were characterized by the presence of five (California and Castilla) and six (Nelson F1) main phenolic compounds (Figure 3). Applying the original standards, compounds 1 and 4 were identified as sinapine and sinapic acid, respectively. Based on the DAD-UV spectra, it was tentatively found that compounds 2, 3, 5, and 6 were derivatives of sinapic acid.

The content of sinapine in the rapeseed extracts ranged from 64.5 mg/g extract (Nelson F1 intensive group) to 81.9 mg/g extract (California intensive group) (Table 3). The lowest content of sinapic acid was determined in the extracts from the Nelson F1 control group—3.8 mg/g extract, and the highest was in the extracts from the California control group—6.3 mg/g extract. The content of sinapine in seeds ranged from 6.2 mg/g seed FM (California spare group) to 7.4 mg/g seed FM (Nelson F1, control group). The lowest content of sinapic acid was found in the Nelson F1 intensively cultivated seeds—0.41 mg/g seed FM. The control and intensively cultivated seeds of the California and Castillo seeds in the control group were characterized by the highest content of sinapic acid—0.54 mg/g seed FM. Interestingly, compound 5 was quantified only in the seeds/extracts of the Nelson F1 cultivar. The content of this compound was relatively high in the range of 16.9–18.3 mg/g extract and 1.69–2.05 mg/g seed FM. Unlike sinapine and sinapic acid, for which no effect of fertilization was found, its content in control seeds was significantly higher than in intensively cultivated and spare seeds.




Figure 3. HPLC chromatograms of rapeseed extracts. The peak numbers correspond to the compounds listed in Table 3.

Table 3. Effect of fertilization on the content of individual phenolic compounds in rapeseed extracts and seeds.

| Compound | Unit | Cultivar | Experimental Group | | |
|--------------------------------|--------------|------------|--------------------|---------------|---------------|
| | | | Control | Intensive | Spare |
| 1 (Sinapine) | mg/g extract | California | 80.1 ± 1.4 a | 81.9 ± 5.1 a | 76.2 ± 4.0 a |
| | | Castilla | 76.3 ± 3.4 a | 73.0 ± 5.6 a | 70.1 ± 1.4 a |
| | | Nelson F1 | 65.8 ± 3.0 a | 64.5 ± 3.3 a | 70.8 ± 5.3 a |
| | mg/g seed FM | California | 6.8 ± 0.1 a | 7.0 ± 0.4 a | 6.2 ± 0.3 a |
| | | Castilla | 7.0 ± 0.3 a | 6.8 ± 0.5 a | 6.3 ± 0.1 a |
| | | Nelson F1 | 7.4 ± 0.3 a | 6.7 ± 0.3 a | 7.0 ± 0.5 a |
| 2 (Sinapic acid derivative) | mg/g extract | California | 8.4 ± 1.4 a | 7.7 ± 0.6 a | 7.6 ± 0.6 a |
| | | Castilla | 7.1 ± 0.9 a | 7.1 ± 1.2 a | 6.4 ± 0.2 a |
| | | Nelson F1 | 4.0 ± 0.1 a | 2.5 ± 0.2 b | 3.1 ± 0.8 ab |
| | mg/g seed FM | California | 0.72 ± 0.03 a | 0.67 ± 0.05 a | 0.61 ± 0.05 a |
| | | Castilla | 0.65 ± 0.08 a | 0.66 ± 0.11 a | 0.58 ± 0.02 a |
| | | Nelson F1 | 0.45 ± 0.01 a | 0.26 ± 0.02 b | 0.31 ± 0.08 b |
| 3 (Sinapic acid derivative) | mg/g extract | California | 5.6 ± 0.3 a | 5.1 ± 0.2 a | 5.1 ± 0.3 a |
| | | Castilla | 5.9 ± 0.5 a | 5.9 ± 1.0 a | 5.4 ± 0.5 a |
| | | Nelson F1 | 5.2 ± 0.1 a | 4.9 ± 0.2 a | 5.0 ± 0.6 a |
| | mg/g seed FM | California | 0.48 ± 0.03 a | 0.44 ± 0.02 a | 0.41 ± 0.02 a |
| | | Castilla | 0.54 ± 0.06 a | 0.49 ± 0.06 a | 0.50 ± 0.01 a |
| | | Nelson F1 | 0.58 ± 0.01 a | 0.51 ± 0.02 b | 0.50 ± 0.04 b |
| 4 (Sinapic acid) | mg/g extract | California | 6.3 ± 0.4 a | 6.2 ± 0.2 a | 6.1 ± 0.3 a |
| | | Castilla | 5.9 ± 0.7 a | 5.2 ± 0.6 a | 5.5 ± 0.1 a |
| | | Nelson F1 | 3.8 ± 0.2 a | 3.9 ± 0.5 a | 4.4 ± 0.8 a |
| | mg/g seed FM | California | 0.54 ± 0.04 a | 0.54 ± 0.03 a | 0.49 ± 0.02 a |
| | | Castilla | 0.54 ± 0.06 a | 0.49 ± 0.06 a | 0.50 ± 0.01 a |
| | | Nelson F1 | 0.43 ± 0.02 a | 0.41 ± 0.05 a | 0.44 ± 0.08 a |
| 5 (Sinapic acid derivative) | mg/g extract | California | - | - | - |
| | | Castilla | - | - | - |
| | | Nelson F1 | 18.3 ± 0.5 a | 16.9 ± 1.0 a | 17.1 ± 0.9 a |
| | mg/g seed FM | California | - | - | - |
| | | Castilla | - | - | - |
| | | Nelson F1 | 2.05 ± 0.06 a | 1.76 ± 0.01 b | 1.69 ± 0.09 b |
| 6 (Sinapic acid derivative) | mg/g extract | California | 8.5 ± 0.5 a | 8.3 ± 0.2 a | 7.6 ± 0.5 a |
| | | Castilla | 6.0 ± 0.7 a | 5.7 ± 0.2 a | 6.4 ± 1.5 a |
| | | Nelson F1 | 3.8 ± 0.4 a | 3.2 ± 0.4 a | 3.5 ± 0.8 a |
| | mg/g seed FM | California | 0.73 ± 0.04 a | 0.72 ± 0.02 a | 0.61 ± 0.04 a |
| | | Castilla | 0.55 ± 0.05 a | 0.53 ± 0.05 a | 0.58 ± 0.05 a |
| | | Nelson F1 | 0.43 ± 0.04 a | 0.33 ± 0.04 b | 0.35 ± 0.08 b |

Means with different letters in the same row are significantly different ($p < 0.05$). FM—fresh matter.

The collected data from the analyses of the total phenolic content and antioxidant assays were subjected to PCA to identify the relationships between these variables and rapeseed varieties cultivated in different ways. The PCA results computed for the values of variables expressed per extract are depicted in Figure 4 and for those expressed per seed FM in Figure 5. In the case of data relating to extracts, the first two components (PC1 and PC2) explained 92.45% of the total variance. The system was divided along PC1 into two groups. The first was formed by extracts from the California variety seeds cultivated in each method (A1–A3) and extracts from the Nelson F1 spare cultivated seeds (C3). Most of the variables (total phenolic content and results of ABTS and FRAP assays) described this group. The second group was formed by the rest of the samples, which were closely associated with the results of the DPPH assay. For data relating to seed FM, PC1 and PC2 significantly explained 96.66% of the total variance. Variables affected the A1 and A2, as

well as seeds of the Nelson F1 variety from the control (C1) and spare (C3) cultivations. To determine direct correlations between variables, a Pearson correlation analysis was performed for data relating to the extracts. Total phenolic content significantly ($p < 0.05$) correlated with the results of the ABTS and FRAP assays (Table 4). A high correlation coefficient (0.672; $p < 0.05$) was also found for the correlation between TEAC and FRAP. These findings are in line with those shown by PCA (Figure 4B).




Figure 4. Plots of principal component analysis (PCA) of the objects (A) and variables (B) for data relating to extracts. A1—California control group; A2—California intensive group; A3—California spare group; B1—Castilla control group; B2—Castilla intensive group; B3—Castilla spare group; C1—Nelson F1 control group; C2—Nelson F1 intensive group; C3—Nelson F1 spare group; TEAC—Trolox equivalent antioxidant capacity; FRAP—ferric-reducing antioxidant potential; DPPH—DPPH radical scavenging activity.




Figure 5. Plots of principal component analysis (PCA) of the objects (A) and variables (B) for data relating to seed FM. A1—California control group; A2—California intensive group; A3—California spare group; B1—Castilla control group; B2—Castilla intensive group; B3—Castilla spare group; C1—Nelson F1 control group; C2—Nelson F1 intensive group; C3—Nelson F1 spare group; TEAC—Trolox equivalent antioxidant capacity; FRAP—ferric-reducing antioxidant potential.

Table 4. Pearson correlations between total phenolic content, Trolox equivalent antioxidant capacity (TEAC), ferric-reducing antioxidant potential (FRAP), and DPPH radical scavenging activity of rapeseed of different varieties cultivated in different conditions.

| Variables | Regression Equation | Correlation Coefficient |
|--------------------------|-------------------------|-------------------------|
| Total phenolics vs. TEAC | $y = 0.0071x + 0.0327$ | 0.814 * |
| Total phenolics vs. FRAP | $y = 0.0293x + 0.5733$ | 0.871 * |
| Total phenolics vs. DPPH | $y = -0.0027x + 0.4418$ | -0.271 |
| TEAC vs. FRAP | $y = 1.740x + 0.667$ | 0.672 * |
| TEAC vs. DPPH | $y = -0.2182x + 0.3864$ | -0.190 |
| FRAP vs. DPPH | $y = -0.076x + 0.4016$ | -0.255 |

* significant at $p < 0.05$.

4. Discussion

The results of total phenolic and sinapine content in rapeseed extracts and seeds are comparable with those reported in the literature. Thiyam et al. [20] reported that the content of total phenolics ranged from 12 to 24 mg/g of rapeseed oil-free meal. According to Mert-Türk et al. [39], the sinapine content in five rapeseed cultivars ranged from 0.33 to 0.43 g/kg. Similar results were reported for sinapic acid content in rapeseed by Siger et al. [40], who found a significant variation in the content of this phenolic compound in rapeseed between the individual harvest years within the same varieties. The variety was a very important factor that influenced the content of phenolic compounds in broccoli [41,42].

Sinapine and sinapic acid are the well-known phenolic compounds of rapeseed. In our research, we also determined four other rapeseed phenolic compounds that were tentatively identified as sinapic acid derivatives. Identification was made based on the similarity of the shape of the UV-DAD spectra with a maxima at a wavelength of 328–330 nm of these compounds with that of sinapic acid. The presence of sinapic acid derivatives in rapeseed has been reported in the literature [43]. Among them, more polar compounds than sinapic acid, including sinaopyl glucose and sinaopyl malate, and less polar compounds, such as disinapic acid and its derivatives, were found. In our study, two sinapic acid derivatives (compounds 2 and 3) were also more polar, and two others (compounds 5 and 6) were eluted after sinapic acid. However, further research on determining the structure of these compounds using more advanced analytical techniques and their full identification is necessary. Ferulic acid and its derivatives, quercetin-sinapoyl di-hexoseptose, and kaempferol 3-sinapoylsphorotrioside-7-glucoside were another phenolic compound identified previously in rapeseeds [43]. Our study did not confirm the presence of these compounds in the analyzed varieties.

The effect of different fertilization systems used in our study on the content of total and individual phenolic compounds and the antioxidant potential of the rapeseeds and their extracts varied depending on the plant variety and antioxidant assay; however, clear trends of change were not found. The results of scientific research also do not indicate a clear effect of fertilization on the content of phenolic compounds in rapeseed and other plants belonging to the Brassicaceae family and their antioxidant potential. Increased nitrogen fertilization of rapeseed, as claimed by Butkutė et al. [22] and Mert-Türk et al. [39], increased the content of phenolic compounds in seeds. De Pascale et al. [44] demonstrated an increased accumulation of phenolic acids, including gallic acid, in *Brassica rapa* plants under the influence of increased sulfur fertilization. Fertilization of red cabbage with 150 kg N/ha (nitrogen in the form of ammonium nitrate) was most affected in terms of the content of phenolic compounds and the antioxidant capacity [45]. Sady et al. [46] found the highest content of phenolics in *Brassica oleracea* var. *capitata* alba L. fertilized with RSM (solution ammonium nitrate and urea 1:1). In contrast, according to Li et al. [47], a significant decrease in phenolic compounds in mustard leaves was caused by increasing the level of nitrogen fertilization. Broccoli showed a decrease in flavonoid contents with

nitrogen fertilization [48]. The maximum concentration of phenolic compounds was found in broccoli [49] and Chinese cabbage [50] in nonfertilized N plants.

In the ecological system used for *Brassica rapa* L. chinensis plant cultivation, the level of phenolic compounds was higher compared to the conventional system [51]. Johnson et al. [52] reported that a liquid organic nitrogen source should be applied rather than traditional mineral fertilization due to the increased accumulation of *p*-coumaric acid in pac choi. In the organic growing cauliflower *Brassica oleracea* L. subsp. Botrytis, compared to a conventional one, 24, 21, 13, 48, and 44% higher contents of ascorbic acid, polyphenols, carotenoids, and volatile substances were found, as well as a higher antioxidant capacity [53]. Conversa et al. [54] also showed a higher level of bioactive compounds in organic *Brassica rapa* L. The influence of different nitrogen fertilizations at doses of 0, 100, 200, and 300 mg/dm³, as well as the type of soil, did not affect the total phenolic content and FRAP results in the two varieties of kale, *Brassica oleracea* L. var. acephala [55]. However, red kale, in comparison with green-leafed kale, exhibited a positive effect of N fertilization. In addition, in other research, the production system did not show any effect on the total phenolic content and total flavonoid content in cauliflower (except for single genotypes) [56] or broccoli [57]. In experiments by Conversa et al. [54], the modifying effect of conventional and organic growing systems on the antioxidant properties of crude or processed *Brassica rapa* L. subsp. Sylvestris was not observed. According to Ibrahim and Jaafar [58], Zhang et al. [59], and Deng et al. [60], nitrogen deficiency or lower nitrogen content can trigger ethylene signaling and upregulate the transcription factor MYB12 to stimulate genes involved in the biosynthesis of phenylpropanoids and flavonoids, including chalcone synthase and L-phenylalanine ammonia-lyase. The results of a meta-analysis by Sun et al. [61] suggest a reduction in the internal C/N ratio by nitrogen application and a lowering of the biosynthesis of phenolic compounds, such as phenolic acids, flavonoids, including anthocyanins, and condensed tannins. Regulation by N at the branch point of 3-dehydroshikimate is proposed by Salminen and Maarit [28] to explain the variation in the abundance pattern of phenylpropanoids.

The principal component analysis (PCA) was employed in our study to further understand the relationship within the system. For the data relating to extracts, the importance of PC2 was relatively low (15.02 %), and PC1 explained the majority of the variance (77.43 %). This significant component discriminated the samples according to the variety rather than the type of fertilization, although in the case of the Nelson F1 variety, the division into seeds from different cultivations occurred along PC1. Considering the PCA results for the seed data, both PC1 (60.25%) and PC2 (36.41) significantly explained the total variance, and they clearly discriminated the Castilla variety from the remaining varieties. For the California and Nelson F1 seeds, samples were discriminated along PC1 and PC2, but no clear clusters were observed based on cultivation type, so we concluded that the effect of fertilization was individual for each variety.

The data from the PCA and the Pearson correlation analysis showed that the results of total phenolics, ABTS, and FRAP assays correlated with each other. The same relation between similar variables was observed for false flax extracts [38] and extracts from seeds of different *Vitis* species [62].

5. Conclusions

This study delved into the impact of varied fertilization techniques on the phenolic compound profile and antioxidant activity of different rapeseed cultivars. We observed significant differences in phenolic content, with the California variety showing the highest content under control conditions. Intensive fertilization reduced the content of phenolic compounds in the Nelson F1 seeds, while economic fertilization decreased the phenolic content in the California variety. Antioxidant assessments through ABTS and FRAP assays revealed cultivar-specific responses. The California variety exhibited higher antioxidant activity in intensively cultivated seeds, whereas the Nelson F1 variety showed stronger antiradical activity in spare-cultivated seeds. HPLC analysis identified key phenolic

compounds, predominantly sinapine and sinapic acid. These findings highlight the complex relationship between fertilization practices, phenolic compound accumulation, and the antioxidant activity of rapeseed. Integrating varietal traits and cultivation practices is crucial for optimizing the nutritional benefits of rapeseed. Future research should explore the biochemical mechanisms influencing phenolic compound synthesis in response to specific fertilization regimens.

Author Contributions: Conceptualization, B.C.-A., R.A., and M.A.J.; methodology, B.C.-A., M.D., A.S., and M.A.J.; software, M.A.J. and K.S.; validation, M.A.J. and K.S.; formal analysis, K.S., M.A.J., K.P., M.K., A.S., and M.D.; investigation, M.D. and K.S.; data curation, M.D., K.S., and B.C.-A.; writing—original draft preparation, R.A. and B.C.-A.; writing—review and editing, R.A., B.C.-A., and M.K.; visualization, M.A.J.; supervision, R.A.; project administration, R.A. and B.C.-A. All authors have read and agreed to the published version of the manuscript.

Funding: This research received no external funding.

Institutional Review Board Statement: Not applicable.

Informed Consent Statement: Not applicable.

Data Availability Statement: Data is contained within the article.

Conflicts of Interest: The authors declare no conflicts of interest.

References

- Zagoskina, N.V.; Zubova, M.Y.; Nechaeva, T.L.; Kazantseva, V.V.; Goncharuk, E.A.; Katanskaya, V.M.; Baranova, E.N.; Aksenova, M.A. Polyphenols in plants: Structure, biosynthesis, abiotic stress, regulation, and practical applications. *J. Mol. Sci.* **2023**, *24*, 13874. <https://doi.org/10.3390/ijms241813874>.
- Pratyusha, S. Phenolic compounds in the plant development and defense: An overview. In *Plant Stress Physiology—Perspectives in Agriculture*; Hasanuzzaman, M., Nahar, K., Eds.; IntechOpen: London, UK, 2022, pp. 1–17. <https://doi.org/10.5772/intechopen.102873>.
- Singh, S.; Kaur, I.; Kariyat, R. The multifunctional roles of polyphenols in plant-herbivore interactions. *Int. J. Mol. Sci.* **2021**, *22*, 1442. <https://doi.org/10.3390/ijms22031442>.
- Shahidi, F.; Hossain, A. Importance of insoluble-bound phenolics to the antioxidant potential is dictated by source material. *Antioxidants* **2023**, *12*, 203. <https://doi.org/10.3390/antiox12102023>.
- Amarowicz, R.; Pegg, R.B. Natural antioxidants of plant origin. *Adv. Food Nutr. Res.* **2019**, *90*, 1–81. <https://doi.org/10.1016/bs.afnr.2019.02.011>.
- Rasouli, H.; Farzaei, M.H.; Khodarahmi, R. Polyphenols and their benefits: A review. *Int. J. Food Prop.* **2017**, *20*, 1700–1741. <https://doi.org/10.1080/10942912.2017.1354017>.
- Loganes, C.; Ballali, S.; Minto, C. Main properties of canola oil components: A descriptive review of current knowledge. *Open Agric. J.* **2016**, *10*, 69–74. <https://doi.org/10.2174/1874331501610010069>.
- Mupondwa, E.; Li, X.; Wanasundara, J.P.D. Technoeconomic prospects for commercialization of *Brassica* (Cruciferous) plant proteins. *J. Am. Oil Chem. Soc.* **2018**, *95*, 903–922. <https://doi.org/10.1002/aocs.12057>.
- Chmielewska, A.; Kozłowska, M.; Rachwał, D.; Wnukowski, P.; Amarowicz, R.; Nebesny, E.; Rosicka-Kaczmarek, J. Canola/rapeseed protein—nutritional value, functionality and food application: A review. *Crit. Rev. Food Sci. Nutr.* **2020**, *61*, 3836–3856. <https://doi.org/10.1080/10408398.2020.1809342>.
- Naczka, M.; Amarowicz, R.; Sullivan, A.; Shahidi, F. Current research developments on polyphenolics of rapeseed/canola: A review. *Food Chem.* **1998**, *62*, 489–502. [https://doi.org/10.1016/S0308-8146\(97\)00198-2](https://doi.org/10.1016/S0308-8146(97)00198-2).
- Szydłowska-Czeraniak, A. Rapeseed and its products—sources of bioactive compounds: A review of their characteristics and analysis. *Crit. Rev. Food Sci. Nutr.* **2013**, *53*, 307–330. <https://doi.org/10.1080/10408398.2010.529959>.
- Rękas, A.; Ścibisz, I.; Siger, A.; Wroniak, M. The effect of microwave pretreatment of seeds on the stability and degradation kinetics of phenolic compounds in rapeseed oil during long-term storage. *Food Chem.* **2017**, *222*, 43–52. <https://doi.org/10.1016/j.foodchem.2016.12.003>.
- Shrestha, K.; Stevens, C.V.; de Meulenaer, B. Isolation and identification of a potent radical scavenger (canolol) from roasted high erucic mustard seed oil from Nepal and its formation during roasting. *J. Agric. Food Chem.* **2012**, *60*, 7506–7512. <https://doi.org/10.1021/jf301738y>.
- Naczka, M.; Amarowicz, R.; Pink, D.; Shahidi, F. Insoluble condensed tannins of canola/rapeseed. *J. Agric. Food Chem.* **2000**, *48*, 1758–1762. <https://doi.org/10.1021/jf9908401>.
- Amarowicz, R.; Naczka, M.; Shahidi, F. Antioxidant activity of crude tannins of canola and rapeseed hulls. *J. Am. Oil Chem. Soc.* **2000**, *77*, 957–961. <https://doi.org/10.1007/s11746-000-0151-0>.

16. Szydłowska-Czerniak, A.; Tułodziecka, A. Antioxidant capacity of rapeseed extracts obtained by conventional and ultrasound-assisted extraction. *J. Am. Oil. Chem. Soc.* **2014**, *91*, 2011–2019. <https://doi.org/10.1007/s11746-014-2557-4>.
17. Szydłowska-Czerniak, A.; Amarowicz, R.; Szlyk, E. Antioxidant capacity of rapeseed meal and rapeseed oils enriched with meal extract. *Eur. J. Lipid Sci. Technol.* **2010**, *112*, 750–760. <https://doi.org/10.1002/ejlt.200900292>.
18. Amarowicz, R.; Fornal, J.; Karamać, M. Effect of seed moisture on phenolic acids in rapeseed oil cake. *Grasas Y Aceites* **1995**, *46*, 354–356. <https://doi.org/10.3989/gya.1995.v46.i6.952>.
19. Amarowicz, R.; Fornal, J.; Karamać, M.; Shahidi, F. Antioxidant activity of extracts of phenolic compounds from rapeseed oil cakes. *J. Food Lipids* **2001**, *8*, 65–74. <https://doi.org/10.1111/j.1745-4522.2001.tb00184.x>.
20. Thiyam, U.; Stöckmann, H.; Zum Felde, T.; Schwarz, K. Antioxidative effect of the main sinapic acid derivatives from rapeseed and mustard oil by-products. *Eur. J. Lipid Sci. Technol.* **2006**, *108*, 239–248. <https://doi.org/10.1002/ejlt.200500292>.
21. Björkman, M.; Klingen, I.; Birch, A.N.E.; Bones, A.M.; Bruce, T.J.B.; Johansen, T.J.; Meadow, R.; Mølmann, J.; Seljåsen, R.; Smart, L.E.; et al. Phytochemicals of Brassicaceae in plant protection and human health—Influences of climate, environment and agronomic practice. *Phytochemistry* **2011**, *72*, 538–556. <https://doi.org/10.1016/j.phytochem.2011.01.014>.
22. Butkutė, B.; Mašauskienė, A.; Šidlauskas, G.; Šliesaravičienė, L. The effect of agronomic factors and growth conditions on protein and fat content in the seed of spring oilseed rape and on the variation of fatty acids. *Agriculture* **2000**, *70*, 160–175.
23. Zhao, F.M.; McGrath, S.P.; Blake-Kalff, M.M.A.; Link, A.; Tucker, M. Crop responses to sulphur fertilization in Europe. Biogeochemistry of sulphur in agricultural systems. Part II. Fertilizers and fertilization. *Proc. Int. Fertl. Soc.* **2003**, *16*, 26–51.
24. Podleśna, A.; Cacak-Pietrzak, G.; Sowiński, M. Response of winter wheat to fertilizing S in the field experiment. *Fert. Fertil.* **2003**, *4*, 169–179.
25. Zhao, X.; Nechols, J.R.; Williams, K.; Wang, W.; Carey, E.E. Comparison of phenolic acids in organically and conventionally grown pac choy (*Brassica rapa* L./chinensis). *J. Sci. Food. Agric.* **2009**, *89*, 940–946. <https://doi.org/10.1002/jsfa.3534>.
26. Romani, A.; Ieri, F. Plant polyphenol content, soil fertilization and agricultural management: A review. *Eur. Food Res. Technol.* **2017**, *243*, 1107–1115. <https://doi.org/10.1007/s00217-016-2826-6>.
27. Nguyen, P.M.; Niemeyer, E.D. Effects of nitrogen fertilization on the phenolic composition and antioxidant properties of basil (*Ocimum basilicum* L.). *J. Agric. Food Chem.* **2008**, *56*, 8685–8691. <https://doi.org/10.1021/jf801485u>.
28. Salminen, J.P.; Maarit, K. Chemical ecology of tannins and other phenolics: We need a change in approach. *Func. Ecol.* **2011**, *25*, 325–338. <https://doi.org/10.1111/j.1365-2435.2010.01826.x>.
29. Cwalina-Ambroziak, B.; Stępień, A.; Kurowski, T.P.; Głosek-Sobieraj, M.; Wiktorski, A. The health status and yield of winter rapeseed (*Brassica napus* L.) grown in monoculture and in crop rotation under different agricultural production systems. *Arch. Agron. Soil Sci.* **2016**, *62*, 1722–1732. <https://doi.org/10.1080/03650340.2016.1171851>.
30. Stępień, A.; Wojtkowiak, K.; Pietrzak-Fiecko, R. Nutrient content, fat yield and fatty acid profile of winter rapeseed (*Brassica napus* L.) grown under different agricultural production systems. *Chilean J. Agric. Res.* **2017**, *77*, 266–272. <https://doi.org/10.4067/S0718-58392017000300266>.
31. Amarowicz, R.; Wanasundara, U.N.; Karamać, M.; Shahidi, F. Antioxidant activity of ethanolic extract of mustard seed. *Nahrung—Food* **1996**, *40*, 261–263. <https://doi.org/10.1002/food.19960400506>.
32. Naczka, M.; Shahidi, F. The effect of methanol-ammonia-water treatment on the content of phenolic acids of canola. *Food Chem.* **1989**, *31*, 159–164. [https://doi.org/10.1016/0308-8146\(89\)90026-5](https://doi.org/10.1016/0308-8146(89)90026-5).
33. Re, R.; Pellegrini, N.; Proteggente, A.; Pannala, A.; Yang, M.; Rice-Evans, C. Antioxidant activity applying an improved ABTS radical cation decolorization assay. *Free Rad. Biol. Med.* **1999**, *26*, 1231–1237. [https://doi.org/10.1016/S0891-5849\(98\)00315-3](https://doi.org/10.1016/S0891-5849(98)00315-3).
34. Benzie, I.F.; Strain, J.J. The ferric reducing ability of plasma (FRAP) as a measure of "antioxidant power": The FRAP assay. *Anal Biochem* **1999**, *239*, 70–76. <https://doi.org/10.1006/abio.1996.0292>.
35. Amobonye, A.; Bhagwat, P.; Ruzengwe, F.M.; Singh, S.; Pillai, S. Pear juice clarification using polygalacturonase from *Beauveria bassiana*: Effects on rheological, antioxidant and quality properties. *Pol. J. Food Nutr. Sci.* **2022**, *72*, 57–67. <https://doi.org/10.31883/pjfn/145704>.
36. Brand-Williams, W.; Cuvelier, M.E.; Berset, C. Use of a free radical method to evaluate antioxidant activity. *LWT Food Sci. Technol.* **1995**, *28*, 25–30. [https://doi.org/10.1016/S0023-6438\(95\)80008-5](https://doi.org/10.1016/S0023-6438(95)80008-5).
37. Sulewska, K.; Rybarczyk-Płońska, A.; Karamać, M. Antioxidant capacity of lentil flour hydrolysates obtained with pancreatin. *Pol. J. Food Nutr. Sci.* **2022**, *72*, 381–391. <https://doi.org/10.31883/pjfn/155932>.
38. Karamać, M.; Gai, F.; Peiretti, P.G. Effect of the growth stage of false flax (*Camelina sativa* L.) on the phenolic compound content and antioxidant potential of the aerial part of the plant. *Pol. J. Food Nutr. Sci.* **2020**, *70*, 189–198. <https://doi.org/10.31883/pjfn/119719>.
39. Mert-Türk, F.; Kemal, Ü.L.; Egesel, Ö. Nitrogen and fungicide applications against *Erysiphe cruciferarum* affect quality components of oilseed rape. *Mycopathologia* **2008**, *165*, 27–35. <https://doi.org/10.1007/s11046-007-9070-3>.
40. Siger, A.; M. Nogala-Kałucka, M.; E. Lampart-Szczapa, E.; Hoffmann, A.A. Phenolic compound contents in new rape varieties. *Oilseed Crops* **2004**, *25*, 263–274.
41. Eberhardt, M.V.; Kobira, K.; Keck, A.S.; Juvik, J.A.; Jeffery, E.H. Correlation analyses of phytochemical composition, chemical, and cellular measures of antioxidant activity of broccoli (*Brassica oleracea* L. var. *italica*). *J. Agric. Food Chem.* **2005**, *53*, 7421–7431. <https://doi.org/10.1021/jf051495k>.
42. Singh, A.; Upadhyay, K.; Prasad, A.; Bahadur, A.; Rai, M. Variability of carotenes, vitamin C, E and phenolics in brassica vegetables. *J. Food Comp. Anal.* **2007**, *20*, 106–112. <https://doi.org/10.1016/j.jfca.2006.08.002>.

43. Reungoat, V.; Allais, F.; Ducatel, H.; Ioannou, I. Extraction and purification processes of sinapic acid derivatives from rapeseed and mustard seed by-products. *Separ. Purific. Rev.* **2022**, *51*, 521–544. <https://doi.org/10.1080/15422119.2021.2004550>.
44. De Pascale, S.; Maggio, A.; Pernice, R.; Fogliano, V.; Barbieri, G. Sulphur fertilization may improve the nutritional value of *Brassica rapa* L. subsp. *sylvestris*. *Eur. J. Agron.* **2007**, *26*, 418–424. <https://doi.org/10.1016/j.eja.2006.12.009>.
45. Biesiada, A.; Nawirska-Olszańska, A.; Kucharska, A.; Sokół-Łętowska, K.; Kędra. The effect of nitrogen fertilization on nutritive value and antioxidative activity of red cabbage. *Acta Sci Pol. Hortorum Cultus* **2010**, *9*, 13–21.
46. Sady, W.; Domagała-Świątek, I.; Rożek, S. Effect of nitrogen fertilization on sugars, ascorbic acid and phenolic compounds in white cabbage (*Brassica oleracea* var. *capitata alba* L.). *Acta Sci. Pol. Hortorum Cultus* **2010**, *9*, 41–51.
47. Li, J.; Zhu, Z.; Gerendas, J. Effects of nitrogen and sulfur on total phenolics and antioxidant activity in two genotypes of leaf mustard. *J. Plant Nutr.* **2008**, *31*, 1642–1655. <https://doi.org/10.1080/01904160802244860>.
48. Fortier, E.; Desjardins, Y.; Tremblay, N.; Bělec, C.; Côté, M. Influence of irrigation and nitrogen fertilization on broccoli polyphenolics concentration. *Acta Hort.* **2010**, *856*, 55–62. <https://doi.org/10.17660/ActaHortic.2010.856.6>.
49. Jones, R.B.; Imsic, M.; Franz, P.; Tomkins, B.T. High nitrogen during growth reduced glucoraphanin and flavonol content in broccoli (*Brassica oleracea* var. *italica*) heads. *Aust. J. Exp. Agric.* **2007**, *47*, 1498–1505. <https://doi.org/10.1071/EA06205>.
50. Zhu, W.; Lin, X.; Jin, C.; Zhang, Y.; Fang, P. Effects of nitrogen application rates on antioxidant contents and antioxidative activities in Chinese cabbage (*Brassica chinensis* L.). *J. Zhejiang Univ. Agric. Life Sci.* **2009**, *35*, 299–306.
51. Zhao, X.; Carey, E.E.; Young, J.E.; Wang, W.; Iwamoto, T. Influences of organic fertilization, high tunnel environment, and post-harvest storage on phenolic compounds in lettuce. *HortScience* **2007**, *42*, 71–76. <https://doi.org/10.21273/HORTSCI.42.1.71>.
52. Johnson, W.A.; Cloyd, R.A.; Nechols, J.R.; Williams, K.A.; Nelson, N.O.; Rotenberg, D.; Kennelly, M.M. Effect of nitrogen source on pac choy (*Brassica rapa* L.) chemistry and interactions with the diamondback moth (*Plutella xylostella* L.). *HortiScience* **2012**, *47*, 1457–1465. <https://doi.org/10.21273/HORTSCI.47.10.1457>.
53. Lo Scalzo, R.; Picchi, V.; Migliori, C.A.; Campanelli, G.; Leteo, F.; Ferrari, V.; Di Cesare, L.F. Variations in the phytochemical contents and antioxidant capacity of organically and conventionally grown Italian cauliflower (*Brassica oleracea* L. subsp. *botrytis*): Results from a three-year field study. *J. Agric. Food Chem.* **2013**, *61*, 10335–10344. <https://doi.org/10.1021/jf4026844>.
54. Conversa, G.; Bonasia, A.; Lazzizzera, C.; Elia, A. Bio-physical, physiological, and nutritional aspects of ready-to-use cima di rapa (*Brassica rapa* L. subsp. *sylvestris* L. Janch. var. *esculenta* Hort.) as affected by conventional and organic growing systems and storage time. *Sci. Horticult.* **2016**, *213*, 76–86. <https://doi.org/10.1016/j.scienta.2016.10.021>.
55. Łata, B. Variability in enzymatic and non-enzymatic antioxidants in red and green-leafy kale in relation to soil type and N-level. *Sci. Horticult.* **2014**, *168*, 38–45. <https://doi.org/10.1016/j.scienta.2014.01.009>.
56. Picchi, V.; Migliori, C.; Lo Scalzo, R.; Campanelli, G.; Ferrari, V.; Di Cesare, L.F. Phytochemical content in organic and conventionally grown Italian cauliflower. *Food Chem.* **2012**, *130*, 501–509. <https://doi.org/10.1016/j.foodchem.2011.07.036>.
57. Valverde, J.; Reilly, K.; Villacreces, S.; Gaffney, M.; Granta, J.; Brunton, N. Variation in bioactive content in broccoli (*Brassica oleracea* var. *italica*) grown under conventional and organic production systems. *J. Sci. Food Agric.* **2015**, *95*, 1163–11761. <https://doi.org/10.1002/jsfa.6804>.
58. Ibrahim, M.H.; Jaafar, H.Z.E. Involvement of carbohydrate, protein and phenylalanine ammonia lyase in up-regulation of secondary metabolites in *Labisia pumila* under various CO₂ and N₂ level. *Molecules* **2011**, *16*, 4172–4190. <https://doi.org/10.3390/molecules16054172>.
59. Zhang, Y.; Ma, X.-M.; Wang, X.-C.; Liu, J.-H.; Huang, B.-Y.; Guo, X.-Y.; Xiong, S.P.; La, G.-X. UPLC-QTOF analysis reveals metabolomic changes in the flag leaf of wheat (*Triticum aestivum* L.) under low-nitrogen stress. *Plant Physiol. Bioch.* **2017**, *111*, 30–38. <https://doi.org/10.1016/j.plaphy.2016.11.009>.
60. Deng, B.; Li, Y.; Xu, D.; Ye, Q.; Liu, G. Nitrogen availability alters flavonoid accumulation in *Cyclocarya paliurus* via the effects on the internal carbon/nitrogen balance. *Sci. Rep.* **2019**, *9*, 2370. <https://doi.org/10.1038/s41598-019-38837-8>.
61. Sun, Y.; Guo, J.; Li, Y.; Luo, G.; Li, L.; Yuan, H.; Mur, L.A.J.; Guo, S. Negative effects of the simulated nitrogen deposition on plant phenolic metabolism: A meta-analysis. *Sci. Total Environ.* **2020**, *719*, 137442. <https://doi.org/10.1016/j.scitotenv.2020.137442>.
62. Pozzo, L.; Grande, T.; Raffaelli, A.; Longo, V.; Weidner, S.; Amarowicz, R.; Karamać, M. Characterization of antioxidant and antimicrobial activity and phenolic compound profile of extracts from seeds of different *Vitis* species. *Molecules* **2023**, *28*, 4924. <https://doi.org/10.3390/molecules28134924>.

Disclaimer/Publisher's Note: The statements, opinions and data contained in all publications are solely those of the individual author(s) and contributor(s) and not of MDPI and/or the editor(s). MDPI and/or the editor(s) disclaim responsibility for any injury to people or property resulting from any ideas, methods, instructions or products referred to in the content.