



Assessment of Biodegradation and Phytotoxicity of Azo Dyes Methyl Orange and Crystal Violet by *Escherichia coli* and *Bacillus subtilis*

Okoh EI ^{a*} and Immanuel OM ^b

^a Department of Microbiology, Ambrose Alli University, Ekpoma, Edo State, Nigeria.

^b Department of Biological Sciences, University of Africa, Toru-Orua, Bayelsa State, Nigeria.

Authors' contributions

This work was carried out in collaboration between both authors. Both authors read and approved the final manuscript.

Article Information

DOI: <https://doi.org/10.9734/sajrm/2024/v18i7369>

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/117428>

Original Research Article

Received: 26/03/2024

Accepted: 29/05/2024

Published: 03/06/2024

ABSTRACT

Azo dyes are widely used colourants, possess array of colours and are generally recalcitrant to microbial degradation. This study aimed to isolate azo dyes degrading bacteria from soil; treat methyl red and crystal violet dyes using the isolates, and assess the phytotoxicity of the treated and untreated dyes on sorghum. Ten bacteria were isolated from the soil, of which two were identified as efficient azo dyes degrading bacteria by efficiency in dye decolouration. The isolates were identified as *Escherichia coli* and *Bacillus subtilis* based on their biochemical characteristics. The isolates were able to degrade methyl orange dye by 16-75 % and crystal violet by 16-78%,

*Corresponding author: Email: okohenivweru@aauekpoma.edu.ng;

Cite as: EI, O., & OM, I. (2024). Assessment of Biodegradation and Phytotoxicity of Azo Dyes Methyl Orange and Crystal Violet by *Escherichia coli* and *Bacillus subtilis*. *South Asian Journal of Research in Microbiology*, 18(7), 1–8. <https://doi.org/10.9734/sajrm/2024/v18i7369>

depending on the concentration of the dye. The isolates were more efficient at degrading the dyes at a low concentration of 0.00025 mg/ml, as indicated by a higher percentage of decolourization (53-79%). The percentage phytotoxicity of the untreated methyl red ranged from 33-59%, while the toxicity of the untreated crystal violet ranged from 62-91%. The toxicity of the treated methyl red ranged from 9-34%, while the toxicity of the treated crystal violet ranged from 12-32%. The results revealed that the isolates were capable of degrading methyl red and crystal violet dyes, and could be useful in the bioremediation of azo dyes polluted soil.

Keywords: *Bacillus subtilis*; *bacteria*; *biodegradation*; *crystal violet*; *Escherichia coli*; *methyl red*; *phytotoxicity*.

1. INTRODUCTION

Dyes are the coloured substances which fix firmly with the substrate. There are different types of dyes classified according to their structures and functions. Azo dyes are the largest group of dyes used in the industry, representing more than half of the annual production [1,2]. Azo dyes are the largest class of synthetic dyes with the greatest variety of colours [3]. Different varieties of azo dyes are in use today in the consumer goods industries because of the diversity of dye components available for the synthesis of a large number of structurally different azo dyes [4,5]. Azo dyes are used by a wide number of industries because these compounds retain their colour and structural integrity under exposure to sunlight, soil, bacteria and sweat; they also exhibit high resistance to microbial degradation [6].

There is a continual demand to develop longer-lasting, more applicable dyes for items such as synthetic fabrics [7,8]. Azo dyes are continually updated to produce colours that reflect the trends dictated by changing social ideas and styles. From an environmental standpoint, the expansion in the demand and application of azo dyes is a threat to the safety of life which is never going away unless alternatives are found or better ways to manage them are devised.

While most azo dyes themselves are non-toxic a significantly larger portion of their metabolites are toxic [6,9,10]. Some dyes are carcinogenic and genotoxic because of the toxic potential of many aromatic amines [5,11]. Although several soil microorganisms can degrade dyes, they are often limited by the production of toxic by-products or the presence of toxic dye additives and also high salt concentrations [12]. Thus, these dyes are likely to persist in the environment and become toxic to biota.

The treatment of dye effluents is of interest due to their toxic and aesthetic impacts on receiving soil and water bodies. Effective and economic treatment of a diversity of effluents containing azo has become a problem. Currently, much research has been focused on chemically and physically degrading azo dyes in wastewaters, such as chemical oxidation, UV oxidation, electrochemical or wet oxidation, activated carbon adsorption, reverse osmosis, or coagulation/flocculation [2,13,14]. Many of these technologies are cost-prohibitive, however, and therefore are not viable options for treating large waste streams.

While much research has been performed to develop effective treatment technologies, no single solution has been satisfactory for remediating the diversity of wastes from dye industries [15]. This is because no single treatment system is adequate for degrading the diverse dye structures. However, the chemical structures of dyes can be modified by microbial metabolism [16,17]. Biodegradation of azo dyes is a viable option for the treatment of dye effluents, as it can bring about complete mineralization of the dye [18]. Several species of species have been reported to effectively decolourize azo dyes in a wide range of salt solutions [19-21]. Microorganisms decolourize dyes by desulphonation of their aromatic moiety coupled with reductive cleavage of the azo bond to prohibit aromatic amine formation, thereby, detoxifying the aromatic amines [20,22]. Also, the adsorption of the dye molecules to microbial cells has been implicated in the decolourization of dyes [23]. Several factors could affect the degradation of dyes by microorganisms, including factors related to the microorganisms themselves.

This study aims to isolate azo dyes degrading bacteria from the soil and assess their ability to degrade methyl orange and basic violet dyes, as

well as to assess the phytotoxicity of the dye on sorghum (*Sorghum bicolor*) seeds.

2. MATERIALS AND METHODS

2.1 Sample Collection

The soil samples were collected from three locations at the dumpsite of a laboratory-scale textile mill and printing press at the University of Port Harcourt, Port Harcourt. These samples were collected at a depth of 10cm in 10g amounts into a sterile Erlenmeyer flasks and composite sample was used.

The dyes used are methyl orange (Acid orange 52) with 85% purity and the dye crystal violet (Basic violet 3) with 95% purity (BDH Chemicals Ltd, England). Stock solutions of each dye were prepared by dissolving 0.1g of dye into 100 ml of the mineral salt medium.

2.2 Isolation of Dye Degrading Bacteria

One gram of each dye was added into a 25ml mineral salt medium, followed by the addition of 1g of soil sample. The flask was shaken properly and incubated in a shaker incubator (Barloworld Scientific Ltd, U.K) at 50rpm at 30°C. The cultures were observed daily for decolourization of the dye. After 14 days of incubation, 1ml of culture from each flask was serially diluted by Ten-fold serial dilution using physiological saline and spread plated onto mineral source agar medium in Petri dishes contained. Colonies which developed after incubation at 30°C were picked and repeatedly sub-cultured onto nutrient agar plates to obtain pure isolates.

2.3 Identification of Isolates

Isolates were identified following Gram reaction, motility, and spore formation, and based on their biochemical profiles following citrate, oxidase, methyl red, Voges-Proskauer, indole, TSI (Triple Sugar Iron), starch hydrolysis, sugar fermentation test.

2.4 Preparation of Standard Inoculum

Isolates were picked from the stock culture and each was inoculated into 10ml sterile nutrient broth contained in test tubes. The cultures were incubated at 37°C for 24 hours. These served as the standard inoculums for the various tests.

2.5 Screening of Isolates for the Degradation of Methyl Orange and Crystal Violets Dyes

All bacterial isolates were screened for the ability to degrade methyl orange and crystal violet dyes in test tubes containing 20mls of stock solution. Isolates that were able to decolourize the dyes were considered as dye degrading bacteria.

2.6 Degradation of Dye

A loopful of each isolate from the standard inoculums was inoculated into 10ml of sterile nutrient broth in test tubes. Cultures were incubated at 37°C for 24 hours. After incubation, 1ml from the culture was inoculated into 50ml of dye of different concentrations (0.00025mg/ml, 0.0005mg/ml, 0.001mg/ml and 0.002 mg/ml) contained in 250ml Erlenmeyer flask and incubated at 37°C.

Degradation was monitored daily for four days as follows: Two millilitres were withdrawn from each flask starting from day one and was centrifuged at 1000rpm for 30 minutes in an 800 Electric Centrifuge (Xiangtian, China). The supernatant was decanted and analysed.

The optical density was read using Spectrophotometer (EMCLAM, Germany) at 505nm for methyl orange and 588nm for crystal violet.

$$\text{Percentage decolourization} = \frac{\text{OD of control} - \text{OD of sample}}{\text{OD of Control}} \times 100$$

2.7 Phytotoxicity Assay

Phytotoxicity was conducted as per the method of Oyedeji and Immanuel [24] with modifications. Ten viable sorghum (*Sorghum bicolor*) seeds were pre-treated to prevent fungal growth by soaking them in sodium hypochlorite (3.5%w/v) solution for 8 minutes and subsequently rinsed ten times with sterile distilled water. Pre-treated seeds were placed on sterile Whatman No.1 filter papers in sterile Petri dishes. The Petri dishes were divided into three sets which consisted of set A (control experiment) which was irrigated with sterile distilled water; set B which was irrigated with untreated dye solution and set C which was irrigated with treated dye solution. The plates were irrigated daily with 5 ml of the dye solutions and were incubated at room

temperature (25°C). All experiments were carried out in triplicates. Seeds were considered to have germinated when root elongation was observed. The lengths of the radicles were recorded after 4 days. Toxic effect was measured in terms of percentage of phytotoxicity as:

$$\frac{\text{Percentage of phytotoxicity}}{\frac{\text{Radicle length of control} - \text{Radicle length of test}}{\text{Radicle length of control}}} \times 100 =$$

3. RESULTS

3.1 Isolation and Identification of Dye Degrading Bacteria

Table 1 shows two methyl red and crystal violet dyes degrading bacteria isolated from the soil samples.

3.2 Percentage Degradation

Fig. 1 shows results for biodegradation of the dyes by decolourization by Isolate A. Isolate A degraded methyl orange dye by 19-75% and crystal violet by 16-66%.

Fig. 2 shows results for biodegradation of the dyes by decolourization by Isolate B. Isolate B degraded methyl orange dye by 16-53 % and crystal violet by 18-79%.

Fig. 3 shows phytotoxicity results for dyes treated with isolate A and untreated dyes. The toxicity of untreated methyl red ranged from 33-59%; while the toxicity of crystal violet ranged from 62-91%. The toxicity of treated methyl red ranged from 9-34% while toxicity of crystal violet ranged from 12-32%.

Table 1. Biochemical characteristics of methyl red and crystal violet degrading bacteria

Gram stain	Shape	Spore	Motility	Oxidase	Citrate	Indole	MR	VP	Starch hydrolysis	TSI	Gas	Glucose	Sucrose	Lactose	Mannitol	Probable Organism
Isolate A	- Short rod	-	+	-	-	+	+	-	-	A	+	A/G	A/G	A/G	A/G	<i>E. coli</i>
Isolate B	+ Rod	+	+	-	+	-	-	+	+	A	+	A/G	A/-	-/-	A/-	<i>Bacillus subtilis</i>

Keys: A= Acid, G= Gas, - = negative), + = positive




Fig. 1. Percentage degradation of methyl orange and crystal violet dyes by Isolate A




Fig. 2. Percentage degradation of methyl orange and crystal violet dyes by Isolate B




Fig. 3. Phytotoxicity of dyes treated with isolate A and untreated dyes




Fig. 4. Phytotoxicity of dyes treated with isolate B and untreated dyes

Fig. 4 shows phytotoxicity results for dyes treated with isolate A and untreated dyes. Toxicity of untreated methyl red ranged from 40-89%; while the toxicity of crystal violet ranged from 64-93%. Toxicity of treated methyl red ranged from 22-56% while toxicity of crystal violet ranged from 10-52%.

4. DISCUSSION

Two azo dyes degrading bacteria were isolated from small-scale textile mill-contaminated soil. The isolates were identified as *E. coli* and *Bacillus subtilis*. Both isolates decolourized the methyl blue and crystal violet dyes. Decolourization of substrates is often associated with the degradation of the dye as decolourizing efficiency was dependent on the growth of the isolate in the flask [11,20]. There was neither growth nor decolourization in the control flasks. This evidence that the decolourization was due to the metabolic activity of the organisms and not due to any abiotic factors, in agreement with Oranusi and Ogugbue [20]. The dyes served as carbon, energy and nitrogen sources for the isolates as alluded to by Oranusi and Mbah [18].

Several bacterial species have been reported to degrade azo dyes including species of *Proteus*, *Pseudomonas*, *Sphingomonas*, *Actinobacillus*, *Mycobacterium*, *Bacillus* and *E. coli* [11,19,20]. *E. coli* and *Bacillus subtilis* have been reported to degrade methyl blue, a cationic dye, in earlier studies, on account of their high metabolic diversity [20,25]. The efficiency of azo dye biodegradation by the two isolates can be influenced by factors such as the initial dye concentration, the presence of co-substrates, and the cultivation conditions such as pH, temperature, and oxygen availability. Therefore, cultural conditions must be optimized to increase the efficiency of dye degradation. Aromatic amines generated by the reductive cleavage of azo dyes are potentially toxic, mutagenic and carcinogenic [5,11]. The reduction in the toxicity of the dyes after treatment as the isolates utilized the dyes as a source of nitrogen might have detoxified the parent compounds or any intermediates formed as alluded to by Chivukula and Renganathan [26] and Abadulla et al. [27].

In the present study, the general effect of the two dyes on germination followed the same patterns. The percentage of phytotoxicity of both dyes increased with an increase in concentration. Although higher concentrations of the dye exert a phytotoxic effect ranging from 47-93%, the lower

concentration of the dye also exerted an adverse phytotoxic effect on the seed by 33-64%. However, the treated dye exerted toxicity of 9-22% at the lowest concentration of 0.00025 mg/ml and toxicity of 32-56% at the highest concentration of 0.002 mg/ml. This reemphasized that indiscriminate disposal of azo dye-containing waste into soil and water bodies poses a major threat to the environment. The widely used phytotoxicity tests in vascular plants are the seed germination test (a direct exposure method) and the root elongation test [28,29]. Seed germination represents an important initial phase in the life cycle of plants. Any adverse effect on the seed germination process would be detrimental to the establishment and healthy growth of plants.

5. CONCLUSION

The present study showed that microbiological treatment of dyes is an efficient way to detoxify methyl red and crystal dyes. The toxic effect of dyes reported in this study suggests the need for remediation of textile dyes in the industry effluents before discharging them into the environment. Whatever treatment is adopted should ensure complete conversion of dye components to avoid the build-up of toxic metabolites.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Stolz A. Basic and Applied Aspects in the Microbial Degradation of Azo Dyes. *Applied Microbiology and Biotechnology*. 2001;56:69-80.
2. Ramalho PA, Paiva S, Cavaco-Paulo A, Casal M, Cardoso MH, Ramalho MT. Azo reductase activity of intact *Saccharomyces cerevisiae* cells is dependent on the Frelp component of the plasma membrane ferric reductase. *Appl. Environ. Technol.* 2005;71(7):3882-3888.
3. Zollinger H. *Colour chemistry: Synthesis, properties and applications of organic dyes and pigments*. 1991;496.
4. McCurdy MW. Chemical reduction and oxidation combined with biodegradation for the treatment of a Textile Dye Wastewater. Master's Thesis, Virginia Polytechnic Institute and State University. 1991;181.

5. Shi Y, Yang Z, Xing L, Zhang X, Li X, Zhang D. Recent advances in the biodegradation of azo dyes. *World Journal of Microbiology and Biotechnology*. 2021;37:137. Available:<https://doi.org/10.1007/s11274-021-03110-6>
6. Ramzan U, Shakoori FR, Zahid MT, Majeed W, Zahra I, Abbas SZ, Hedfi A, Hassa S, Shakoori AR, Al Mutery A. Biodegradation and Decolorization of Textile Azo Dyes by *Paramecium caudatum* Isolated from Industrial Wastewater. *Journals Water*. 2022;14:21. DOI:10.3390/w14213553
7. Padmavathy S, Sandhya S, Swaminathan K, Subrahmanyam YV, Chakrabarti T, Kaul SN. Aerobic Decolorization of Reactive Azo Dyes in Presence of Various Cosubstrates. *Chem. Biochem. Eng. Q*. 2003;17(2):147-151.
8. Moreira TA, Viacava C, Gladys V. Fed-batch Decolourization of Poly R-478 by *Trametes Versicolor*. *Braz. Arch. Biol. Technol*. 2004;47(2):45-65.
9. Ganesh R. Fate of Azo Dyes in Sludge. Master's Thesis, Virginia Polytechnic Institute and State University. 1992;193.
10. Prasad ASA, Satyanarayana VSV, Rao KVB. Biotransformation of Direct Blue 1 by a moderately halophilic bacterium *Marinobacter* sp. strain HBRA and toxicity assessment of degraded metabolites. *J. Hazard. Mater*. 2013;262:674–684.
11. Ayed L, Mahdhi A, Cheref A, Bakhrouf A. Decolorization and degradation of azo dye Methyl Red by an isolated *Sphingomonas paucimobilis*: Biotoxicity and metabolites characterization. *Desalination*. 2011;274:272–277. DOI: 10.1016/j.desal.2011.02.024
12. Suzuki Y, Yoda T, Ruhul A, Sugiura W. Molecular cloning and characterization of the gene coding for azoreductase from *Bacillus* sp. OY1-2 isolated from soil. *J. Biol. Chem*. 2001;276:9059–9065. DOI: 10.1074/jbc.M008083200
13. Edwards JC. Investigation of color removal by chemical oxidation for three reactive textile dyes and spent textile dye wastewater. Master's Thesis, Virginia Polytechnic Institute and State University. 2000; 55.
14. Naresh B, Jaydip J, Prabhat B, Rajkumar P. Recent biological technologies for textile effluent treatment. *Int. Res. J. Biol. Sci*. 2013;2:77–82.
15. Mani S, Chowdhary P, Bharagava RN. Textile wastewater dyes: Toxicity profile and treatment approaches. In *Emerging and Eco-Friendly Approaches for Waste Management*; Springer: Berlin/Heidelberg, Germany. 2019;219–244.
16. Sawhney R, Kumar A. Congo red (azo dye) decolourization by local isolate VTII inhibiting dye effluent exposed soil. *Int J. Environ. Sci*. 2011; 1:1261-7.
17. Rajee O, Patterson J. Decolourization of azo dye (orange MR) by an autochthonous bacterium *Micrococcus* sp. DBS 2. *Indian J. Microbiol*. 2011;51:159-63.
18. Oranusi NA, Mbah AN. Utilization of Azo and Triphenylmethyl Dyes as a Source of Carbon, Energy and Nitrogen by *Bacillus* sp. *African J. Appl. Zool. Environ. Biol*. 2005;7:87-94.
19. Jones JJ, Falkinham III JO. Decolourization of malachite green and crystal violet by waterborne pathogenic Mycobacteria. *Appl. Environ. Microbiol*. 2003;47(7):2323–2326.
20. Oranusi NA, Ogugbue CJ. Degradation of sulphonated azo dyes by *Pseudomonas* sp. *J. Appl. Sci. & Environ. Management*. 2011;5(2):13-17.
21. Dwivedi P, Kaur J, Dhanjal NIK, Cameotra SS. Biodecolorization of pararosaniline chloride dye using a bacterial isolate, *Arthrobacter* species. *J. Adv. Dev. Res*. 2012;3:77-81.
22. Elisangela F, Fabio Z, Cristiano DG, Regina DL, Artur CP. Biodegradation of textile azo dyes by a facultative *Staphylococcus arlettae* strain VN-11 using a sequential microaerophilic/aerobic process. *Int. Biodeterior. Biodegradation*. 2009;63:280-288.
23. Kadam AA, Lade HS, Patil SM, Govindwar SP. Low cost CaCl₂ pretreatment of sugarcane bagasse for enhancement of textile dyes adsorption and subsequent biodegradation of adsorbed dyes under solid state fermentation. *Bioresour. Technol*. 2013;32:276–284.
24. Oyedeji AA, Immanuel OM. Phytoremediation potential of *Senna fistula* L. in the decontamination of crude oil-polluted soil. *J. Appl. Sci. Environ. Manage*. 2023;27(12):2967-2972.
25. Ngo ACR, Tischler D. Microbial Degradation of Azo Dyes: Approaches and Prospects for a Hazard-Free Conversion by Microorganisms. *Int. J. Environ. Res. Public Health*. 2022;19:4740.

- DOI: 10.3390/ijerph19084740
26. Chivukula M, Renganathan V. Phenolic azo dye oxidation by laccase from *Pyricularia oryzae*. Appl. Environ. Microbiol. 1995;61:4373-4377.
 27. Abadulla E, Tzonov T, Costa S, Robra KH, Cavaco PA. Decolourization and detoxification of textile dyes with a laccase from *Trametes hirsute*. Appl. Environ. Microbiol. 2000;66(8):3357–3362.
 28. Oyedeji AA, Oyedeji FO, Immanuel OM, Adekoya OO. Assessing the potential of *Parkia biglobosa* (Jacq.) R.Br. ex. G. Don - A leguminous plant species commonly found in Nigeria to decontaminate crude oil-polluted soil in Terrestrial Ecosystem J. Appl. Sci. Environ. Manage. 2024; 28(2): 589-594.
 29. Oyedeji AA, Oyedeji FO, Immanuel OM. Role of *Pentaclethra macrophylla* Benth in crude oil-polluted soil remediation. World News of Natural Sciences. 2024;53:145-156.

© Copyright (2024): Author(s). The licensee is the journal publisher. This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:

<https://www.sdiarticle5.com/review-history/117428>