



Survey of the Intestinal Bacterial Compositions of Three Sympatric Passeriformes Species

Hisako Masuda ^{a*}, Fattaneh Nejad Zarabadi ^a
and Lina Rifai ^a

^a School of Sciences, Indiana University Kokomo, 2300 S Washington St. Kokomo, USA.

Authors' contributions

This work was carried out in collaboration among all authors. Authors LR and HM contributed to the study conception and design. Material preparation and data collection were performed by authors LR, FNZ and HM. Analysis was performed by authors LR and HM. The first draft of the manuscript was written by authors LR and HM. All authors commented on previous versions of the manuscript. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/JAMB/2022/v22i12689

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/93668>

Original Research Article

Received: 17/09/2022
Accepted: 19/11/2022
Published: 30/11/2022

ABSTRACT

Aims: The gut microbiota of avian species, especially wild birds, is studied less than that of mammals. Moreover, due to ethical concerns, a vast majority of microbiota studies have been limited to their feces. In this study, we investigated the microbial composition in the upper intestinal tract of sympatric, heterospecific bird species.

Study Design: We obtained samples from three wild bird species belonging in the same Order Passeriformes: Cedar Waxwing (*Bombycilla cedrorum*), American Robin (*Turdus migratorius*) and Dark-eyed Junco (*Junco hyemalis*). The samples were obtained from wild birds soon after they died due to physical injury. Microbial diversity was analyzed via high-throughput sequencing of 16S rRNA gene fragments.

Place and Duration of Study: Indiana University Kokomo campus, Indiana. United States, between October 2019 and December 2020.

*Corresponding author: E-mail: masudah@iuk.edu;

Methodology: Illumina Miseq sequencing of V3-V4 region of the 16S rRNA gene was employed to determine microbial composition. The α and β diversity were determined by Shannon's diversity index and principal coordinate analysis, respectively. Pair-wise comparison of Bray-Curtis dissimilarity between birds of same species, different species, collection dates and years were analyzed.

Results: The proteobacterial sequences were predominant in all samples. However, the comparison at the level of bacterial class revealed a significant intraspecies variation in both α and β diversity. Additionally, some microbiota exhibited higher interspecies similarity than intraspecies similarity, despite these birds not sharing the same foraging guilds.

Conclusion: Our study revealed high variability in the microbial compositions within the upper part of the digestive tract of closely related Passeriform birds in their natural environment.

Keywords: Gut microbiota; Cedar Waxwing; American Robin; Dark-eyed Junco; 16S rRNA.

1. INTRODUCTION

Gut microbial composition greatly impacts the physiology of the host. The microbiota can play a crucial role in physiological processes, such as digestion, vitamin production, protection against pathogens, as well as maturation of hosts' immune system and brain functions, including those that affect behavior [1-7].

Diet has been shown to have substantial impact on microbiota of animals, including birds [8-11]. Some studies have utilized artificial food to study the effects of a single component (e.g., the amount of dietary fiber or fat content) on microbiota [8,12]. Others have examined the impact of specific aspects of diet (e.g., calories) on microbiota [13]. Studying the impact of diet on microbiota of animals in natural environments is challenging, as organisms consume a variety of foods, often with seasonal variations.

Only a few bird species (2%) are strictly herbivorous [14,15], while 32% of all animals are herbivores [15]. Herbivory in birds is most likely rare due to the high metabolic needs of birds and fermentation being a relatively slow process. Many birds tend to switch their diets after their breeding seasons and/or during the colder months when less arthropod prey is available. During the colder months, their diet often become more frugivorous or granivorous [16,17]. This type of diet switch leads to an increase in gut length and change in microbial composition [18-20].

In addition to diet, other environmental and genetic factors can also alter gut microbiota. Hosts' social and physical environments, geographical distance, age, sex, active migration, and foraging strata have all been shown to contribute to the variations of

microbiota composition albeit at different degrees [21-23]. Additionally, host's genomes can have a significant impact on microbiota composition [24-26].

Many studies of gut microbiota have so far been limited to humans and commercial animals, while the study of microbiota of wild animals remains scarce. Additionally, the vast majority of these studies have been on mammalian species (i.e. for every avian study there are 10 mammalian ones), and the majority of those that have been conducted on avian species were done on domesticated birds [27].

Our primary aim for this study was to expand our current catalog of gut microbiota diversity of wild bird species, which is an understudied aspect of bird biology. Secondly, we examined the microbiota within the gut, rather than feces, to directly study their gut microbiota. For this purpose, we studied the microbiota of three wild bird species collected at the same location during the same season. Each bird species belongs to a different family within the same order, the Passeriformes: Cedar Waxwings (*Bombycilla cedrorum*, Bombycillidae), American Robins (*Turdus migratorius*, Turdidae), and Dark-eyed Juncos (*Junco hyemalis*, Passerellidae).

All three species of birds we tested are known to switch diets during the fall and winter season. American Robins switch from a mostly insectivorous diet to a mostly frugivorous one. Cedar Waxwings will switch from a mostly frugivorous to an almost exclusively frugivorous. Dark-eyed Juncos will switch from an omnivorous diet to a near exclusive granivorous diet [28]. Both Cedar Waxwings and American Robins have been observed to feed on cedar berries (Cedar Waxwings mostly), as well as fruits of crabapple and hawthorn [29-31]. Since

these birds have different forage patterns, our study also aimed to compare microbial compositions of sympatric, heterospecific species that belong to the same Order during the cold months.

While fecal samples are the predominant method from which to study gut microbiomes, differences do exist between microbial composition in the gut and in feces [32]. Even within the gut, the composition in gizzard, upper intestine, and lower intestine are not identical [33]. Since we are interested in directly measuring gut microbial composition, we collected samples from the upper intestinal tract of birds in natural habitat. To reduce the chance of postmortem change in microbial composition, samples were collected immediately after their death due to accidental collision with windows.

2. MATERIALS AND METHODS

2.1 Samples

Four Cedar Waxwings (samples CW1-4), three American Robins (samples R1-3), and two Dark-eyed Juncos (samples DJ1-2) were collected on the Indiana University Kokomo campus (Kokomo, Indiana USA), except for sample DJ2 which was collected one mile south of campus. The samples were collected as part of daily monitoring for birds that are injured or have died due to window collisions. To minimize the effects of postpartum change in bacterial composition, only recently deceased birds were processed further for DNA extraction. Injured birds were transported immediately to a bird rehabilitation center. Only birds that were collected in October and November of 2019 and 2020 were analyzed. The information and sample names are summarized in the Table 1. After collecting recently deceased birds, they were either immediately processed or placed in the freezer until dissection. Once birds were dissected, the intestines were removed in sections using sterile dissection equipment. The upper intestine was then placed into a sterile container and kept at -80°C until DNA extraction.

2.2 DNA Extraction and Sequencing

The DNA was extracted from 100 mg of each sample using the E.Z.N.A. Stool DNA kit (Omega-Biotek Inc., GA, USA) according to the

manufacturer's instructions. RNA was removed by RNase treatment at 37°C . The extracted DNA was visualized with 1% agarose gel electrophoresis and stained with ethidium bromide solution. The concentration and quality of the DNA was calculated from the measurement of absorbance at 260 nm and 280 nm using the spectrophotometer. A_{260}/A_{280} between 1.80 and 1.90 were further processed for sequencing. 16S rRNA DNA library of V3-V4 region was constructed and sequenced with Illumina MiSeq at Psamogen Inc. (MD, USA). Between 101,826 and 172,376 raw reads were generated.

2.3 Bioinformatic and Statistical Analyses

The Illumina MiSeq sequence reads were assembled and processed using micca [34]. Reads were trimmed to remove barcode and primer sequences. Low quality reads with average quality scores of less than 25 were discarded. Assembled reads from each sample were classified into operational taxonomic unit (OTU) of 97% sequence similarity. Taxonomic information was assigned to each OTU using the Ribosomal Database Project (RDP) classifier [35]. The rarefaction curve was created by the vegan package of R [36]. A single OTU matched a chloroplast-derived sequence. This sequence was removed from further analysis. A phyloseq object was created using phyloseq [37] and further analyzed by R [38]. The reads were rarified to an even depth. The multiple sequence alignment was performed using msa command using the Nearest Alignment Space Termination method [39]. Green gene core set was used as a template. The rooted phylogenetic tree was constructed.

The pairwise dissimilarities were calculated using Jaccard distance, Bray-Curtis dissimilarity index, weighted and unweighted UniFrac distances with packages vegan and phyloseq [36,37]. Principal coordinate analysis (PCoA), RDA, and CCA methods were employed to analyze the dissimilarity. Since similar conclusions were drawn from each result, not all of the results are shown. Hierarchical clustering of dissimilarity was calculated using hclust with the Ward's method. Permanova and pairwise Adonis function of vegan was utilized for beta diversity hypothesis testing. PCoA, heatmap and bar plots were constructed using vegan, ggplot2, and ggbiplot in R [40,41].

Table 1. The summary of sample types and collection dates

Sample ID	Collection date	Common name	Order	Family	Species	Diet change when transitioning from summer to fall [28]
CW1	10/8/2020	Cedar Waxwings	Passeriformes	Bombycillidae	<i>Bombycilla cedrorum</i>	Mostly frugivorous to almost exclusively frugivorous
CW2	11/10/2020					
CW3	11/16/2020					
CW4	10/7/2020					
R1	10/30/2020	American Robins	Passeriformes	Turdidae	<i>Turdus migratorius</i>	Mostly insectivorous to a mostly frugivorous
R2	11/6/2019					
R3	11/28/2019					
DJ1	10/12/2020	Dark-eyed Juncos	Passeriformes	Passerellidae	<i>Junco hyemalis</i>	Omnivorous to almost exclusively granivorous
DJ2	11/25/2019					

3. RESULTS

Between 101,826 and 172,376 reads were obtained for each sample by Illumina MiSeq sequencing. The rarefaction curve shows that enough sequencing depth was achieved to capture the diversity within each sample (Fig. 1). The reads were classified into 633 total OTUs. The alpha diversity of each sample was analyzed using the Shannon's diversity index (Fig. 2). There were large variations in diversity index in all species (Fig. 2a). There were no statistically significant differences in diversity between species (Mann Whitney U test, $P = 0.27$ for Cedar Waxwing vs Dark-eyed Junco; $P = 0.4$ for Dark-eyed Junco vs American Robin; and $P =$

0.86) for Cedar Waxwing vs. American Robin). Also, no significant difference was detected for alpha diversity of samples collected in the year 2019, compared to the year 2020 ($P = 0.73$, Fig. 2b). The collection month (October vs. November) also did not impact the alpha diversity ($P = 0.29$).

The beta diversity of the samples was then compared using the principal coordinate plot (Fig. 3). The principal coordinate plot, displaying the weighted and unweighted UniFrac distance, does not show distinct clustering of samples based on taxonomic group (species) nor based on the collection year or month (Fig. 3).




Fig. 1. Rarefaction curve of the Illumina reads
Sample IDs are listed in the Table 1




Fig. 2. Comparison of alpha diversity index

The Shannon diversity index of OTUs in each sample is grouped according to the (a) bird species and (b) sample collection year. There were no differences between the groups with statistical significance




Fig. 3. Principal coordinate analysis of beta diversity based on weighted or unweighted UniFrac distance

Unweighted (a and c) or weighed (b and d) UniFrac distance methods were used. The samples are colored according to the species (pink: Cedar Waxwings, green: Dark-eyed Juncos, and blue: American Robins). The shape represents the month (a and b) or year (c and d) of sample collection date

Each sample contained 60 to 315 unique OTUs. The taxonomy was assigned for each OTU. In four samples (CW1, CW2, R1, and R2), a single OTU composed a significant portion of the microbiota (74%, 98%, 71%, and 84% of total sequences, respectively). The most abundant OTU in a Cedar Waxwing sample (CW2) and an American Robin sample (R1) belong to the genus *Escherichia/Shigella* (OTU1). The most abundant OTU of the sample CW1 is *Helicobacter* (OTU2). Another *Helicobacter* sequence (OTU5) was also abundant in other Cedar Waxwing samples: in CW1 (11%) and CW3 (29%). *Yersinia* sequence (OTU3) was only abundantly present in sample R2 (57%). The second most abundant OTU (OTU10), which comprised 26% of the sample R2, was also *Yersinia* sequence.

There are many other proteobacterial sequences that were prevalently present, including *Campylobacter* (OTU4), Enterobacteriaceae (OTU6 and OTU11), and *Aeromonas* (OTU7). The phyla Firmicutes, Tenericutes, Deinococcus-Thermus and Actinobacteria were also found in multiple samples at a high abundance. The taxonomy and relative abundance of OTU in each sample are visualized in Fig. 4. The most abundant bacterial phyla were Proteobacteria

(colored brown in the Fig. 4a) in all samples. Firmicutes (colored light green) were also highly represented in samples CW4, DJ1, and DJ2, and R3, respectively.

Despite proteobacteria being uniformly the most abundant bacterial phyla, different classes of proteobacteria were abundant in each sample (Fig. 4b). In CW1 and CW3, epsilonproteobacteria was the most abundant class. As discussed above, the most abundant sequences in these two samples were that of genus *Helicobacter* (OTU2 and 5) and *Campylobacter* (OTU4), which both belong to epsilonproteobacteria (Fig. 4c). In other samples, gammaproteobacterial sequences were abundant (Fig. 4b). *Escherichia/Shigella* (OTU1) and *Yersinia* (OTU3 and 10) belong to the class gammaproteobacteria (Fig. 4c). Additionally, the most abundant OTU in CW4 (OTU6) was a gammaproteobacterial sequence with no known homolog at the genus level. Fig. 5a summarizes the fraction of each OTU present in each sample. Samples CW4, R3, and DJ2 contained many unique OTUs that were not present in others. Other samples contained relatively smaller numbers of OTUs, but these OTUs are present in most samples.




Fig. 4. The relative abundance of OTUs classified into unique taxonomic groups Taxonomic classification at (a) phylum, (b) class, and (c) genus is shown. The taxonomic units that represent greater than 1% abundance are included. The sample IDs are summarized in the Table 1




Fig. 5. The heatmap and cluster dendrogram of OTUs in each sample (a) Heatmap of the OTU with the abundances greater than 10^{-5} are shown. (b) Ward's hierarchical agglomerative clustering method was applied to the Bray-Curtis distance between samples. Abbreviations: CW: Cedar Waxing; DJ: Dark-eyed Junco; and R: American Robin. (c) Pair-wise comparison of Bray-Curtis dissimilarity index. The dissimilarity of microbiota of samples belonging to the same species (right) and dissimilarity between samples belonging to different species (left) are shown




Fig. 6. The Venn diagram of unique OTUs observed in each species

The numbers of OTUs that were present in at least one sample from each species were counted

Pairwise comparison of Bray-Curtis dissimilarity was measured. The dendrogram of Bray-Curtis distance shows clustering of 1) CW1 and CW3, 2) CW2 and R1, 3) CW4 and DJ1, and 4) R3 and DJ2 (Fig. 5b). An intraspecies pair (CW1 and CW3) has a similar microbiota comparison. However, there were certain pair of samples with low dissimilarity (e.g. CW2 and R1; CW4 and DJ1). We then pooled a pairwise comparison of Bray-Curtis dissimilarity into 1) between the intra-species and 2) inter-species samples (Fig. 5c). This comparison did not show a statistically significant difference (Mann Whitney U test, $P = 0.87$) (Fig. 5c). A pair of microbiota from different species (R1 and CW2) had low dissimilarity index of 0.282. This is lower than the smallest dissimilarity index when two of the CW samples were compared (0.537 for CW1 and CW3). Pairwise comparison based on collection year or month also did not show a significant difference ($P = 0.48$ and $P = 0.30$, respectively).

Fig. 6 displays the number of OTUs that are unique to each species as well as commonly found in two or three bird species. The values represent the number of unique OTUs that were found in at least one sample in each species. Out of 632 total OTUs found in this study, 73 OTUs (12% of total) were detected in all three species. Dark-eyed Juncos, Cedar Waxwings, and American Robins carried 92, 129, and 242 unique OTUs that were not found in other species, respectively.

4. DISCUSSION

In this study, we discovered that three sympatric Passeriformes species in the wild exhibited high intraspecies variation of microbiota composition

in the upper intestinal tract. While uniformly the most abundant phylum was proteobacteria, a high dissimilarity of composition at the genus and class level was observed. Because we relied on our sample collection from birds that accidentally died due to collision with windows, the numbers of our sample were limited. In particular, we were able to collect only two Dark-eyed Junco samples, limiting our ability to draw conclusions with statistical significance. However, our approach allows an ethical means to gain samples from the intestinal tract of birds in their natural habitats and study their microbiota.

Our personal observations on campus of both Cedar Waxwings and American Robins have confirmed that during the fall and winter months they will flock to trees to feed on berries. If diet is the major determining factor in birds for microbiota similarities, then the gut microbiota of Dark-eyed Juncos should have been more unique compared to the two other species. However, overall gut microbial composition of Dark-eyed Juncos was not significantly different from the two berry-eating species. Our finding mirrors that of Song and colleagues, who found that the microbiota of birds and bats have very little correlation to host diet [42].

Our results match a trend of weaker association between host's phylogeny and microbiota variation in birds that have been observed previously [43,44]. The striking similarity between birds of different species suggests the possibility that interspecies exchange of gut microbes happens among sympatric birds. Further investigation to search for the cause of observed high intraspecies dissimilarity, as well as interspecies similarity, such as examining the

birds' individual migratory history, food content in the gut, and sex, is needed.

Flighted birds and bat microbiota generally have fewer bacteria responsible for food fermentation, while the number of proteobacteria is higher. A relatively high amount of proteobacteria have been observed in feces of Passeriform species [22]. In wild neotropical birds, the proportion of proteobacteria is higher in bird consuming fruits than in birds consuming grains [33]. Our study also detected a high proportion of proteobacteria in all samples except for a relatively smaller proportion in a robin sample (R3). Whether or not the difference between R3 and other samples reflects the types of diet immediately before sampling requires further investigation.

The most common OTUs belonged to the genus *Escherichia/Shigella*, *Helicobacter*, *Yersinia*, *Campylobacter*, unknown genus of Enterobacteriaceae, and *Aeromonas*. Although our study does not allow identification of specific function, some of the members of these genera are known pathogens [45]. The presence of human pathogens in bird feces has been described [46]. The roles of proteobacteria in the gut of flighted animals are of interest not only for understanding their diet but also for epidemiological concerns.

The postmortem change in gut microbial composition has been studied in humans [47]. Increases in *Bacteroides* and *Lactobacillus* have been detected, but changes were minimum within the first 24 hours. No significant increase in the abundance of proteobacteria was observed. It remains unknown whether the same sets of bacterial species will change abundance in birds postmortem as in humans. Since our samples were relatively fresh (from a few minutes up until a few hours), the postmortem change should be low. However, a future study to determine the time-dependent change postmortem of microbial composition in bird species will be needed.

5. CONCLUSION

In conclusion, our study has shown that birds of Passeriformes in natural habitats carry highly diverse sets of gut bacteria. This suggests the significant effects of diet and/or other environmental factors on their microbiota. Due to the presence of bacterial strains with known associations with pathogenicity in humans (e.g. *Escherichia/Shigella*, *Helicobacter*, and

Campylobacter), this is an area that deserves further attention and studies.

ACKNOWLEDGEMENT

The authors thank Dr. Peter Tupa for proofreading the manuscript. This research was supported by Grants-in Aid and Applied Learning Grants from Indiana University Kokomo. The authors also acknowledge the funding support from the IUK School of Sciences.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Diaz Heijtz R, Wang S, Anuar F, Qian Y, Björkholm B, Samuelsson A, et al. Normal gut microbiota modulates brain development and behavior. *Proc Natl Acad Sci U S A*. 2011;108:3047-52. DOI:10.1073/pnas.1010529108
2. Al-Asmakh M, Stukenborg JB, Reda A, Anuar F, Strand ML, Hedin L, et al. The gut microbiota and developmental programming of the testis in mice. *PLoS One*. 2014;9:e103809. DOI:10.1371/journal.pone.0103809
3. Rosenbaum M, Knight R, Leibel RL. The gut microbiota in human energy homeostasis and obesity. *Trends in Endocrinology & Metabolism*. 2015;26:493-501.
4. Buffie CG, Pamer EG. Microbiota-mediated colonization resistance against intestinal pathogens. *Nature Reviews Immunology*. 2013;13:790-801.
5. Chung H, Pamp SJ, Hill JA, Surana NK, Edelman SM, Troy EB, et al. Gut immune maturation depends on colonization with a host-specific microbiota. *Cell*. 2012;149:1578-93. DOI:10.1016/j.cell.2012.04.037
6. LeBlanc JG, Milani C, De Giori GS, Sesma F, Van Sinderen D, Ventura M. Bacteria as vitamin suppliers to their host: a gut microbiota perspective. *Current Opinion in Biotechnology*. 2013;24:160-168.
7. Hoban AE, Stilling RM, Moloney G, Shanahan F, Dinan TG, Clarke G, et al. The microbiome regulates amygdala-dependent fear recall. *Molecular Psychiatry*. 2018;23:1134-1144.

8. Carmody Rachel N, Gerber Georg K, Luevano Jesus M, Jr., Gatti Daniel M, Somes L, Svenson Karen L, et al. Diet Dominates Host Genotype in Shaping the Murine Gut Microbiota. *Cell Host & Microbe*. 2015;17:72-84.
DOI:10.1016/j.chom.2014.11.010
9. David LA, Maurice CF, Carmody RN, Gootenberg DB, Button JE, Wolfe BE, et al. Diet rapidly and reproducibly alters the human gut microbiome. *Nature*. 2014; 505:559-63.
DOI:10.1038/nature12820
10. Sonnenburg ED, Smits SA, Tikhonov M, Higginbottom SK, Wingreen NS, Sonnenburg JL. Diet-induced extinctions in the gut microbiota compound over generations. *Nature*. 2016;529:212-215.
11. Griffin NW, Ahern PP, Cheng J, Heath AC, Ilkayeva O, Newgard CB, et al. Prior dietary practices and connections to a human gut microbial metacommunity alter responses to diet interventions. *Cell Host & Microbe*. 2017;21:84-96.
DOI:https://doi.org/10.1016/j.chom.2016.12.006
12. Tap J, Furet JP, Bensaada M, Philippe C, Roth H, Rabot S, et al. Gut microbiota richness promotes its stability upon increased dietary fibre intake in healthy adults. *Environ Microbiol*. 2015;17:4954-64.
DOI:10.1111/1462-2920.13006
13. Dong TS, Luu K, Lagishetty V. A high protein calorie restriction diet alters the gut microbiome in obesity. *Nutrients*. 2020;12:3221.
DOI:10.3390/nu12103221
14. Burin G, Kissling WD, Guimarães PR, Şekercioğlu ÇH, Quental TB. Omnivory in birds is a macroevolutionary sink. *Nature Communications*. 2016;7:1-10.
15. Román-Palacios C, Scholl JP, Wiens JJ. Evolution of diet across the animal tree of life. *Evolution Letters*. 2019;3:339-347.
16. Biebach H. Energetics of winter and migratory fattening. *Avian energetics and nutritional ecology*. Springer. 1996;280-323.
17. Karasov WH. Digestive plasticity in avian energetics and feeding ecology. *Avian energetics and nutritional ecology*. Springer, New York. 1996;61-84.
18. McWilliams SR, Karasov WH. Phenotypic flexibility in digestive system structure and function in migratory birds and its ecological significance. *Comparative Biochemistry and Physiology Part A: Molecular & Integrative Physiology*. 2001;128:577-591.
19. Bodawatta KH, Freiberga I, Puzejova K, Sam K, Poulsen M, Jønsson KA. Flexibility and resilience of great tit (*Parus major*) gut microbiomes to changing diets. *Animal Microbiome*. 2021;3:20.
DOI:10.1186/s42523-021-00076-6
20. Teyssier A, Matthysen E, Hudin NS, de Neve L, White J, Lens L. Diet contributes to urban-induced alterations in gut microbiota: experimental evidence from a wild passerine. *Proc Biol Sci*. 2020; 287:2019-2182.
DOI:10.1098/rspb.2019.2182
21. Yatsunenkov T, Rey FE, Manary MJ, Trehan I, Dominguez-Bello MG, Contreras M, et al. Human gut microbiome viewed across age and geography. *Nature*. 2012;486:222-227.
22. Hird SM, Sánchez C, Carstens BC, Brumfield RT. Comparative gut microbiota of 59 neotropical bird species. *Frontiers in Microbiology*. 2015;6:1403.
23. Risely A, Waite DW, Ujvari B, Hoyer BJ, Klaassen M. Active migration is associated with specific and consistent changes to gut microbiota in *Calidris* shorebirds. *Journal of Animal Ecology*. 2018;87:428-437.
24. Kemis JH, Linke V. Genetic determinants of gut microbiota composition and bile acid profiles in mice. *PLoS Genet*. 2019; 15:e1008073.
DOI:10.1371/journal.pgen.1008073
25. Ochman H, Worobey M, Kuo CH, Ndjango JB, Peeters M, Hahn BH, et al. Evolutionary relationships of wild hominids recapitulated by gut microbial communities. *PLoS Biol*. 2010; 8:e1000546.
DOI:10.1371/journal.pbio.1000546
26. Amato KR, Sanders JG, Song SJ, Nute M, Metcalf JL, Thompson LR, et al. Evolutionary trends in host physiology outweigh dietary niche in structuring primate gut microbiomes. *The ISME Journal*. 2019;13:576-587.
27. Grond K, Sandercock BK, Jumpponen A, Zeglin LH. The avian gut microbiota: community, physiology and function in wild birds. *Journal of Avian Biology*. 2018; 49:e01788.

28. Martin AC, Zim HS, Nelson AL. American wildlife & plants: a guide to wildlife food habits: the use of trees, shrubs, weeds, and herbs by birds and mammals of the United States: Courier Corporation; 1961.
29. Witmer MC. Annual diet of cedar waxwings based on US biological survey records (1885-1950) compared to diet of American Robins: contrasts in dietary patterns and natural history. *The Auk*. 1996;113:414-430.
30. Sallabanks R. Fruit fate, frugivory, and fruit characteristics: a study of the hawthorn, *Crataegus monogyna* (Rosaceae). *Oecologia*. 1992;91:296-304.
31. Wheelwright NT. The diet of American Robins: an analysis of US Biological Survey records. *The Auk*. 1986;103:710-725.
32. Waite DW, Taylor MW. Characterizing the avian gut microbiota: membership, driving influences, and potential function. *Frontiers in Microbiology*. 2014;5. DOI:10.3389/fmicb.2014.00223
33. García-Amado MA, Shin H, Sanz V, Lentino M, Martínez LM, Contreras M, et al. Comparison of gizzard and intestinal microbiota of wild neotropical birds. *PLoS One*. 2018;13:e0194857. DOI:10.1371/journal.pone.0194857
34. Albanese D, Fontana P, De Filippo C, Cavalieri D, Donati C. MICCA: a complete and accurate software for taxonomic profiling of metagenomic data. *Scientific Reports*. 2015;5:9743. DOI:10.1038/srep09743
35. Wang Q, Garrity GM, Tiedje JM, Cole JR. Naive Bayesian classifier for rapid assignment of rRNA sequences into the new bacterial taxonomy. *Appl Environ Microbiol*. 2007;73:5261-7. DOI:10.1128/aem.00062-07
36. Dixon P. VEGAN, a package of R functions for community ecology. *Journal of Vegetation Science*. 2003;14:927-930. DOI:https://doi.org/10.1111/j.1654-1103.2003.tb02228.x
37. McMurdie PJ, Holmes S. phyloseq: an R package for reproducible interactive analysis and graphics of microbiome census data -. *PLoS One*. 2013;8:e61217. Available:https://bioconductor.org/packages/release/bioc/html/phyloseq.html DOI:10.1371/journal.pone.0061217
38. Ihaka R, Gentleman RR. A language for data analysis and graphics. *Journal of Computational and Graphical Statistics*. 1996;5:299-314. DOI:10.1080/10618600.1996.10474713
39. DeSantis TZ, Jr, Hugenholtz P, Keller K, Brodie EL, Larsen N, Piceno YM, et al. NAST: a multiple sequence alignment server for comparative analysis of 16S rRNA genes. *Nucleic Acids Research*. 2006;34:W394-W399. DOI:10.1093/nar/gkl244
40. Wickham H. ggplot2: Elegant graphics for data analysis. Available:https://cran.r-project.org/web/packages/ggplot2/index.html (Accessed 30 Jan 2021), 2016
41. Vincent QV. 2011. ggbiplot: A ggplot2 based biplot. R package version 0.55. Available:http://github.com/vqv/ggbiplot (Accessed 30 Jan 2021).
42. Song SJ, Sanders JG, Delsuc F, Metcalf J, Amato K, Taylor MW, et al. Comparative analyses of vertebrate gut microbiomes reveal convergence between birds and bats. *MBio*. 2020;11:e02901-19.
43. Youngblut ND, Reischer GH, Walters W, Schuster N, Walzer C, Stalder G, et al. Host diet and evolutionary history explain different aspects of gut microbiome diversity among vertebrate clades. *Nature Communications*. 2019;10:2200. DOI:10.1038/s41467-019-10191-3
44. Teyssier A, Lens L, Matthysen E, White J. Dynamics of gut microbiota diversity during the early development of an avian host: evidence from a cross-foster experiment. *Frontiers in Microbiology*. 2018;9. DOI:10.3389/fmicb.2018.01524
45. Kirk MD, Pires SM, Black RE, Caipo M, Crump JA, Devleeschauwer B, et al. World Health Organization estimates of the global and regional disease burden of 22 foodborne bacterial, protozoal, and viral diseases, 2010: A data synthesis. *PLoS Med*. 2015;12:e1001921. DOI:10.1371/journal.pmed.1001921
46. Byappanahalli MN, Nevers MB, Korajkic A, Staley ZR, Harwood VJ. Enterococci in the environment. *Microbiology and Molecular Biology Reviews* : MMBR. 2012;76:685-706. DOI:10.1128/MMBR.00023-12

47. Hauther KA, Cobaugh KL, Jantz LM, Sparer TE, DeBruyn JM. Estimating time since death from postmortem human gut microbial communities. *Journal of Forensic Sciences*. 2015;60:1234-40. DOI:10.1111/1556-4029.12828

© 2022 Masuda et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:
The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/93668>