



## **Association between Prenatal Viral Infections and the Occurrence of Schizophrenia in Adulthood**

**Ioanna Kotsiri<sup>a</sup>, Alexandros Spyranitis<sup>b</sup>, Charalampos Panotopoulos<sup>c</sup> and Emmanouil Magiorkinis<sup>d\*</sup>**

<sup>a</sup> *Department of Internal Medicine, Asklepieion General Hospital, Voulas, Athens, Greece.*

<sup>b</sup> *General Hospital Ag. Anargiroi Kifisia, Athens, Greece.*

<sup>c</sup> *Department of Clinical Cardiology, General Hospital of Sparti, Sparti, Greece.*

<sup>d</sup> *Department of Laboratory Medicine, Sotiria General Hospital for Chest Diseases, Athens, Greece.*

### **Authors' contributions**

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

### **Article Information**

DOI: 10.9734/JAMMR/2022/v34i2031480

### **Open Peer Review History:**

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/87749>

**Review Article**

**Received 06 May 2022**

**Accepted 11 July 2022**

**Published 13 July 2022**

### **ABSTRACT**

Schizophrenia is a severe mental disorder that affects approximately 1% of the population worldwide. Genetics and environmental factors appear to contribute to the pathogenesis of schizophrenia. Epidemiological studies have focused on the relationship between viral infection during pregnancy and an increased risk of developing schizophrenia in offspring. The aim of our review is to summarize whether viral infections during pregnancy can lead to psychiatric disorders and especially to schizophrenia in adulthood. For that purpose we undertaken a thorough search in Pubmed using as keywords viral infections, schizophrenia, perinatal infections and virus. Epidemiological studies have focused on the relationship between perinatal viral infections and an increased risk of developing schizophrenia in offspring. Influenza virus, various members of the Herpesviridae family, retroviruses, borna virus are discussed as examples of viral infections associated with schizophrenia. These viral infections appear to interfere with the normal maturation of the brain and may lead to the subsequent onset of schizophrenia. Another epidemiological aspect of schizophrenia is the fact that it presents a seasonal prevalence with peaks in offsprings which wer born during winter and spring. It is also possible for an infection occurring in infancy to reactivate later, as is known to happen with herpes viruses. Overall, various studies showed that

\*Corresponding author: E-mail: [mayiork@med.uoa.gr](mailto:mayiork@med.uoa.gr), [emmanouil.magiorkinis@gmail.com](mailto:emmanouil.magiorkinis@gmail.com);

prenatal maternal infections appear to increase the risk of developing schizophrenia in the offspring, and this is due to the body's immune response to the infection, and not in exposure to specific viral or bacterial pathogens.

**Keywords:** *Pregnancy and viral infections; schizophrenia; congenital infections; viral infections and schizophrenia; prenatal viral infections.*

## 1. INTRODUCTION

Schizophrenia is a severe mental disorder characterized by cognitive disturbance. It affects approximately 1% of the population worldwide. The disorder has a diverse clinical picture and the first symptoms appear during puberty or early adulthood. People with schizophrenia may demonstrate various symptoms such as hallucinations, disorganized thinking, delusions, social withdrawal, and decreased emotional expression [1,2]. Additional symptoms that can be seen in schizophrenia are sleeping and eating disorders, and an increased risk of co-morbidities such as obesity, diabetes, arterial hypertension, and coronary disease [3].

The symptoms of schizophrenia according to eleventh revision of the International Classification of Diseases (ICD-11) the diagnostic criteria of schizophrenia or other primary psychotic disorders (Schizoaffective disorder, Schizotypal disorder, acute and transient psychotic disorder (ATPD), delusional disorder) are categorized into positive and negative. The diagnosis of schizophrenia requires at least two or more symptoms and at least one of the two symptoms must be a positive symptom. Positive symptoms are hallucinations, delusions, disorganized speech and abnormal movements. Negative symptoms are lack of motivation and initiative blunt feeling, social withdrawal, neglect of the self, alogia, anhedonia, and lack of emotion, on a 4-point scale ranging from 0 = not present to 3 = present and severe [4].

Schizophrenia subtypes are omitted in ICD 11. Schizophrenia symptom characteristics and the duration criterion in ICD-11 have largely remained the same compared to ICD-10. Symptoms must be present for at least one month. The changes from ICD-10 to ICD-11. ICD-10 included four subtypes of ATPD. ICD10 Acute Polymorphic Psychotic Disorder Without Symptoms of Schizophrenia largely corresponds to ICD-11 Acute and Transient Psychotic Disorder. ICD10 Acute polymorphic Psychotic

Disorder with Symptoms of Schizophrenia and ICD-10 Acute schizophrenia-like Psychotic Disorder account for presentations of schizophrenia-like symptoms that are too short to qualify for a diagnosis of schizophrenia, and are thus included in the diagnosis of Other Primary Psychotic Disorder in ICD-11 [4].

The causes of schizophrenia are mostly unknown. It is believed that genetics may play the main role in the onset of the disorder, due to its increased frequency among first-degree relatives. However, its appearance among monozygotic twins is approximately 50% which shows that environmental factors may play a role in the manifestation of schizophrenia and causing changes in the structure of the fetal brain increasing the risk of schizophrenia in individuals with a genetic predisposition [1,5]. The obstetric complications during labor, poor or inadequate prenatal diet, prenatal exposure of pregnant women to viral infections, the period of labor, labor in urban settings, smoking during adolescence are some of the environmental factors which may affect fetal brain development and lead to neurodevelopmental disorders which will increase the risk of appearance of schizophrenia in adulthood life [6-9].

Therapy of schizophrenia at an early stage are mainly antipsychotics. They work by blocking the effect of the chemical dopamine on the brain. Antipsychotics reduce the stress and irritability that the patient may feel when experiencing an episode of schizophrenia, while a few days later they can also reduce the symptom of hallucinations and delusions. There are first generation and second generation antipsychotics (FGAs). First-generation antipsychotics are effective in treating positive symptoms but have no effect on negative symptoms. Second-generation antipsychotics (SGAs) are effective in treating both the positive and negative symptoms of schizophrenia. SGAs have the additional effect of competing with 5HT serotonin receptors along with D2 receptors that cause fewer extrapyramidal symptoms compared to FGAs. Clozapine is intended for treatment-resistant schizophrenia [10].

## 1.1 Pathophysiology of Schizophrenia

It is difficult to understand the pathophysiology of schizophrenia since its causes are not fully known.

It is thought that there is a disorder of the neurotransmitter system and mainly the dopamine system, which shows increased production and activity in the limbic system of the brain. The serotonin level is increased in these patients due to stress and block the dopaminergic neurons. The glutamine neurotransmitter presents reduced activity at the prefrontal cortex of the brain. Schizophrenia is the result of a disturbance in the balance of neurotransmitters such as dopamine, glutamic and gamma-aminobutyric acid (GABA) or by changes in the biochemical pathways associated with inflammation (kynurenic acid). In patients with schizophrenia, changes in dopamine function within the striatum cause delusions and psychosis [11-14].

The neurodevelopmental disorder theory is supported by many researchers [15-19]. These disorders affect the temporal structures such as the hippocampus [19] which appears in the stages of prenatal development and many years before the onset of psychosis [17]. These disorders may affect the multiplication of neuronal cells and the formation of axial connections leading to insufficient development of the central neural system (CNS) (16) Brain CT and MRI findings showed enlargement of lateral cerebral ventricles and decreased volume of cerebral cortex mainly in the frontal and temporal lobes, as well as reduced hippocampal volume [20,21].

According to the neurodevelopmental disorder theory both genetic and environmental factors play a significant role in abnormal cerebral maturation [22].

In the pathophysiology of schizophrenia, it seems that immune functions are disrupted due to increased activation of the immune system by viral or bacterial infections, or immunogenic factors. The increased activity of the immune system may lead to excess release of inflammatory cytokines, which in turn may affect CNS growth of the fetus [23,24]. In schizophrenia abnormal interactions are observed between astroglia and microglia. The pro-inflammatory cytokines of microglia stimulate the production of kynurenic acid in astrocytes.

These observations identify cell populations associated with the risk of schizophrenia and should be investigated [25].

Various studies have shown higher levels of proinflammatory cytokines during pregnancy [26,27]. However, the increased cytokines levels during gestation, such as TNF- $\alpha$ , IL-1 $\beta$  and IL-6 may adversely affect the development of the fetal brain [28]. The cytokines IL-1 $\beta$ , IL-6 and, TNF- $\alpha$  may cross the placenta, whereas they may be synthesized by the mother, the placenta itself, or from the fetus. The levels of these maternal cytokines have been shown to be elevated in human pregnancies in which the offspring exhibit schizophrenia [29,30]. Most likely, in both prenatal and postnatal infections, the risk of schizophrenia is rather increased by the immune response (inflammatory cytokines, discs), instead of the specific pathogen [31,32].

There are indications of common genetic risk factors for both schizophrenia and depression. In addition to hereditary and environmental factors, genetic factors also seem to play an important role in the pathogenesis of emotional disorders [33]. The genome study of five psychiatric disorders such as schizophrenia, bipolar disorder, Attention Deficit Hyperactivity Disorder (ADHD) and Autism Spectrum Disorders (ASDs) showed that single nucleotide variants in the CACNA1C and CACNB2 genes, which encode calcium channel subunits, are highly associated with these diseases [34]. Several studies have identified a promising association with candidate genes, including COMT, DISC1, RGS4, PPP3CC, ZDHHC8, AKT1, neuregulin, dysbindin, G72 / G30, TRAR4, and alpha-7 nicotinic receptors. These genes are involved in the regulation of dopamine, contributing to the underlying cause of schizophrenia [35,36].

## 1.2 Environmental Factors

Several genetic, psychological, and environmental factors have been implicated at the manifestation of schizophrenia. Epidemiological studies as well as studies from monozygotic twins show that the factors affect the premature development of the brain [37].

**The season of childbirth:** In their studies Hare *et al* and Machon *et al* reported that children who were born towards the end of the winter or early spring, showed an increased incidence of schizophrenia [38].

These findings correspond with other studies which showed that childbirths during these months showed an increased rate of occurrence of schizophrenia by 5% -15%. This is happening probably due to an increase of incidence of viral infections during the winter and early spring period [39-41].

**Stress during pregnancy:** Situations creating intense stress during the pregnancy are associated with an increased risk of developing psychiatric disorders in the offspring due to the disturbance of the hypothalamic-pituitary-adrenal axis [42]. Maternal stress during the gestational period may cause placental vasoconstriction and delayed intrauterine growth of the fetus [43].

**Poor nutrition:** Various nutrient deficiency and especially of folic acid which is essential for cerebral development may lead to an increased risk of neurodevelopmental disorder [44].

**Place of birth:** The urban childbirth and upbringing of children are recognized as possible factors in the manifestation of adult schizophrenia later. Several mechanisms could explain this correlation, such as increased exposure to infectious agents in densely populated areas [45,46].

**Obstetric complications:** Various studies have shown that complications during gestation, or during labor and delivery may increase the risk of schizophrenia in offspring [47,48].

Most studies of the patients case-control showed a high rate of schizophrenia in people with complications during childbirth and especially in those with a genetic vulnerability [47,49-51]. This could be due to the hypoxia of the fetus which leads to irreversible injury of the hippocampus and of the amygdala [43,48].

Schizophrenia has also been linked to other complications of pregnancy such as low birth weight, [52] congenital malformations, small circumference of the head, suffocation during childbirth, uterine atony, perinatal bleeding, and ischemic injuries [53,54].

**Prenatal viral infections:** Viral infections during the 1<sup>st</sup> and the 2<sup>nd</sup> trimester of gestation can lead to neurodevelopmental disorders and possibly to the development of psychiatric illness during adolescence or adulthood [55,56]. In our review we focused on the impact of prenatal viral infections which, according to the available

literature, have been shown to have a connection with the onset of schizophrenia.

### 1.3 Influenza Virus

Infection with the influenza virus during pregnancy seems to be a risk factor for appearance of schizophrenia in offspring during adulthood. There are various studies showing that most people with schizophrenia are born during the winter months in late winter and early spring where flu outbreaks occur. There is much discussion as to whether direct offspring infection leads to neurodevelopmental disorders that ultimately lead to schizophrenia or whether this is due to the production of maternal cytokines in response to infection [39,57].

Mednick *et al.* examined the possibility whether the fetal viral infection is a causative agent of schizophrenia in adults. In their study, they included 1781 patients who were born during the flu A2 epidemic in 1957 in Helsinki and were diagnosed with schizophrenia. Their study showed an increased risk of schizophrenia among people whose second trimester of fetal life coincided with the flu epidemic, compared to those whose coincided the first or third trimester [58].

Another study by Limosin *et al.*, compared 973 schizophrenic patients with a corresponding number of non-schizophrenic patients and their non-schizophrenic brothers. They showed that there was an association between exposure to influenza virus and the manifestation of schizophrenia. The number of schizophrenic patients who exposed to the flu virus, especially during the second trimester of pregnancy, was much higher than that of controls [59].

As for England and Wales, O'Callaghan *et al.* reported an 88% increase in schizophrenic adults which were born during the period from February to mid-March. The study included 616 patients which were born between 1983 and 1988 and suffered from schizophrenia. They divided them in two groups: patients with a family history of psychiatric disorder and patients without a family history. The study showed a significant increase only in those who had no family history of any psychiatric disorder supporting the view that an environmental factor is responsible for the occurrence of this disorder [60].

Brown *et al.* used serological methods to document prenatal exposure to influenza and the

risk of schizophrenia. Their study included 64 patients who had been diagnosed with schizophrenia and 125 controls, all born during the period 1959-1966. Maternal serum samples were tested for influenza antibodies in order to estimate the prevalence in population during the same period. Serum samples came from pregnant women whose offspring developed schizophrenia compared with pregnant women whose offspring did not develop schizophrenia. The results of this study showed that the risk of schizophrenia increased in people exposed intrauterine to the flu virus during the first trimester of pregnancy compared to those in the second or third trimester [61].

Torrey *et.al.*, on the other hand, examined birth data in 43,778 people with schizophrenia and compared them to 10,496,686 births between 1950 and 1959 in ten US states, and concluded that there was no significant increase in the schizophrenic birth rate just before during or after the 1957 flu epidemic [62].

## 2. HERPEVIRIDAE FAMILY

The exposure to herpes viruses is a possible risk factor for development of schizophrenia. Members of the herpeviridae family associated with schizophrenia include herpes simplex viruses (HSV) 1 and 2, cytomegalovirus (CMV), Epstein-Barr virus (EBV), human herpes virus 6 (HHV-6) and the varicella-zoster virus (VZV). Herpes viruses are not totally eradicated after their initial infection but remain dormant incorporated in host genome and are reactivated during immunosuppression [63].

### 2.1 CMV

CMV is found in saliva, blood, urine, vaginal fluids, and breast milk of infected individuals. CMV infection is transmitted through close personal contact and from mother to fetus. It is more common to be found among individuals with low socioeconomic status [64]. Maternal infection during pregnancy or reactivation of the virus leads to congenital infection of the fetus. Congenital CMV infection is characterized by hepatosplenomegaly, jaundice, microcephaly, congenital mental retardation, hearing and vision loss, and cerebral calcifications [65,66]. CMV is the most common cause of damage to the fetal nervous system and, therefore, it is believed that there may be a causal link between congenital infection and schizophrenia [67]. Torrey *et al.* concluded that there is a positive causal

relationship between CMV and schizophrenia [68]. Similar results were presented by Dickerson *et al* who studied 323 individuals with schizophrenia in whom elevated CMV antibody titers were found [69]. Albrecht *et al.* also reported elevated CMV antibody titers in 19 of 60 schizophrenic patients in cerebrospinal fluid [70]. On the other hand, several studies showed a negative correlation between CMV infection and schizophrenia [71,72].

Serum antibody titers against CMV were also studied from patients with schizophrenia. Those studies showed no significant differences in serum anti-CMV antibodies between schizophrenic patients and controls [72-74]. Additionally, increased levels of antibodies were also found in CSF IgM against CMV, but not IgG without being able to determine whether it was a primary infection or reactivation of the virus [75].

In a study by Hoffmann *et al.* a possible reduction in brain volume in CMV-infected embryos has been investigated by the use of MRI. The study included 27 CMV-infected embryos and 52 uninfected embryos. Infected fetuses had a significantly lower brain volume than controls. This, combined with a genetic predisposition, may increase the risk of developing schizophrenia later in life [76].

### 2.2 HSV

HSV is transmitted through nasal or oral secretions and can cause mild or no symptoms. The most serious complication caused by the virus is encephalitis with severe morbidity which can lead to psychosis and cognitive impairment [77].

Several studies have shown an association between HSV-1 infection and cognitive impairment in schizophrenia. However, no association was found between HSV-2, HSV-6, CMV, EBV, and VZV infection [78-80].

Similarly, a study by Brown *et.al.* examined whether maternal exposure to HSV-2 was associated with the risk of schizophrenia in adulthood. 60 prenatal serum samples were examined and the authors found no correlation between prenatal exposure to HSV-2 and the risk of schizophrenia [81].

Mohagheghi *et.al.* studied IgM and IgG serum titers against HSV-1, HSV-2, and CMV in 45 patients with schizophrenia and 45 controls. The results showed that anti-CMV and anti-HSV1

and anti-HSV2 IgG antibodies were significantly increased in patients with schizophrenia compared to controls [82]. Mortensen *et al.* studied the possible correlation between maternal exposure to HSV-2 and the risk of schizophrenia in their adult offspring. They examined the blood serum of 602 schizophrenic patients and 602 controls. The study showed an increased risk of schizophrenia in the offspring of mothers positive for serum HSV-2 IgG [83]. In another study conducted by Thomas *et al.*, they investigated whether intrauterine exposure to HSV-1 is associated with an increased risk of developing schizophrenia and reduced cognitive function. The study included 171 patients with schizophrenia, 27 with schizophreniform disorders and 100 controls. All were evaluated for exposure to HSV-1 using serum HSV-1 antibody titers. The results did not show a significant relationship between exposure to HSV-1 and the risk of schizophrenia. On the other hand, significant cognitive impairment was found among patients exposed to HSV-1 compared to non-exposed individuals [79].

### 2.3 EBV

Another virus that belongs to the herpeviridae family is EBV which can cause persistent infection. In the study by Dickerson *et al.*, serum IgG EBV antibodies virus were measured in 432 people with schizophrenia and 311 controls. People with schizophrenia had elevated levels of antibodies to EBV viruses compared to the control population [84]. The same research group in a recent study shown that there is a significant association between decreased cognitive performance in patients with schizophrenia as measured by the MATRICS Consensus Cognitive Battery (MCCB) and the levels of antibodies to EBV (anti-EBNA-1 and anti-VCA) [85]. Khandaker et al. [86] shown a connection between EBV infection in childhood and an increased subsequent risk of psychotic experiences in adolescence.

### 3. RETROVIRUSES

Retroviruses are enveloped RNA viruses and exist as exogenous or endogenous entities. Exogenous retroviruses infect cells through a specific receptor for the virus, while endogenous retroviruses are present in the genome of host cells and are inherited through successive generations [87]. A typical retrovirus contains 3 genes, the order of which is gag-pol-env [88]. With the help of a reverse transcriptase, the

viruses can convert their RNA into DNA, and this retroviral DNA can be incorporated into the cell of the host cell. Retroviruses are able to induce immunodeficiency and neurological disorders such as multiple sclerosis and schizophrenia. Human endogenous retroviruses (HERVs), which belong to the human family of endogenous W-type retroviruses (HERV-W), have attracted the most attention. Several studies have since then investigated whether there is a causal link between HERV and schizophrenia, with the HERV-W family showing the greatest correlation indications [89].

In their study, Huang *et al.* collected blood samples from 58 individuals with schizophrenia and 38 controls. Using RT-PCR, they examined the RNA levels of the HERV pol gene in the blood. Retroviral pol genes were found to be elevated in 20 out of 58 subjects with schizophrenia. Additionally, the Western blot method showed elevated antibodies against ERV9 pol protein in the serum of the patients with schizophrenia and HERV+, but not in controls leading to the conclusion that the activation of certain retroviral proteins of the HERV-W family in some patients can lead to schizophrenia [90]. In a different study, the same group tried to detect genes with mRNA sequencing similar to the human endogenous human gene HERV-W env in the plasma of 118 subjects with recent onset of schizophrenia and 106 physiological controls/The authors detected the gene in 42 out of 118 individuals with schizophrenia but none was found in the participants of the control group of the study. Using human glioma cells U 251, they found that HERV-Wenv overexpression regulates the neurotrophic factor derived from the brain (BDNF), a gene associated with schizophrenia, the N2-receptor receptor type of neurotrophic tyrosine kinase type 2 (NTRK2, also known as TrkB) and dopamine receptor D3 and increased cAMP phosphorylation. Gene tests showed that HERV-W env activates BDNF production in human U251 glial cells. The HERV transcriptional activation is associated with the development of schizophrenia in some patients, and HERV-W env regulates the expression of schizophrenia-related genes [91].

Retroviruses incorporate into DNA and can infect brain cells, and causing damage to brain function, combined with genetic factors. Yolken *et al.* investigated the expression of RNA using the RT-PCR method in the frontal cortex in the postmortem brains of four individuals with

schizophrenia, four subjects with bipolar disorder, and six subjects control group. HERV-W expression was increased significantly in subjects with schizophrenia compared with control participants [80]. Perron *et al.* in their study quantified HERV-W gag and envelope (env) proteins in the serum of 49 subjects with schizophrenia and 49 healthy individuals, with a dedicated immunoassay set-up with specific monoclonal antibodies to either antigen. A positive antigenemia for envelope protein was found in 47% of patients, while 49% of patients had a positive antigen for gag compared with 3% for env and 4% for gag in the healthy subjects. A significant correlation was also found between gag or env antigenemia and C-reactive protein, suggesting a pathophysiology caused by inflammation [92]. Karlsson *et al.* identified similar nucleotide sequences in retroviral pol genes in CSF in 10 of 35 patients (29%) diagnosed with schizophrenia and schizophreniform disorders with recent onset, as well as in CSF of one out of 20 subjects with chronic. These retroviral sequences which are mainly related to the HERV-W family were not detected in any of the 32 participants in the study. Moreover, in the same study, brain tissue was taken postmortem from the frontal cortex of 5 subjects with schizophrenia. Additionally, similar frontal cortex tissue was obtained from six individuals with no history of psychiatric illness. Increased RNA transcription was found for retrovirus HERV-W in the brain tissue of individuals with schizophrenia compared with the healthy ones [93]. Yao *et al.* reported elevated transcription levels of HERV-W gag in mononuclear blood cells obtained from 30 patients with psychosis, including schizophrenia, schizophreniform disorder, and schizoaffective disorder compared to 26 healthy subjects in the control group. Increased levels of HERV-W gag were observed, but no difference was found between HERV-W env transcription levels [94].

#### 4. BORNA VIRUS

Bornavirus (BDV) is a negative ss RNA virus that belongs to the Bornaviridae family, and naturally infects horses, sheep, poultry, and cattle. It is a neurotropic virus that infects the brain and can cause nerve cell degeneration in the chronic phase of the disease [95]. BDV infection of CNS in animals can cause sporadic neurological diseases such as encephalitis, meningitis, and various abnormalities in movement and behavior, and sometimes with psychological manifestations similar to schizophrenia [96].

Several studies have focused on whether BDV disease can cause psychiatric disorders in humans such as schizophrenia and depression. Most studies have not shown that BDV infection causes psychiatric manifestations in humans and especially the manifestation of schizophrenia.

Yong *et al.* showed in their study that BDV infection may in some cases be associated with schizophrenia [90]. They measured BDV p24 antibodies in the sera of 116 schizophrenic patients and control subjects, and the results showed that infection is possibly associated with schizophrenia [97].

The study of Iwata *et al.*, however, showed opposite results. BDV p24 RNA was measured in monocytes of the peripheral blood (PBMCs) of psychiatric patients, 49 of whom had mood disorders, 77 schizophrenia, and 84 control individuals. The results did not show a greater prevalence of BDV p24 RNA in patients with psychiatric illness compared to the control group. Similar results were found by the study of Selten *et al.* who showed that the role of BDV infection in the pathogenesis of schizophrenia is unlikely [98,99]. The same negative correlation was shown by a study conducted in Iran, where the possibility of BDV infection-causing psychiatric disorders was considered. They examined samples of peripheral blood for the detection of BDV RNA nucleoprotein P40 from 120 patients (60 with bipolar disorder (BD) and 60 with schizophrenia) and 75 controls. The results showed no link between BDV infection and the pathogenesis of these psychiatric disorders [100]. In another study, which included 40 individuals with the first onset of schizophrenia and 40 controls, the researchers examined whether there was a correlation between BDV infection and schizophrenia measuring BDV RNA in PBMCs isolated from patients with the first onset of schizophrenia. The results did not show a causal link between the Bornavirus and schizophrenia [101].

##### 4.1 Other Viral Infections

A small study by Severance *et al.* in 106 patients with early onset of psychotic symptoms and 196 controls showed a higher seropositivity for HKU1, NL63 and OC43 coronaviruses [102]. A possible role of parvovirus B19 infection has also been implicated in some studies, but the results should be further investigated in larger studies [103,104].

**Table 1. Antiviral drugs used during pregnancy**

Antiviral drug	Target	References
<i>Acyclovir</i>	HSV	[111]
<i>Valacyclovir</i>	HSV, CMV	[111]
<i>CMV hyperimmune globulin</i>	CMV	[111]
<i>Lamivudine (3TC)</i>	HBV, HIV-1	[111]
<i>Tenofovir disoproxil fumarate (TDF)</i>	HBV, HIV-1	[111]
<i>Tenofovir Alafenamide (TAF)</i>	HIV-1	[112]
<i>Abacavir (ABC)</i>	HIV-1	[112]
<i>Emtricitabine (FTC)</i>	HIV-1	[112]
<i>Zidovudine (AZT)</i>	HIV-1	[112]
<i>Efavirenz (EFV)</i>	HIV-1	[112]
<i>Nevirapine (NVP)</i>	HIV-1	[112]
<i>Doravirine (DOR)</i>	HIV-1	[112]
<i>Rilpivirine (RPV)</i>	HIV-1	[112]
<i>Atazanavir (ATV)</i>	HIV-1	[112]
<i>Darunavir (DRV)</i>	HIV-1	[112]
<i>Lopinavir/Ritonavir (LPV/r)</i>	HIV-1	[112]
<i>Fostemsavir (FTR)</i>	HIV-1	[112]
<i>Ibalizumab-uiyk (IBA)</i>	HIV-1	[112]
<i>Maraviroc</i>	HIV-1	[112]
<i>Bictegravir/FTC/TAF</i>	HIV-1	[112]
<i>Cabotegravir</i>	HIV-1	[112]
<i>Elvitegravir</i>	HIV-1	[112]
<i>Dolutegravir</i>	HIV-1	[112]
<i>Raltegravir</i>	HIV-1	[112]
<i>Cobisistat</i>	HIV-1	[112]
<i>Ritonavir</i>	HIV-1	[112]
<i>DAAAs (anti-NS5B, anti-NS5A)</i>	HCV	[111]

Eagles in an early paper in 1992 suggested a possible connection between poliovirus infection and schizophrenia, but his hypothesis was never tested [105,106]. A possible role of Zika virus infection has been proposed in the pathogenesis of schizophrenia but there are no data supporting this hypothesis [107,108].

Given the current pandemic of SARS-CoV-2 some researchers claim that COVID-19 infection during pregnancy may have a neurodevelopmental impact on the offspring. It is well known that SARS-CoV-2 triggers a cascade of inflammatory cytokines and chemokines, many of which have been implicated in brain development such as IL1 $\beta$ , IL-2, IL-4, IL-6, IL-8 and IL-10 as well as IFN- $\alpha$ , IFN- $\gamma$  and TNF- $\alpha$  [109]. Studies will be needed in order to define whether COVID-19 infection will lead to an increase of schizophrenia prevalence to the offsprings [110].

#### 4.2 Treatment of Antiviral Infections during Pregnancy

Treatment of antiviral infections during pregnancy aim to decrease symptoms of the

mother, ameliorate maternal disease and avoid or minimize vertical transmission and congenital infection. There are several drugs considered safe during pregnancy as shown at Table 1.

## 5. CONCLUSIONS

Results from various studies show a positive association between viral infections during pregnancy and subsequent development of schizophrenia in offspring. Prenatal viral infections can change the normal development of the unborn baby's nervous system through the activation of the maternal immune system. Prenatal maternal infections appear to increase the risk of developing schizophrenia in the offspring, and this is due to the body's immune response to the infection (increased levels of inflammatory cytokines, antibodies), and not in exposure to specific viral or bacterial pathogens.

## CONSENT

It is not applicable.

## ETHICAL APPROVAL

It is not applicable.



## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

- Lakhan SE, Vieira KF. Schizophrenia pathophysiology: Are we any closer to a complete model? *Annals of general psychiatry*. 2009;8:12.
- Riedmuller R, Muller S. Ethical Implications of the Mild Encephalitis Hypothesis of Schizophrenia. *Frontiers in psychiatry*. 2017;8:38.
- Vahia VN. Diagnostic and statistical manual of mental disorders 5: A quick glance. *Indian Journal of Psychiatry*. 2013; 55(3):220-3.
- Gaebel W, Stricker J, Kerst A. Changes from ICD-10 to ICD-11 and future directions in psychiatric classification. *Dialogues in clinical neuroscience*. 2020; 22(1):7-15.
- Wong AH, Van Tol HH. Schizophrenia: from phenomenology to neurobiology. *Neuroscience and biobehavioral reviews*. 2003;27(3):269-306.
- Boog G. Obstetrical complications and subsequent schizophrenia in adolescent and young adult offsprings: is there a relationship? *European journal of obstetrics, gynecology, and reproductive biology*. 2004;114(2):130-6.
- Chung Y, Cannon TD. Brain imaging during the transition from psychosis prodrome to schizophrenia. *The Journal of nervous and mental disease*. 2015;203 (5):336-41.
- Marcelis M, Navarro-Mateu F, Murray R, Selten JP, Van Os J. Urbanization and psychosis: a study of 1942-1978 birth cohorts in The Netherlands. *Psychological medicine*. 1998;28(4):871-9.
- Murray RM, Lewis SW. Is schizophrenia a neurodevelopmental disorder? *British medical journal*. 1987;295(6600):681-2.
- Livsannyam E, Jain MS, Pormento MKL, Siddiqui H, Balagtas ARA, Emuze BO, et al. Neurobiology of Schizophrenia: A Comprehensive Review. *Cureus*. 2022; 14(4):e23959.
- Eggers AE. A serotonin hypothesis of schizophrenia. *Medical hypotheses*. 2013; 80(6):791-4.
- Guan J, Cai JJ, Ji G, Sham PC. Commonality in dysregulated expression of gene sets in cortical brains of individuals with autism, schizophrenia, and bipolar disorder. *Translational psychiatry*. 2019; 9(1):152.
- Hensler JG, Artigas F, Bortolozzi A, Daws LC, De Deurwaerdere P, Milan L, et al. Catecholamine/Serotonin interactions: systems thinking for brain function and disease. *Advances in pharmacology*. 2013; 68:167-97.
- Kesby JP, Eyles DW, McGrath JJ, Scott JG. Dopamine, psychosis and schizophrenia: the widening gap between basic and clinical neuroscience. *Translational psychiatry*. 2018;8(1):30.
- Bloom FE. Advancing a neurodevelopmental origin for schizophrenia. *Archives of general psychiatry*. 1993;50(3):224-7.
- Keshavan MS. Development, disease and degeneration in schizophrenia: a unitary pathophysiological model. *Journal of psychiatric research*. 1999;33(6):513-21.
- Pantelis C, Maruff P. The cognitive neuropsychiatric approach to investigating the neurobiology of schizophrenia and other disorders. *Journal of psychosomatic research*. 2002;53(2):655-64.
- Walker EF, Diforio D, Baum K. Developmental neuropathology and the precursors of schizophrenia. *Acta psychiatrica Scandinavica Supplementum*. 1999;395:12-9.
- Zaidel DW, Esiri MM, Harrison PJ. The hippocampus in schizophrenia: lateralized increase in neuronal density and altered cytoarchitectural asymmetry. *Psychological medicine*. 1997;27(3):703-13.
- Sanfilippo M, Lafargue T, Rusinek H, Arena L, Loneragan C, Lautin A, et al. Volumetric measure of the frontal and temporal lobe regions in schizophrenia: relationship to negative symptoms. *Archives of general psychiatry*. 2000;57(5):471-80.
- Thaker GK, Carpenter WT, Jr. Advances in schizophrenia. *Nature medicine*. 2001;7(6): 667-71.
- Tsuang M. Schizophrenia: genes and environment. *Biological psychiatry*. 2000; 47(3):210-20.
- Horvath S, Mirnics K. Immune system disturbances in schizophrenia. *Biological psychiatry*. 2014;75(4):316-23.
- Khandaker GM, Cousins L, Deakin J, Lennox BR, Yolken R, Jones PB. Inflammation and immunity in schizophrenia: implications for

- pathophysiology and treatment. *The lancet Psychiatry*. 2015;2(3):258-70.
25. Abashkin DA, Kurishev AO, Karpov DS, Golimbet VE. Cellular Models in Schizophrenia Research. *International journal of molecular sciences*. 2021;22(16).
  26. Buka SL, Tsuang MT, Torrey EF, Klebanoff MA, Bernstein D, Yolken RH. Maternal infections and subsequent psychosis among offspring. *Archives of general psychiatry*. 2001;58(11):1032-7.
  27. Buka SL, Tsuang MT, Torrey EF, Klebanoff MA, Wagner RL, Yolken RH. Maternal cytokine levels during pregnancy and adult psychosis. *Brain, behavior, and immunity*. 2001;15(4):411-20.
  28. Dammann O, Leviton A. Infection remote from the brain, neonatal white matter damage, and cerebral palsy in the preterm infant. *Seminars in pediatric neurology*. 1998;5(3):190-201.
  29. Mattei D, Djodari-Irani A, Hadar R, Pelz A, de Cossio LF, Goetz T, et al. Minocycline rescues decrease in neurogenesis, increase in microglia cytokines and deficits in sensorimotor gating in an animal model of schizophrenia. *Brain, behavior, and immunity*. 2014;38:175-84.
  30. Rosenblat JD, Cha DS, Mansur RB, McIntyre RS. Inflamed moods: a review of the interactions between inflammation and mood disorders. *Progress in neuro-psychopharmacology & biological psychiatry*. 2014;53:23-34.
  31. Benros ME, Mortensen PB, Eaton WW. Autoimmune diseases and infections as risk factors for schizophrenia. *Annals of the New York Academy of Sciences*. 2012; 1262:56-66.
  32. Estes ML, McAllister AK. Maternal immune activation: Implications for neuropsychiatric disorders. *Science*. 2016;353(6301):772-7.
  33. Kalueff A, Demin K, Volgin A. Genomic mechanisms of anxiety and depression pathogenesis in experimental models. *VM Bektherev Rev Psych Med Psychol*. 2019;4(1):48-50.
  34. Cross-Disorder Group of the Psychiatric Genomics C. Identification of risk loci with shared effects on five major psychiatric disorders: a genome-wide analysis. *Lancet*. 2013;381(9875):1371-9.
  35. Cannon TD. The inheritance of intermediate phenotypes for schizophrenia. *Current opinion in psychiatry*. 2005;18(2): 135-40.
  36. Riley B, Kendler KS. Molecular genetic studies of schizophrenia. *European journal of human genetics: EJHG*. 2006;14(6):669-80.
  37. Bauman MD, Lesh TA, Rowland DJ, Schumann CM, Smucny J, Kukis DL, et al. Preliminary evidence of increased striatal dopamine in a nonhuman primate model of maternal immune activation. *Translational psychiatry*. 2019;9(1):135.
  38. Hare EH, Price JS, Slater E. Schizophrenia and season of birth. *The British journal of psychiatry : The Journal of Mental Science*. 1972;120(554):124-5.
  39. Pallast EG, Jongbloet PH, Straatman HM, Zielhuis GA. Excess seasonality of births among patients with schizophrenia and seasonal ovopathy. *Schizophrenia bulletin*. 1994;20(2):269-76.
  40. Susser EB, Brown A, Matte TD. Prenatal factors and adult mental and physical health. *Canadian Journal of Psychiatry Revue canadienne de psychiatrie*. 1999; 44(4):326-34.
  41. Torrey EF, Miller J, Rawlings R, Yolken RH. Seasonality of births in schizophrenia and bipolar disorder: A review of the literature. *Schizophrenia research*. 1997;28 (1):1-38.
  42. Mueller BR, Bale TL. Sex-specific programming of offspring emotionality after stress early in pregnancy. *The Journal of neuroscience: The Official Journal of the Society for Neuroscience*. 2008;28(36): 9055-65.
  43. DeLisi LE, Goldin LR, Gershon ES. Studies of biological factors associated with the inheritance of schizophrenia: a selective review. *Journal of psychiatric research*. 1987;21(4):507-13.
  44. Brown AS, Susser ES. Prenatal nutritional deficiency and risk of adult schizophrenia. *Schizophrenia bulletin*. 2008;34(6):1054-63.
  45. Pedersen CB, Mortensen PB. Family history, place and season of birth as risk factors for schizophrenia in Denmark: a replication and reanalysis. *The British journal of psychiatry: The Journal of Mental Science*. 2001;179:46-52.
  46. Sham PC, MacLean CJ, Kendler KS. Risk of schizophrenia and age difference with older siblings. Evidence for a maternal viral infection hypothesis? *The British journal of psychiatry: The Journal of Mental Science*. 1993;163:627-33.

47. DeLisi LE, Weber RJ, Pert CB. Are there antibodies against brain in sera from schizophrenic patients? Review and prospectus. *Biological psychiatry*. 1985; 20(1):110-5.
48. McNeil TF. Perinatal risk factors and schizophrenia: selective review and methodological concerns. *Epidemiologic reviews*. 1995;17(1):107-12.
49. Jacobsen B, Kinney D. Perinatal complications in adopted and non-adopted schizophrenics and their controls: Preliminary results. *Acta psychiatrica Scandinavica Supplementum*. 1980;62 (S285):337-46.
50. Owen MJ, Lewis SW, Murray RM. Obstetric complications and schizophrenia: a computed tomographic study. *Psychological medicine*. 1988;18(2):331-9.
51. Parnas J, Schulsinger F, Teasdale TW, Schulsinger H, Feldman PM, Mednick SA. Perinatal complications and clinical outcome within the schizophrenia spectrum. *The British Journal of Psychiatry: The Journal of Mental Science*. 1982;140:416-20.
52. Woerner MG, Pollack M, Klein DF. Birth weight and length in schizophrenics personality disorders and their siblings. *The British journal of psychiatry: The Journal of Mental Science*. 1971;118 (545):461-4.
53. Gilmore J, Murray R. Prenatal and perinatal factors. *The American psychiatric publishing textbook of schizoprhenia*. Washington: American Psychiatric Publishing Inc. 2006;55-67.
54. Stilo SA, Murray RM. The epidemiology of schizophrenia: replacing dogma with knowledge. *Dialogues in clinical neuroscience*. 2010;12(3):305-15.
55. Fatemi SH, Folsom TD. The neurodevelopmental hypothesis of schizophrenia, revisited. *Schizophrenia bulletin*. 2009;35(3):528-48.
56. Wright P, Takei N, Rifkin L, Murray RM. Maternal influenza, obstetric complications, and schizophrenia. *The American journal of psychiatry*. 1995;152(12):1714-20.
57. Watson CG, Kucala T, Tilleskjoer C, Jacobs L. Schizophrenic birth seasonality in relation to the incidence of infectious diseases and temperature extremes. *Archives of general psychiatry*. 1984; 41(1):85-90.
58. Mednick SA, Machon RA, Huttunen MO, Bonett D. Adult schizophrenia following prenatal exposure to an influenza epidemic. *Archives of general psychiatry*. 1988;45(2):189-92.
59. Limosin F, Rouillon F, Payan C, Cohen JM, Strub N. Prenatal exposure to influenza as a risk factor for adult schizophrenia. *Acta psychiatrica Scandinavica*. 2003;107(5):331-5.
60. O'Callaghan E, Gibson T, Colohan HA, Walshe D, Buckley P, Larkin C, et al. Season of birth in schizophrenia. Evidence for confinement of an excess of winter births to patients without a family history of mental disorder. *The British journal of psychiatry : the journal of mental science*. 1991;158:764-9.
61. Brown AS, Begg MD, Gravenstein S, Schaefer CA, Wyatt RJ, Bresnahan M, et al. Serologic evidence of prenatal influenza in the etiology of schizophrenia. *Archives of general psychiatry*. 2004;61(8):774-80.
62. Torrey EF, Bowler AE, Rawlings R. An influenza epidemic and the seasonality of schizophrenic births. *Psychiatry and biological factors: Springer*. 1991;109-16.
63. Bray PF, Bale JF, Anderson RE, Kern ER. Progressive neurological disease associated with chronic cytomegalovirus infection. *Annals of neurology*. 1981;9(5): 499-502.
64. Hanshaw JB. Congenital cytomegalovirus infection: a fifteen year perspective. *The Journal of infectious diseases*. 1971;123 (5):555-61.
65. Bhumbra NA, Nankervis GA. Cytomegalovirus infection. Overview and new developments. *Postgraduate medicine*. 1983;73(4):62-9.
66. Boppana SB, Rivera LB, Fowler KB, Mach M, Britt WJ. Intrauterine transmission of cytomegalovirus to infants of women with preconceptional immunity. *The New England Journal of Medicine*. 2001;344 (18):1366-71.
67. Torrey EF, Leweke MF, Schwarz MJ, Mueller N, Bachmann S, Schroeder J, et al. Cytomegalovirus and schizophrenia. *CNS Drugs*. 2006;11:879-85.
68. Torrey EF, Yolken RH, Albrecht P. Cytomegalovirus as a possible etiological agent in schizophrenia. *Adv Biol Psych*. 1983;12:150-60.
69. Dickerson F, Kirkpatrick B, Boronow J, Stallings C, Origoni A, Yolken R. Deficit schizophrenia: association with serum antibodies to cytomegalovirus.

- Schizophrenia bulletin. 2006;32(2):396-400.
70. Albrecht P, Torrey EF, Boone E, Hicks JT, Daniel N. Raised cytomegalovirus-antibody level in cerebrospinal fluid of schizophrenic patients. *Lancet*. 1980;2(8198):769-72.
  71. Gotlieb-Stematsky T, Zonis J, Arlazoroff A, Mozes T, Sigal M, Szekely AG. Antibodies to Epstein-Barr virus, herpes simplex type 1, cytomegalovirus and measles virus in psychiatric patients. *Archives of virology*. 1981;67(4):333-9.
  72. Lycke E, Norrby R, Roos BE. A serological study on mentally ill patients with particular reference to the prevalence of herpes virus infections. *The British journal of psychiatry: The Journal of Mental Science*. 1974; 124(0):273-9.
  73. Delisi LE, Smith SB, Hamovit JR, Maxwell ME, Goldin LR, Dingman CW, et al. Herpes simplex virus, cytomegalovirus and Epstein-Barr virus antibody titres in sera from schizophrenic patients. *Psychological medicine*. 1986;16(4):757-63.
  74. Krech U. Complement-fixing antibodies against cytomegalovirus in different parts of the world. *Bulletin of the World Health Organization*. 1973;49(1):103-6.
  75. Kaufmann CA, Weinberger DR, Yolken RH, Torrey EF, Pofkin SG. Viruses and schizophrenia. *Lancet*. 1983;2(8359):1136-7.
  76. Hoffmann C, Grossman R, Bokov I, Lipitz S, Biegon A. Effect of cytomegalovirus infection on temporal lobe development in utero: quantitative MRI studies. *European neuropsychopharmacology : the journal of the European College of Neuropsychopharmacology*. 2010;20(12):848-54.
  77. Steiner I, Kennedy PG, Pachner AR. The neurotropic herpes viruses: herpes simplex and varicella-zoster. *The Lancet Neurology*. 2007;6(11):1015-28.
  78. Dickerson FB, Boronow JJ, Stallings C, Orioni AE, Ruslanova I, Yolken RH. Association of serum antibodies to herpes simplex virus 1 with cognitive deficits in individuals with schizophrenia. *Archives of general psychiatry*. 2003;60(5):466-72.
  79. Thomas P, Bhatia T, Gauba D, Wood J, Long C, Prasad K, et al. Exposure to herpes simplex virus, type 1 and reduced cognitive function. *Journal of psychiatric research*. 2013;47(11):1680-5.
  80. Yolken RH, Torrey EF, Lieberman JA, Yang S, Dickerson FB. Serological evidence of exposure to Herpes Simplex Virus type 1 is associated with cognitive deficits in the CATIE schizophrenia sample. *Schizophrenia research*. 2011;128 (1-3):61-5.
  81. Brown AS, Schaefer CA, Quesenberry CP, Jr., Shen L, Susser ES. No evidence of relation between maternal exposure to herpes simplex virus type 2 and risk of schizophrenia? *The American journal of psychiatry*. 2006;163(12):2178-80.
  82. Mohagheghi M, Eftekharian MM, Taheri M, Alikhani MY. Determining the IgM and IgG antibodies titer against HSV1, HSV2 and CMV in the serum of schizophrenia patients. *Human antibodies*. 2018;26(2): 87-93.
  83. Mortensen PB, Pedersen CB, Hougaard DM, Norgaard-Petersen B, Mors O, Borglum AD, et al. A Danish National Birth Cohort study of maternal HSV-2 antibodies as a risk factor for schizophrenia in their offspring. *Schizophrenia research*. 2010; 122(1-3):257-63.
  84. Dickerson F, Jones-Brando L, Ford G, Genovese G, Stallings C, Orioni A, et al. Schizophrenia is Associated With an Aberrant Immune Response to Epstein-Barr Virus. *Schizophrenia bulletin*. 2019; 45(5):1112-9.
  85. Dickerson F, Katsafanas E, Orioni A, Squire A, Khushalani S, Newman T, et al. Exposure to Epstein Barr virus and cognitive functioning in individuals with schizophrenia. *Schizophrenia research*. 2021;228:193-7.
  86. Khandaker GM, Stochl J, Zammit S, Lewis G, Jones PB. Childhood Epstein-Barr Virus infection and subsequent risk of psychotic experiences in adolescence: a population-based prospective serological study. *Schizophrenia research*. 2014;158(1-3):19-24.
  87. Christensen T. HERVs in neuropathogenesis. *Journal of neuroimmune pharmacology: The official journal of the Society on NeuroImmune Pharmacology*. 2010;5(3):326-35.
  88. Gurling HM. Testing the retrovirus hypothesis of manic depression and schizophrenia with molecular genetic techniques. *Journal of the Royal Society of Medicine*. 1988;81(6):332-4.
  89. Aftab A, Shah AA, Hashmi AM. Pathophysiological Role of HERV-W in Schizophrenia. *The Journal of neuropsychiatry and clinical neurosciences*. 2016;28(1):17-25.

90. Huang WJ, Liu ZC, Wei W, Wang GH, Wu JG, Zhu F. Human endogenous retroviral pol RNA and protein detected and identified in the blood of individuals with schizophrenia. *Schizophrenia research*. 2006;83(2-3):193-9.
91. Huang W, Li S, Hu Y, Yu H, Luo F, Zhang Q, et al. Implication of the env gene of the human endogenous retrovirus W family in the expression of BDNF and DRD3 and development of recent-onset schizophrenia. *Schizophrenia bulletin*. 2011;37(5):988-1000.
92. Perron H, Mekaoui L, Bernard C, Veas F, Stefanis I, Leboyer M. Endogenous retrovirus type W GAG and envelope protein antigenemia in serum of schizophrenic patients. *Biological psychiatry*. 2008;64(12):1019-23.
93. Karlsson H, Bachmann S, Schroder J, McArthur J, Torrey EF, Yolken RH. Retroviral RNA identified in the cerebrospinal fluids and brains of individuals with schizophrenia. *Proceedings of the National Academy of Sciences of the United States of America*. 2001;98(8):4634-9.
94. Yao Y, Schroder J, Nellaker C, Bottmer C, Bachmann S, Yolken RH, et al. Elevated levels of human endogenous retrovirus-W transcripts in blood cells from patients with first episode schizophrenia. *Genes, brain, and behavior*. 2008;7(1):103-12.
95. Gosztonyi G, Ludwig H. Borna disease - neuropathology and pathogenesis. *Borna disease*: Springer. 1995;39-73.
96. Bilzer T, Planz O, Lipkin WI, Stitz L. Presence of CD4+ and CD8+ T cells and expression of MHC class I and MHC class II antigen in horses with Borna disease virus-induced encephalitis. *Brain pathology*. 1995;5(3):223-30.
97. Yang AY, Zhang FM, Li JH, Li GM, Ma PL, Gu HX, et al. [Detection of Borna disease virus-p24 specific antibody in the sera of schizophrenic patients of China by means of Western-blot]. *Zhonghua shi yan he lin chuang bing du xue za zhi = Zhonghua shiyan he linchuan bingduxue zazhi = Chinese journal of experimental and clinical virology*. 2003;17(1):85-7.
98. Iwata Y, Takahashi K, Peng X, Fukuda K, Ohno K, Ogawa T, et al. Detection and sequence analysis of borna disease virus p24 RNA from peripheral blood mononuclear cells of patients with mood disorders or schizophrenia and of blood donors. *Journal of virology*. 1998;72(12):10044-9.
99. Seltzer JP, van Vliet K, Pleyte W, Herzog S, Hoek HW, van Loon AM. Borna disease virus and schizophrenia in Surinamese immigrants to the Netherlands. *Medical microbiology and immunology*. 2000;189(2):55-7.
100. Shatizadeh-Malekshahi S, Ahmadkhaniha HR, Kiani SJ, Nejati A, Janani L, Yavarian J. No molecular evidence of Borna disease virus among schizophrenia and bipolar disorder patients in Iran. *Iranian Journal of Microbiology*. 2017;9(2):112-8.
101. Soltani H, Mohammadzadeh S, Makvandi M, Pakseresht S, Samarbafe-Zadeh A. Detection of Borna Disease Virus (BDV) in Patients with First Episode of Schizophrenia. *Iranian Journal of Psychiatry*. 2016;11(4):257-61.
102. Severance EG, Dickerson FB, Viscidi RP, Bossis I, Stallings CR, Origoni AE, et al. Coronavirus immunoreactivity in individuals with a recent onset of psychotic symptoms. *Schizophrenia bulletin*. 2011;37(1):101-7.
103. Hammond CJ, Hobbs JA. Parvovirus B19 infection of brain: possible role of gender in determining mental illness and autoimmune thyroid disorders. *Medical hypotheses*. 2007;69(1):113-6.
104. Hobbs JA. Detection of adeno-associated virus 2 and parvovirus B19 in the human dorsolateral prefrontal cortex. *Journal of neurovirology*. 2006;12(3):190-9.
105. Eagles JM. Are polioviruses a cause of schizophrenia? *The British journal of psychiatry : The Journal of Mental Science*. 1992;160:598-600.
106. Squires RF. How a poliovirus might cause schizophrenia: a commentary on Eagles' hypothesis. *Neurochemical research*. 1997;22(5):647-56.
107. Job B, Wiwanitkit V. Zika Virus Outbreak, assisted reproduction patients and pregnancy. *JBRA assisted reproduction*. 2018;22(1):75.
108. Pierson TC, Diamond MS. The emergence of Zika virus and its new clinical syndromes. *Nature*. 2018;560(7720):573-81.
109. Reyes-Lagos JJ, Abarca-Castro EA, Echeverria JC, Mendieta-Zeron H, Vargas-Caraveo A, Pacheco-Lopez G. A Translational Perspective of Maternal Immune Activation by SARS-CoV-2 on the

- Potential Prenatal Origin of Neurodevelopmental Disorders: The Role of the Cholinergic Anti-inflammatory Pathway. *Frontiers in psychology*. 2021; 12:614451.
110. Massrali A, Adhya D, Srivastava DP, Baron-Cohen S, Kotter MR. Corrigendum: Virus-Induced Maternal Immune Activation as an Environmental Factor in the Etiology of Autism and Schizophrenia. *Frontiers in neuroscience*. 2022;16:943903.
111. Rogan SC, Beigi RH. Treatment of Viral Infections During Pregnancy. *Clinics in perinatology*. 2019;46(2):235-56.
112. Recommendations for the Use of Antiretroviral Drugs During Pregnancy and Interventions to Reduce Perinatal HIV Transmission in the United States; 2022.

© 2022 Kotsiri et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*

*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/87749>