



Effect of Boiling Times on Polyphenol, Flavonoid, Tannin, Vitamin C and β -carotene contents of African Asparagus (*Laccosperma secundiflorum*): Their Contribution to Overall Antioxidant Activity

**Kouassi Loukou Viviane-Sipo^{a*}, N'gbo Martin Luthère King^b,
Sinh Josi-Noelline^c, Disseka William Kwithony^a
and Faulet Meuwiah Betty-Ahonzo^a**

^a *Laboratory of Biocatalysis and Food Bioprocessing, University Nangui Abrogoua, Abidjan, Côte d'Ivoire.*

^b *Agriculture, Fishery Resources and Agro-Industry, Department of the University of San Pedro, Côte d'Ivoire.*

^c *National Institute of Youth and the Sports, Abidjan, Côte d'Ivoire.*

Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/IJBCRR/2021/v30i1030294

Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/79895>

Original Research Article

**Received 21 October 2021
Accepted 26 December 2021
Published 27 December 2021**

ABSTRACT

Aims: *Laccosperma secundiflorum* is a very important rattan species for certain populations in the Ivory Coast. They use the apical meristem in their food and the stems to make various items such as furniture, carpets and baskets etc. However, there is a gap in the study of its nutritional value and its nutritional potential before and after boiling.

Place and Duration of Study: Department of Food Science and Technology (UFR-STA), University nangui abrogoua, between December 2019 and November 2020.

Methodology: African asparagus was boiled for 5, 10 and 15 minutes respectively. The effect of boiling on the total polyphenols, flavonoids, tannins, vitamin C and β -carotene contents, and further on the antioxidant activity of the heart of rattan palm (African asparagus) was studied quantitatively at this different times. Standardized methods were adopted for phytochemical, antioxidant activity of this samples.

*Corresponding author: E-mail: viviane.kouassi3@gmail.com;

Results: Vitamin C content, which was estimated initially at 37.04 g/100 dry matter, decreased as the cooking time in water increased. Indeed, it is noted 40, 72, and 81% of loss respectively for the cooking times of 5, 10, and 15 min in the water. Also, these different heat treatments affected the β -carotene content of African asparagus with losses ranging from 76.80 to 84%. Polyphenols were also affected by cooking. The values obtained were 5104.80 mg GAE/100g for fresh African asparagus, 5284.09 mg GAE/100g for 5 min cooking (CE5), 5233.31 mg GAE/100g for 10 min cooking (CE10), and 3536.44 mg GAE/100g for 15 min cooking (CE15). On the other hand, the tannin content of fresh African asparagus was 1954.84 mg/100g DM. However, this rate decreased continuously with heating and was estimated at 1699.93 mg/100g DM, 1429.977548 mg/100g DM, and 1035.42 mg/100g DM respectively for CE5, CE10, and CE15 samples with losses of 13.04, 26.85 and 47.03% respectively. For flavonoids, the losses were varied from 7.41 and 19.92%. In our study, the scavenging activity was found to be highest in the fresh sample, followed by CE15 and CE5.

Conclusion: Finally, boiling reduces the levels of the different parameters studied, but the antioxidant activity of African asparagus increased at the end of the 15 min heat treatment. However, a cooking time of less than or equal to 5 min in water can be advantageous for the consumer.

Keywords: African asparagus; *Laccosperma secundiflorum*; antioxidant activity; boiling; palm heart.

1. INTRODUCTION

During this last decade, high consumption of fruits and vegetables associated with decreased risk of diseases such as cardiovascular pathologies, obesity, diabetes, neurodegenerative diseases, and cancer has been observed in numerous epidemiological studies [1]. This high consumption of fruits and vegetables is due to the richness of these foods in antioxidants (ascorbic acid, tocopherols, carotenoids, and polyphenols) which are molecules with preventive effects against these diseases because they participate in the neutralization of free radicals. These free radicals are permanently generated by our body or formed in response to environmental aggressions. Polyphenols are micronutrients that are particularly abundant in cereals, fruits, and vegetables [2]. Their interest lies in their antioxidant properties, especially their capacity to trap free radicals [3]. These plants also have multiple properties, among others antioxidants because of antioxidants compounds such as vitamins, carotenoids, phenolic compounds ...in their edible parts [4,5]. Several plants, fruits, and vegetables are consumed in Côte d'Ivoire in lean periods [6], and most of them are cooked before consumption, for example the African asparagus. However, culinary practices induce significant changes of chemical compositions, the concentration and bioavailability of bioactive compounds in these vegetables. Positive as well as negative effects have been reported based on differences in processing conditions and morphological and nutritional characteristics of

vegetable species [7]. Knowing how and why changes occur can help the consumer, the food processor, and even the chef to limit waste and therefore improve the nutritional quality of food. This study aimed to evaluate the effect of boiling water cooking on the total content of polyphenols, flavonoids, tannins, vitamin C and β -carotene, and thus on the antioxidant activity of African asparagus.

2. MATERIALS AND METHODS

2.1 Sampling

The African asparagus *Laccosperma secundiflorum* was collected in the Agneby Tiassa region, more precisely in the Sikensi area, 5°40'40" north latitude 4°34'33" south longitude. The African asparagus was transported in polypropylene plastic bags directly to the laboratory for analysis (Fig.1).

2.2 Methods

2.2.1 Heat treatment of samples

The boiling of African asparagus was done according to the method described by Randrianatoandro [8]. 1.5 kg of African asparagus cut into 5 cm "sticks" were immersed in 1 L of boiling water in a stainless steel container for 5, 10, and 15 min. The cooking solution was discarded and the boiled samples were cooled, drained at ambient temperature, and subjected to the same treatment used for raw samples.



Fig. 1. African asparagus

A: African asparagus enveloped in leaf sheaths, B: African asparagus with leaf sheaths removed

The cooked African asparagus samples were oven-dried (Biobase, China, Shandong) at 45°C for 48 h. They were then powdered using a Binatone-type blender (BLG-555, China, Hong Kong) and sieved using a sieve with a mesh size of 500 µm (AFNOR -NFX 11504. The obtained powders were stored in stomacher bags and kept at 4°C in a refrigerator (NASCO, DF2-28, China) for further analysis. The fresh sample was used as a control.

2.2.2 Biochemical analysis of the samples

2.2.2.1 Determination of Vitamin C (Vit C) content

Vitamin C contained in analysed samples was determined by titration using 2,6-dichlorophenol indophenol [9]. This method involves stabilizing vitamin C with metaphosphoric acid/acetic acid and then oxidizing it with 2,6-dichlorophenol indophenol (2,6-DCPIP) which is then reduced. The vitamin C content was obtained from this mathematical relationship:

$$\text{Vitamin C (mg/100g)} = \frac{(V_e - V_0) \times 20}{(V_s - V_0) \times 10} \times 100$$

V_e : Volume of 2,6-dichlorophenol-indophenol poured for the sample;

V_0 : Volume of 2,6-dichlorophenol-indophenol poured for the determination of metaphosphoric acid

V_s : Volume of 2,6-dichlorophenol-indophenol poured for the determination of the vitamin C stock solution

2.2.2.2 Determination of β-carotene content

The β-carotene were extracted and quantified by using a spectrophotometric method [10]. African asparagus samples (10g) were homogenized in ethanol (40 mL). The mixture was introduced into a separatory funnel containing 50 mL of hexane. The hexane phase was evaporated for 24 hours. Another 10 ml of hexane was added to this phase. After the rest of 24 hours, the optical density (OD) was read using a spectrophotometer (MS-V5100 visible spectrophotometer, Germany) at 450 nm against a blank solution. The standard solution was prepared with 10 mg of trans-β-carotene dissolved in pure hexane to obtain a 100 µg/mL solution.

2.2.2.3 Determination of total polyphenol content

Folin-ciocalteu method was used to determine the total phenols content [11]. To a test tubes were added 1 ml of methanolic extract and 1 ml of Folin-ciocalteu reagent. The tube was left to stand for 3 min and then 1 mL of sodium carbonate solution (20%, w/v) was added. The contents of the tube were made up to 10 mL with distilled water. After 30 min in the dark, the absorbance of gallic acid as standard and the methanolic extract was measured at 725 nm using a spectrophotometer (MS-V5100 visible

spectrophotometer, Germany) against a blank. A standard range was performed with a 1 mg/mL gallic acid solution.

2.2.2.4 Determination of total flavonoid content

Flavonoid quantification was carried out using aluminium chloride colorimetric method [12]. Into test tubes were successively added, 0.5 mL of methanolic extract, 0.5 mL of distilled water, 0.5 mL of aluminum chloride (10%), and 0.5 mL of potassium acetate (1 M). The final volume was made up of 2 mL of distilled water. The test tube was then incubated in the dark for 30 minutes. The absorbance of standard (quercetin) and the methanolic extract was measured spectrophotometrically at 415 nm. A calibration curve was made using a 0.1 mg/mL quercetin standard solution.

2.2.2.5 Determination of total tannin content

Tannins of samples were quantified using vanillin reagent method [13]. Into test tube, 1 mL of the methanolic extract was homogenised with 5 mL of vanillin reagent (0.1 mg/mL vanillin in 70% (v/v) sulphuric acid). The mixture was then incubated in the dark for 20 minutes at room temperature. The absorbance was measured at 500 nm using a spectrophotometer (MS-V5100 visible spectrophotometer, Germany) against a blank solution. A calibration range was performed using a 0.1 mg/mL tannic acid standard solution.

2.2.2.6 Measurement of antioxidant activity by DPPH radical

Antioxidant activity assay was carried out using the 2,2-diphenyl-1-picrylhydrazyl (DPPH) spectrophotometric method [14]. A solution of DPPH was freshly prepared (about 0.3 mM). The methanolic extract (2 mL) with varying concentrations (2-20 µg/mL) and DPPH solution (1 mL) were mixed in each test tube. The test tube was then incubated in the dark for 30 minutes at room temperature. The decrease in absorbance was measured at 517 nm using a spectrophotometer (MS-V5100 visible spectrophotometer, Germany). Vitamin C was used as the standard. The values of the inhibitory concentrations (IC50) of the different extracts were obtained by projection from the graph of percentage inhibition versus extract concentrations and are expressed in mg/ml. The percentage inhibition of radicals was calculated using the following formula:

$$AA (\%) = \frac{[DO_c - (DO_e - DO_b)] \times 100}{DO_c}$$

AA: antioxidant activity

DOC: absorbance of control tube (1 mL DPPH + 2 mL methanol)

DOe: absorbance of test tube (2 mL methanol extract + 1 mL DPPH)

DOB: absorbance of blank tube (1 mL methanol + 2 mL methanolic extract)

2.2.3 Statistical analysis of the results

The statistical analysis was applied to the data obtained during the biochemical evaluations. All tests relating to the different analyses were carried out in triplicate and the numerical values obtained were expressed as the arithmetic mean affected by the standard deviation. The one-factor ANOVA variance analysis was performed on all the results obtained to determine the existence of significant differences between the averages calculated according to the DUNCAN test using the STATISTICA software version 7.1. The graphs were built using Excel software.

3. RESULTS AND DISCUSSION

3.1 Results

3.1.1 Vitamin C and β-carotene

Ascorbic acid and β-carotene contents are shown in Figs. 2A and 2B. A decrease in the contents of these two parameters was observed with increasing cooking time. As for vitamin C, the losses in content oscillate between 40 and 81%. As for β-carotene, it was varied from 76.80 and 84%.

3.1.2 Phytochemical composition of boiled African asparagus

The obtained results for the polyphenol content were 5104.80, 5284.09, 5233.31, and 3536.44 mg GAE/100gMS respectively for the FRESH, CE5, CE10, and CE15 samples (Fig. 3A). No significant differences ($p > 0.05$) was noted between the means of the cooked samples at 5 and 10 min. However, these means were found to be statistically different from those of the fresh African asparagus and the CE15 sample. Furthermore, a slight increase in polyphenol content of 3.51% and 2.51% for CE5 and CE10, respectively. Then, a 30.72% drop in the averages was recorded at the 15th minute of cooking.




Fig. 2. Vitamin C (A) and β-carotene (B) content of raw and water-cooked African asparagus (EC) at different times (5, 10 and 15 min)

EC5 : cooking 5 min in water ; EC10 : cooking 10 min in water ; EC15 : cooking 15 min in water

The tannin contents of African asparagus during cooking in water were 1954.84, 1699.93, 1429.98, and 1035.42 mgTA/100gMS for samples FRESH, CE5, CE10, and CE15, respectively (Fig. 3C). Cooking with water results in a decrease in the tannin content of the samples with losses of 13.04, 26.85, and 47.03%, respectively for CE5, CE10, and CE15. Analysis of variance shows a significant difference ($p < 0.05$) between the four samples.

Flavonoid contents ranged from 49.43 mg QE/100g (FRESH) to 39.58 mg QE/100g (CE15) during cooking. At 5 min, 10 min, and 15 min of cooking in water, losses of 7.41 %, 12.15 %, and 19.92 %, respectively were observed. Also, a significant difference ($p < 0.05$) was observed between fresh and cooked samples (Fig. 3B).

3.1.3 Antioxidant activity of water-cooked boiling African asparagus

The percentages of free radical scavenging activity are presented in the figure below. The antiradical activity increases with the concentration of the sample. On the other hand, the activity of cooked samples is lower than that of fresh sample. Antioxidant activity values vary between 59.81% and 91.16%. They are equivalent to $91.16 \pm 0.23\%$, $84.7 \pm 0.5\%$, $82.59 \pm 1.89\%$, $81.37 \pm 0.32\%$ for raw African asparagus and samples CE5, CE10, CE15, respectively (Fig. 4). In contrast to fresh African asparagus, a variation in antioxidant capacity of boiled samples was observed (Fig. 4).




Fig. 3. Total polyphenol (A), flavonoid (B), and tannin (C) contents of fresh and boiled African asparagus

EC5: cooking 5 min in water; EC10: cooking 10 min in water; EC15: cooking 15 min in water




Fig. 4. DPPH free radical scavenging of fresh and water-cooked African asparagus extracts (EC) at different times (5, 10 and 15 min)

EC5: cooking 5 min in water; EC10: cooking 10 min in water; EC15: cooking 15 min in water; vit C: vitamin C

An increase in the percentage of DPPH radical inhibition was observed with increasing concentrations of fresh and cooked African asparagus extracts in water. Vitamin C was used as a standard.

Vitamin C and African asparagus extracts showed good DPPH radical scavenging activity. The IC₅₀ values of the raw extracts, samples CE5, CE10 and CE15 were 10.07 µg/mL, 12.07 µg/mL, 15.2 µg/mL and 10.33 µg/mL respectively. These different concentrations were higher than that of vitamin C (2.53µg/mL). The different extracts, as well as vitamin C, show significantly different IC₅₀ values at the 5% threshold. The raw sample extract showed the highest radical scavenging activity, followed by CE15 and CE5. On the other hand, under the same conditions, CE10 sample shows the lowest inhibitory activity (Table).

Table 1. IC₅₀ values of the different samples tested

Treatments	IC ₅₀
RawCE5	10.07
CE10	12.07
CE15	15.20
Vit C	10.33
	2.53

CE5 : cooking 5 min in water ; CE10 : cooking 10 min in water ; CE15 : cooking 15 min in water.

3.2 Discussion

The analysis of the heat treatment results shows the effect of cooking in water on the nutritional composition of African asparagus. The different cooking times to which the studied African asparagus are subjected significantly influence their studied nutritional parameters. Indeed, boiling resulted in a loss of more than 40-81% of the vitamin C content of African asparagus with the greatest loss (81%) observed in samples cooked for 15 minutes. These results corroborate those of Acho *et al.* [15] who recorded significant vitamin C losses ranging from 28.61 to 78.02% at 15 min of boiling in five of the leafy vegetables consumed in southern Côte d' Ivoire. The level of β-carotene also decreases during heat treatment and losses vary between 76 and 84%. These losses are attributed to the sensitivity of β-carotene to oxygen or leaching of nutrients from the vegetables during blanching or boiling [16]. Raw and cooked African asparagus contains more than 900 µg/100g of β-carotene, which is higher than the recommended daily intake for adults [17,18]. Indeed, β-carotene deficiency remains a public health problem affecting 19 million pregnant women in Africa [19] where the estimated requirement of this vitamin is 800 µg/day [20]. Thus, the consumption of African asparagus would be beneficial for this segment of the population. Boiling resulted in losses of up to 9.94% (total polyphenols), 19.92% (flavonoids) and 47.03% (tannins) after 15 minutes of

cooking. The decrease in flavonoid and tannin content is attributed to leaching of phenolic compounds during heat treatment [21]. The analysis of nutritional properties reveals that fresh African asparagus was rich in tannins with a content equal to (1954,84 mgTA/100gMS). However, this content decreased with increasing cooking time. Similar results were observed in relation to the study carried out by some authors that showed a loss of tannic acid after different cooking processes due not only to the degradation of tannins [7,22]. Tannins are involved in tissue regeneration. They help stop bleeding and help fight infections [23], especially in pre- and post-natal, as is the case in southern Ivory Coast. Regular consumption of African asparagus in households could thus prevent several diseases such as gastrointestinal disorders, high blood pressure and malaria [24].

As for the total polyphenol content, it increases slightly during the first 10 minutes of cooking. This result is in agreement with those of Kao *et al.* [25] obtained during their work on the boiling of Thai basil leaves and potatoes. Similarly, Turkmen *et al.* [5] have shown that short cooking time increases the polyphenol content of some green vegetables such as green beans, spinach and peppers. This increase in polyphenol content in the first few minutes of cooking is thought to be due to cell wall disruption, which released soluble phenolics from insoluble ester bonds, or to the heat-induced breakdown of dietary fiber-bound polyphenols into free phenolics [26, 27, 28]. But, the decrease in total phenolics after 10 minutes of cooking, could be explained by leaching of soluble or free polyphenols in the cooking water due to prolonged exposure to heat. However, these levels remain higher than those of Brou *et al.* [7] (291,43 to 909,79 mg/100g) recorded during the cooking of oil palm heart (*Elaeis guineensis jacq.*) Thus, the high content of total polyphenols found in fresh and cooked African asparagus would be useful for children, prone to allergies caused by anti-nutritional substances. They could also participate in the prevention of cardiovascular diseases and cancer in adults [29]. The decrease in antioxidant activity of African asparagus during the first 10 minutes of cooking is thought to be due to a leaching of some compounds such as tannins, β -carotene, vitamin C and flavonoids possessing high antioxidant activity [7]. However, the increase in antioxidant activity of African asparagus after 10 minutes of cooking is thought to be due to the synthesis of new compounds

such as Maillard reaction products with antioxidant activity [30].

4. CONCLUSION

African asparagus (*Laccosperma secundiflorum*) is a food rich in polyphenols, tannin and flavonoids, vitamin C and beta carotene which are compounds known for their antioxidant potential and health benefits. However, cooking significantly reduces the content of these compounds. However, after less than 10 minutes of cooking, more than half of almost all the compounds are still present in African asparagus. Therefore, to benefit from the antioxidant potential of African asparagus, it would be advisable to cook them in boiling water at times between 5 and 10 minutes.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Lenoir L. Protective effect of odorous verbena polyphenols in a model of colonic inflammation in rats. Human medicine and pathology. PhD thesis in Nutrition. University of Auvergne. Life and Health Sciences. 2011; 1.
2. Bravo L. Polyphenols: Chemistry, Dietary Sources, Metabolism, and Nutritional Significance. Nutrition Reviews. 1998;56 (11):317-333.
3. Mehinagic E, Bourles E, & Jourjon F. Fruit compounds of nutritional interest: impact of transformation processes on polyphenols. Swiss journal Viticulture, Arboriculture, Horticulture. 2011;43(6): 364-368.
4. Minussi RC, Rossi M, Bologna L, Cordi L, Rotilio D, Pastore GM. Phenolic compound and total antioxidant potential of commercial wines. Food Chemistry. 2003;82:409-416.
5. Turkmen N, Sari F, & Velioglu YS. The effect of cooking methods on total phenolics and antioxidant activity of selected green vegetables. Food Chemistry. 2005;93:713-718.
6. DaGiau S. "Ethnobotany, morpho-anatomy and phytogeography of useful palm (*Arecaceae*) species from Ivory Coast, with emphasis in rattan general. Master in *plant*

- biology, Faculty of Sciences (University of Geneva). 2014; 171.
7. Brou MR, Ekissi GSE, Fagbohoun JB, Faulet MB, Kouamé PL. Impacts of Boiling Times on Physicochemical and Nutritive Composition from Heart of Oil Palm Tree (*Elaeis guineensis* Jacq.) Consumed as Vegetable in Côte d'Ivoire. *Advances in Research*. 2018;16(5):1-16.
 8. Randrianatoandro VA. Identification and characterization of flat sources of micronutrients consumed in urban areas (*Manjakaray, Madagascar*): study of leafy vegetable dishes, thesis, Madagascar, Manjakaray. 2010; 150.
 9. Pongracz G, Weiser H, Matzinger D. Tocopherols-Antioxidant. *Fat Science Technology*. 1971;97:90-104.
 10. Tee ES, Kuladevan R, Young SI, Khor SC, Zakiah HO. Nutrient analysis of foods. Determination of Vitamine C, β -carotene and Riboflavin Contents in Five Green Vegetables Organically and Conventionally Grown. *Malaysian Journal of Nutrition*. 1996; 9(1):31-39.
 11. Singleton VL. Analysis of total phenols and other oxydant substrates and antioxydants by means of Follin-ciocalteur eagent. *Methode in Enzymology*. 1999; 299:152-178.
 12. Meda A, Lamien CE, Romito M, Millogo J & Nacoulma OG. Determination of total phenolic, flavonoid and proline contents in Burkina Faso honeys as well as their radical scavenging activity. *Food Chemistry*. 2005; 91:571-577.
 13. Baidridge Z, Tomlins K, & Westby A. Analysis of condensed tannins using acidified vanillin. *Journal of Food Science*.1996;29:77-79.
 14. Choi CW, Kim SC, Hwang SS, Choi BK, Ahn HJ, Lee MZ, Park SH & Kim SK. Antioxidant activity and free radical scavenging capacity between Korean medicinal plant and flavonoids by assay guided comparison. *Plant Sciences*. 2002; 163:1161-1168.
 15. Acho FC, Zoué LT, Koua YAG, Kra AKS, Niamké LS. Effect of cooking on nutritive and antioxidant properties of leafy vegetables consumed in Southern Côte d'Ivoire. *International Journal of Research in Biosciences*. 2014;3:75-87. Available : <http://www.ijrbs.in> ISSN 2319-2844
 16. Tessier FJ. Effect of cooking food on vitamin loss. In: *Correspondences in Metabolisms, Hormones, Diabetes and Nutrition*. 2012; 16:5-6.
 17. George DPR. *Newlife style: Enjoy it*. Editorial Safeliz. Spain. 1999;39-100.
 18. Akanya HO. *Retinol: The vitamin of life*. Federal University of Technology, Minna. Inaugural Lecture Series No. 5. Scan Prints Nig. Ltd. 2004;12.
 19. WHO. Global prevalence of vitamin A deficiency in population at risk 1995 – 2005. WHO Global Database on vitamin A Deficiency. Genève, suisse ; 2009.
 20. WHO/FAO. *Vitamin and mineral requirements in human nutrition 2 Nd ed* Genève. Organisation mondiale de la santé ; 2011b. Available :<http://www.who.int/nutrition/publications/micronutriments/9241546123/en/index.html>
 21. Wong SP, LP Leong, JHW Koh. Antioxidant activities of aqueous extracts of selected plants. *Alimentary Chemistry, food and sciences technology programs, department of chemistry, National University of Singapore 117543, Singapore*; 2006.
 22. Zhang D, Hamauzu Y. Phenolics, ascorbic acid, carotenoids and antioxidant activity of broccoli and their changes during conventional and microwave cooking. *Food Chemistry*. 2004b;88:503-509.
 23. Khanbaba K, Ree TR. Tannins: classification and Definition. *Natural Product Reports*. 2001; 18:641-649.
 24. Sereme A, Millogo-Rasolodimby J, Guinko S, Nacro M. Therapeutic property of tannin plants from Burkina Faso. *African Pharmacopoeia and Traditional Medicine*. 2008; 15:41-49.
 25. Kao FJ, Chiu YS, Chiang WD. Effect of water cooking on the antioxidant capacity of carotenoid-rich vegetables in Taiwan. *J. Food Drug Anal*. 2013;2:202–209. DOI: 10.1016/j.jfda.2013.09.010.
 26. Stewart AJ, Bozonnet S, Mullen W, Jenkins GI, Lean MEJ, Crozier A. Occurrence of flavonols in tomatoes and tomato-based products. *Journal of Agriculture and Food Chemistry*. 2000; 48(7):2663–2669. DOI: 10.1021/jf000070p.
 27. Dewanto V, Wu X, Liu RH. Processed sweet corn has higher antioxidant activity. *Journal of Agriculture and Food Chemistry*. 2002;50:4959-4.
 28. Adefegha SA, Oboh G. Enhancement of total phenolics and antioxidant properties

- of some tropical green leafy vegetables by steam cooking. J Food Process Preserv. 2011;35:615-622.
29. Talbi H, Boumaza A, El-mostafa K, Talbi J & Hilali A. Evaluation de l'activité antioxydante et la composition physico-chimique des extraits méthanolique et aqueux de la *Nigella sativa* L. (Evaluation of antioxidant activity and physico-chemical composition of methanolic and aqueous extracts of *Nigella sativa* L.). Journal of Materials and Environmental Science. 2015; 6(4):1111-1117.
30. Morales FJ, Babel M-B. Antiradical efficiency of Maillard reactin mixtures in a hydrophilic media. Journal of Agriculture and Food Chemistry. 2002;50: 2788–92.

© 2021 Kouassi et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<https://www.sdiarticle5.com/review-history/79895>