



# Co-occurrence of Intestinal Parasites among School Children of Akonolinga, Centre Region of Cameroon: Emergency Need to Reduce the Health Divide

Oyono Martin Gael <sup>a,b\*</sup>, Fosso Samuel <sup>c</sup>,  
Njua-Yafi Clarisse <sup>a,d</sup>, Lehman Léopold Gustave <sup>e</sup>  
and Bilong Bilong Charles Félix <sup>a</sup>

<sup>a</sup> Laboratory of Parasitology and Ecology, University of Yaoundé I, P.O Box: 812, Yaoundé, Cameroon.

<sup>b</sup> Institute of Medical Research and Medicinal Plants Studies, P.O. Box 13033, Yaoundé, Cameroon.

<sup>c</sup> Laboratory of Parasitology, Mycology and Parasitic Immunology, Yaoundé Teaching Hospital, P.O. Box: 1364, Yaoundé, Cameroon.

<sup>d</sup> Department of Animal Production Technology, University of Bamenda, P.O Box 39, Bamilli, Cameroon.

<sup>e</sup> Laboratory of Animal Biology and Physiology, University of Douala, P.O. Box: 2701, Douala, Cameroon.

## Authors' contributions

*This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.*

## Article Information

DOI: 10.9734/IJTDH/2022/v43i221364

### Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/92955>

Original Research Article

Received: 01/09/2022  
Accepted: 02/11/2022  
Published: 08/11/2022

\*Corresponding author: Email: [oyonomartingael2.0@gmail.com](mailto:oyonomartingael2.0@gmail.com), [oyonomartin@impm-cm.org](mailto:oyonomartin@impm-cm.org);

## ABSTRACT

**Background:** Intestinal parasitosis pose a serious problem to public health and development, especially for the underprivileged population of low and middle-income countries. School children, who are the most affected, can harbour several parasites at the same time. There is a need to adopt efficient strategies for the elimination of intestinal parasitic infections as a public health problem by 2030.

**Methods:** This cross-sectional study was conducted to highlight the level of co-occurrence of intestinal parasites among pupils of the Akonolinga locality of the Centre region of Cameroon. Pupils were randomly recruited from 5 public schools; stool samples were collected from those whose parents agreed to participate in the study. Helminth eggs and protozoan cysts were assessed by direct smear and the formol-ether concentration technique.

**Results:** Out of the 416 pupils recruited, 252 were infected by at least one of the ten intestinal parasites identified. The most frequent soil-transmitted helminths were *Ascaris lumbricoides* and *Trichuris trichiura* with prevalences of 21.4 % and 18.5 %, respectively. *Entamoeba coli* was the main protozoan followed by *E. histolytica/dispar* with prevalences of 29.3% and 23.8%, respectively, and *Embadomonas intestinalis* was found in only one pupil (0.2%). The co-occurrence of intestinal parasites reached 50% and children with up to four different parasites were detected. The pair of intestinal protozoan parasites, *E. histolytica/dispar* and *E. coli* were the most observed with a frequency of 9.5%. Living in a rural setting and young age were important determinants for most of the parasite infections and co-occurrence of parasites.

**Conclusion:** The co-occurrence of intestinal parasites among school children of Akonolinga is high, and this situation is more alarming in rural areas and in younger children. The living conditions and the co-occurrence of parasites must be integrated into public policies for fighting against intestinal parasite infections.

**Keywords:** Intestinal parasites; co-occurrence; school children; health divide.

## 1. INTRODUCTION

Ecological communities include a complex network of biotic interactions such as predation, mutualism, competition, facilitation and parasitism [1]. For parasites, hosts can be considered as ecosystems that have easily defined resources [2]. In most cases, the host is infected simultaneously by several parasite species; as a result of common ecological and environmental requirements, infection routes, host exposure, host susceptibility, behavioural, sociological, and economic factors that enable the co-occurrence of parasite-host systems [3,4]. Co-infections have important public health, epidemiological and evolutionary implications. They can lead to different outcomes for both the host and the parasites compared to single parasite infection. For instance, co-infections can increase the parasite virulence evolution rate and lead to higher morbidity and mortality of the hosts. The within-host interactions between parasites can regulate their coexistence with mutually beneficial, antagonistic or neutral impacts on one another, leading to higher or lower reproductivity [5,6].

Intestinal helminths and protozoa are among the most important causes of infections in developing

countries, especially in tropical and subtropical countries carrying high burdens of morbidity and mortality [7]. Epidemiological evidence suggests that the soil-transmitted helminths, *Ascaris lumbricoides*, *Trichuris trichiura* and Hookworms, affect more than 1.5 billion people; more than 267 million preschool children and 568 million school children live in areas where these parasites are intensively transmitted [8]. Regarding protozoa species, *Cryptosporidium* spp., *Entamoeba histolytica* and *Giardia intestinalis* are the most common protozoans infecting children in sub-Saharan Africa [9]. School-aged children are the most affected, due to their habits of playing or handling infected soil, eating with soiled hands, unhygienic toilet practices, drinking and eating contaminated water and food respectively [10]. Intestinal parasites consume nutrients from the children they infect, thus retarding their physical and cognitive development. They destroy tissues and organs, cause abdominal pain, diarrhoea, intestinal obstruction, anaemia, ulcers, and other health problems which can lead to slow cognitive development and impaired learning [10,11].

In 2015, the United Nations adopted 17 Sustainable Developments Goals (SDGs) to end poverty, protect the planet, and ensure that by

2030 all people enjoy peace and prosperity [12]. In order to attain the SDG 3: « Ensure healthy life and promote well-being for all at all ages», WHO proposed a road map for NTDs for 2021–2030 [13]; so, there is a need to adopt efficient strategies. Hence, research on the co-occurrence of parasites, increasingly recognized as the norm, could be a prerequisite to designing effective strategies to guarantee the elimination of intestinal parasitic infections as a public health problem. In Cameroon, intestinal parasites are distributed all over the country and several studies have been conducted to prove this [3,14-16]. However, there are still several localities for which epidemiological information is needed. In this paper we examined the co-occurrence and health impact of intestinal parasites among school children of the Akonolinga sub-division, Centre Region of Cameroon.

## 2. MATERIALS AND METHODS

### 2.1 Study Area

This study was carried out between September 2017 and July 2018 in the locality of Akonolinga in the centre region of Cameroon. Akonolinga is located 100 km east of Yaoundé (the political capital of Cameroon). It is the Head Quarters of the Nyong and Mfoumou Division, and situated at 3°46' North latitude, 12°15' East longitude, and 673 meters above sea level (a.s.l.). It covers an area of about 1,420 Km<sup>2</sup> with approximately 18.000 inhabitants [17]. The environment is predominantly tropical forest and the division is irrigated by the Nyong River system which flows from east to west.

### 2.2 Study Population

The study population consisted of school children of the Akonolinga locality. Four hundred and sixteen (416) pupils aged 4 to 15 years old were randomly sampled in five primary schools: three in a rural area (Essang-Ndibi, Kpwele, Eboa) and two in an urban area (Loum, centre ville).

### 2.3 Study Design

A questionnaire was used for the collection of socio-demographic and behavioural data. Each child received information on how to collect stool samples in order to avoid contamination with urine or other contaminants. Early-morning stool samples were collected in plastic screw-cap vials, labelled with each participant's information. All stool samples collected were put in an ice bag and transported to the Parasitology laboratory at

the Yaoundé University Teaching Hospital, for parasitological analysis. Children whose parents or guardians did not give their consent, those who were recently enrolled in school, and those who had taken anti-helminthic treatment at least one month before, were excluded from the study as well as participants who didn't provide stool samples.

### 2.4 Parasitological Examination and Morphological Identification

Stool samples were analysed using direct smears in saline and Lugol solution, and also by the formalin-ether concentration technique as described by Allen and Ridley [18]. Briefly, about a pea size of stool samples were emulsified in 7 ml of formalin, sieved through a wire gauze into an evaporating basin, and then transferred into a centrifuge tube. Then, 3 ml of ether were added, the mixture agitated vigorously for 30 seconds and centrifuged at 3,000 rpm for 60 seconds. The layer of debris accumulated at the interphase was removed by passing a swab stick gently round the circumference of the tube. The supernatant was discarded and the pellet was homogenised and mounted on a glass slide and cover slip for microscopic examination. Samples were observed at a magnification of x200 and x400 using the CyScope® microscope (Partec-Sysmex GmbH, Görlitz, Germany) and the identification of parasite eggs or cysts was done using photos and dimension ranges according to the WHO Bench Aid for the diagnosis of intestinal parasites. All stool samples were analysed in duplicate.

### 2.5 Statistical Analysis

Data obtained were entered and stored in a Microsoft Excel spread sheet then exported to the SPSS (Statistical Package for Social Sciences) version 25.0 software for subsequent analyses. Prevalence was defined as the number of individuals infected with a particular parasite species x 100, divided by the total number of examined individuals [19], while the co-occurrence rate was defined as the number of simultaneous appearances of different parasite species x 100, divided by the total number of infected individuals. The prevalences of different intestinal parasites were compared between age groups, genders, and living areas, using the chi-square test ( $\chi^2$ ). The logistic regression model was performed to determine the risk factor(s) of co-occurrence of intestinal parasite infections. Prevalence and co-occurrence rates were tested at 5% significance level ( $P < 0.05$ ).

### 3. RESULTS

A total of 416 stool samples were collected from school children and examined during the current study. From the total number recruited, 164 (39.4%) children were uninfected while 252 (60.6 %) were infected with at least one intestinal parasite (protozoa and/or helminth). Intestinal protozoans were present in 199 (47.8%) cases, and helminths in 135 (32.5%) cases.

Amongst the Ten (10) parasite species recorded, four (04), six (05) and one (01) were helminths, protozoans and chromista, respectively. Helminth species included *Ascaris lumbricoides* Linnaeus, 1758, *Trichuris trichiura* Linnaeus, 1771, *Hymenolepis nana* Siebold, 1852, and Hookworms (*Necator americanus* Stiles, 1902 / *Ancylostoma duodenale*, Dubini, 1843). Protozoan species included *Entamoeba histolytica/dispar* Schaudinn 1903/Brumpt, 1925, *E. coli* Loesch, 1875, *Endolimax nana* Kuenen et Swellengrebel, 1917, *Giardia intestinalis* Kofoid et Christiansen, 1915, and *Embadomonas intestinalis* Grassi, 1879. *Blastocystis* spp Alexieff, 1911 is the chromista specie found. The most prevalent species among helminths and protozoa were *A. lumbricoides* (21.4%) and *E. coli* (29.3%) respectively as shown in Fig. 1.

The prevalence of intestinal parasites according to living areas, age groups and gender are presented in Table 1. The prevalence of *E. coli*,

*G. intestinalis*, *A. lumbricoides* and *T. trichiura* were significantly higher in the rural areas compared to the urban areas. Although not significant ( $P > 0.05$ ), the prevalence of *E. coli*, *Blastocystis* spp., *Em. intestinalis*, *En. nana*, *T. trichiura* and Hookworms, were higher in children aged 4 to 8 years old. Conversely, the prevalence of *E. histolytica/dispar* ( $P = 0.042$ ) and *A. lumbricoides* ( $P = 0.255$ ) tended to increase significantly with age. In most cases (except *Em. intestinalis* and *En. nana*), females were more often infected than males.

Co-occurrence of parasite species was recorded in 126 (50%) of the 252 pupils found infected by at least one parasite species. The majority of simultaneous infections consisted of two parasite species (33.7%, 85/252), then three parasite species (13.1 %; 33/252), and only 3.2% (8/252) cases with four parasite species. Fig. 2 illustrates the pattern of the different types of parasitic co-occurrence. The most prevalent (co-occurrence > 5%) parasitic co-infections were in decreasing order *E. histolytica/dispar* + *E. coli* (9.5 %; 24/252), *A. lumbricoides* + *E. coli* (6.3 %; 16/252), *T. trichiura* + *E. coli* (6.3 %; 16/252) and *A. lumbricoides* + *T. trichiura* + *E. coli* (6.0 %; 15/252). Apart from *H. nana*, all other parasite species were involved in co-infections; *A. lumbricoides* (87.3 %; 110/126) was the most involved in parasite co-infections, followed by *E. coli* (71.4 %, 90/126) and *T. trichiura* (47.6 %; 60/126).





Fig. 1. Prevalence of intestinal parasites in pupils of the Akonolinga locality

**Table 1. Prevalence of intestinal parasites according to living areas, age groups and gender**

Parasite species	Living area			Age group				Gender		
	Rural (N <sub>1</sub> = 209)	Urban (N <sub>2</sub> = 207)	P	[4 – 8] (x <sub>1</sub> = 135)	[8 – 12] (x <sub>2</sub> = 169)	[12 – 16] (x <sub>3</sub> = 112)	P	Female (n <sub>1</sub> = 226)	Male (n <sub>2</sub> = 190)	P
<i>E. coli</i>	89 (42.6)	33 (16.0)	0.001*	42 (31.1)	51 (30.2)	29 (25.9)	0.636	74 (32.7)	48 (25.3)	0.095
<i>E. histolytica/dispar</i>	46 (22)	53 (25.6)	0.389	22 (16.3)	45 (26.6) \$	32 (28.6) \$	0.042*	61 (27.0)	38 (20.0)	0.095
<i>G. intestinalis</i>	13 (6.2)	04 (1.9)	0.027*	04 (2.9)	10 (5.9)	03 (2.7)	0.294	04 (1.7)	13 (6.8) \$	0.027*
<i>Blastocystis</i> spp.	06 (2.8)	00 (0.0)	/	03 (2.2)	01 (0.6)	02 (1.4)	0.465	04 (1.7)	02 (1.1)	0.541
<i>Em. intestinalis</i>	01 (0.5)	00 (0.0)	/	01 (0.7)	-	-	/	-	01 (0.5)	/
<i>En. nana</i>	08 (3.9)	05 (2.4)	0.407	06 (4.4)	04 (2.4)	03 (2.7)	0.556	06 (2.6)	07 (3.7)	0.547
<i>A. lumbricoides</i>	70 (33.5)	19 (9.2) \$	0.000*	23 (17.0)	42 (24.8)	24 (21.4)	0.255	57 (25.2) \$	32 (16.8)	0.039*
<i>T. trichiura</i>	65 (31.1)	12 (5.8) \$	0.000*	30 (22.2)	33 (19.5)	14 (12.5)	0.133	47 (20.8)	30 (15.8)	0.19
Hookworms	02 (0.9)	02 (0.9)	/	03 (2.2)	01 (0.6)	00 (0.0)	/	03 (1.3)	01(0.5)	/
<i>H. nana</i>	00 (0.0)	01 (0.5)	/	-	01 (0.5)	-	/	01 (0.5)	00 (0,0)	/

N<sub>1</sub>: number of schoolchildren recruited in rural area; N<sub>2</sub>: number of schoolchildren recruited in urban area; x<sub>1</sub>: number of schoolchildren aged 4 to 7 years; x<sub>2</sub>: number of schoolchildren aged 8 to 11 years; x<sub>3</sub>: number of schoolchildren aged 12 to 15 years; n<sub>1</sub>: number of female in the study population; n<sub>2</sub>: number of male in the study population; E: Entamoeba; G: Giardia; Em: Embadomonas; En: Endolimax; A: Ascaris; T: Trichiura; H: Hymenolepis; \*: P-value significant; \$: Prevalence significantly high



**Fig. 2. Patterns of different types of parasitic co-occurrences in school children of Akonolinga**

**Table 2. Risk factors associated with single infection and co-occurrent intestinal parasitic infections in school children of the locality of Akonolinga**

Variable		S.E.	Khi Square	P-value	OR	95 % CI
<i>A. lumbricoides</i>	Female	0.26	3.29	0.07	0.62	0.37 – 1.04
	Age		3.68	0.16		
	[4 – 7] years	0.35	3.66	0.06	0.51	0.26 – 1.02
	[8 – 11] years	0.32	1.02	0.31	0.72	0.38 – 1.36
	Rural	0.30	32.6	0.000*	<b>1.80</b>	<b>1.10 – 3.30</b>
<i>T. trichiura</i>	Female	0.27	1.13	0.28	0.75	0.43 – 1.27
	Age		2.01	0.36		
	[4 – 7] years	0.38	1.08	0.29	1.48	0.71 – 3.11
	[8 – 11] years	0.37	0.000	0.98	0.99	0.48 – 2.06
	Rural	0.34	34.6	0.000*	<b>1.30</b>	<b>1.07 – 2.60</b>
<i>E. histolytica</i>	Female	0.24	3.34	0.06	0.64	0.41 - 1.03
	Age		6.58	0.03		
	[4 – 7] years	0.32	5.19	0.02*	<b>0.48</b>	<b>0.26 – 0.90</b>
	[8 – 11] years	0.28	0.03	0.84	0.94	0.54 – 1.65
	Rural	0.24	0.74	0.38	1.23	0.77 – 1.98
<i>E. coli</i>	Female	0.23	2.29	0.13	0.70	0.45 – 1.11
	Age		0.87	0.64		
	[4 – 7] years	0.30	0.001	0.96	0.98	0.54 – 1.79
	[8 – 11] years	0.29	0.58	0.44	0.79	0.45 -1.43
	Rural	0.24	33.0	0.000*	<b>2.40</b>	<b>1.50 – 4.00</b>
<i>T. trichiura + E. coli</i>	Female	0.37	0.90	0.34	0.70	0.39 – 1.46
	Age		1.73	0.42		
	[4 – 7] years	0.48	0.87	0.34	0.64	0.25 – 1.63
	[8 – 11] years	0.46	1.68	0.19	0.55	0.23 – 1.35
	Rural	0.47	14.5	0.000*	<b>1.60</b>	<b>1.16 – 4.20</b>
<i>E. histolytica/dispar + E. coli</i>	Female	0.35	0.70	0.40	0.74	0.37 – 1.48
	Age		0.10	0.94		
	[4 – 7] years	0.47	0.01	0.90	0.94	0.38 – 2.37
	[8 – 11] years	0.45	0.09	0.75	0.87	0.36 – 2.10
	Rural	0.40	10.7	0.001*	<b>2.60</b>	<b>1.20 – 5.80</b>
<i>A. lumbricoides + E. coli</i>	Female	0.34	0.000	0.99	1.00	0.51 – 1.96
	Age		0.74	0.69		
	[4 – 7] years	0.47	0.74	0.38	0.66	0.26 – 1.68
	[8 – 11] years	0.43	0.26	0.60	0.80	0.34 – 1.87
	Rural	0.44	15.36	0.00*	<b>1.70</b>	<b>1.17 – 4.20</b>
<i>A. lumbricoides + T. Trichiura + E. coli</i>	Female	0.47	1.95	0.16	1.94	0.76 – 4.96
	Age		2.81	2.45		
	[4 – 7] years	0.62	1.94	0.16	0.42	0.12 – 1.42
	[8 – 11] years	0.56	2.26	0.13	0.42	0.14 – 1.29
	Rural	0.65	9.64	0.000*	<b>1.13</b>	<b>1.30– 4.70</b>

Legend: **SE**: Standard Error; **OR**: Odds Ratio; **E**: Entamoeba; **G**: Giardia; **Em**: Embadomonas; **En**: Endolimax; **A**: Ascaris; **T**: Trichiura; **H**: Hymenolepis; \*: P-value significant

Table 2 outlines some of the risk factors involved in single infections and co-occurrent intestinal parasitic infections. Only single and concomitant parasitic infections with frequency  $\geq 5\%$  were included in the model of logistic regression. Rural areas appeared as the main risk factor for both single and co-occurrent intestinal parasitic infections. This was the case with single infections by: *A. lumbricoides* (OR = 1.80; 95%

CI: 1.10 – 3.30;  $P < 0.001$ ), *T. trichiura* (OR = 1.30, 95%CI: 1.07 – 2.60,  $P < 0.001$ ), and *E. coli* (OR = 2.40, 95%CI: 1.50 – 4.00,  $P < 0.001$ ), and with concomitant infections by: *T. trichiura + E. coli* (OR = 1.16, 95%CI: 1.60 – 4.20,  $P < 0.001$ ), *E. histolytica/dispar + E. coli* (OR = 2.60, 95% CI: 1.20 – 5.80,  $P < 0.001$ ), *A. lumbricoides + E. coli* (OR = 1.70, 95%CI: 1.17 – 4.20,  $P < 0.001$ ) and *A. lumbricoides + T. trichiura + E. coli* (OR =

1.13, 95%CI: 1.30 – 4.70,  $P < 0.001$ ). The 4 - 7 years old age group was significantly ( $P < 0.05$ ) associated with *E. histolytica/dispar* infection.

#### 4. DISCUSSION

It is increasingly recognized that host co-infection with multiple pathogens is the norm, and that a better understanding of the nature and extent of multi-parasitism can have important epidemiological, clinical and control implications [20]. The human intestinal tract harbours a community of micro-organisms including viruses, bacteria, fungi, protozoa and helminths [21]. Recent research has revealed interactions between micro-organisms in the gut, with effects on the parasites themselves as well as on host nutrition and organ pathology [22-24].

In low and middle-income countries, children bear the greatest burden of intestinal parasitic infections in terms of morbidity and mortality. In children, intestinal parasitic infections, especially helminthiasis, may cause a number of negative health outcomes such as malnutrition, anaemia, impaired growth and cognitive development. Additionally, these infections may contribute substantially to increase the burden of poverty, impaired mental and educational development in children and damage economic productivity [25-27].

Our results revealed the circulation of at least 10 intestinal parasite species in the locality of Akonolinga. Many of them, nematodes (*A. lumbricoides*, Hookworms and *T. trichiura*) and Amoebae (*E. histolytica/dispar* and *E. coli*), were found at different infection rates in many other localities in Cameroon: Littoral [28], West [15], South [29], and in the South-west [30] regions. The variation in the prevalence may be explained by the geographical and ecological factors, behavioural habits and living conditions of the study populations. High prevalences of intestinal parasitic infections are still common in school children as indicated by this current study which showed a high intestinal parasite infection prevalence rate of 60.6%. These results suggest that the de-worming program put in place may be insufficient or ineffective in eradicating the infection, particularly in this population. Mass drug administration may have a limited effect and doesn't prevent the re-infestation [31]. The implementation of complementary community-based health interventions such as health education or WASH (Water, sanitation and hygiene) programs are strongly recommended in

order to achieve the elimination of intestinal parasitosis and accelerate research on vaccine discovery. The prevalences of many parasites were significantly higher in the rural setting compared to the urban setting, especially for *A. lumbricoides*, *T. trichiura*, *E. coli* and *G. intestinalis*. In low and middle-income countries, rural communities face many difficulties such as poverty, poor sanitation, untreated water and inadequate drainage systems, poor healthcare, promiscuity and dwellings made up of temporary materials [32,33]. All these features contribute to poor living conditions in rural areas, and thus a widespread and a stable transmission of several intestinal parasite species whose transmission stages have the capacity to resist for a long time in the environment [34]. These parasitic infections cause socio-economic devastation and hamper healthcare in rural vulnerable communities [35-37]. There is therefore an emergent need to reduce the health divide through the promotion of mass de-worming or parasite control, health education and access to healthcare facilities in rural areas. This is a prerequisite to the elimination of intestinal parasitic infections as a public health problem, towards the attainment of sustainable health development goals.

This study also revealed that co-occurrence of intestinal parasites occurs in one out of two cases (50%) and that a single individual can simultaneously harbour up to four parasite species. Our findings are similar to published data and confirm that in general, multiparasitism is the rule rather than the exception [38-40]. Both single parasite and multiparasite infections had common risk factors mostly related to the children's individual conditions and habits. In fact, the simultaneous presence of many parasite species can lead to synergistic or antagonistic interspecific interactions between parasites. Such interactions can have important repercussions on the host's health because they can alter host susceptibility to other parasites, infection duration, transmission risks, clinical symptoms and consequently treatment and prevention strategies [4].

#### 5. CONCLUSION

In the Akonolinga locality, the co-occurrence of intestinal parasites among school children is quite frequent, mainly in the rural areas where the highest prevalence of many intestinal parasite infections was recorded. To ensure a healthy life and well-being for all, it is important to



reduce the health divide through the promotion of mass deworming or parasite control, health education and access to health care facilities in rural areas. Furthermore, it's also important to develop new approaches that will help highlight the co-occurrence of parasites, and integrate this neglected reality into strategies for the elimination of intestinal parasitic infections as a public health problem by 2030.

## 6. STUDY LIMITATION

The fact that only a single stool sample was analysed per child, may underestimate the prevalence of intestinal parasites and this could be a possible limitation of this study. However, to minimize this situation, each sample was also examined by using the formalin-ether concentration technique; thus, increasing the diagnostic accuracy.

## ETHICAL APPROVAL AND CONSENT

This study was approved by the National Ethical Committee of Research for Human Health (Ethical Clearance N°: 2018/01/968/CE/CNERSH/SP) and the Directorate of the Yaoundé University Teaching Hospital (Research Authorization N°: 894/AR/CHUY/DG/DGA/DMT). In this cross-sectional study; before samples were collected, informed consent was obtained from the parents and/or legal guardians of the pupils after they understood the purpose of the study.

## ACKNOWLEDGEMENTS

The authors would like to thank Dr. Shey BONG AKEE for her technical assistance.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Fontaine C, Guimarães Jr P, Kéfi S, Loeuille N, Memmott J, van Der Putten W, Van Veen F, Thébault E. The ecological and evolutionary implications of merging different types of networks. *Ecology Letters*. 2011;14:1170–1181.
2. Rynkiewicz E, Pedersen A, Fenton A. An ecosystem approach to understanding and managing within-host parasite community dynamics. *Trends of Parasitology*. 2015; 31:212–221.
3. Brooker S, Clements A, Bundy D. Global epidemiology, ecology and control of soil-transmitted helminth infections. *Advances in Parasitology*. 2006;62:221–261.
4. Staudacher O, Heimer J, Steiner F, Kayonga Y, Havugimana J, Ignatius R. Soil-transmitted helminths in Southern Highland Rwanda: Associated factors and effectiveness of school-based preventive chemotherapy. *Tropical Medicine & International Health*. 2014;19:812–24.
5. Vasco D, Wearing H, Rohani P. Tracking the dynamics of pathogen interactions: Modelling ecological and immune-mediated processes in a two-pathogen single-host system. *Journal of Theoretical Biology*. 2007;245:9–25.
6. Susi H, Barrès B, Vale P, Laine A. Co-infection alters population dynamics of infectious disease. *Nature Communications*. 2015;6:5975.
7. Pullan R, Smith J, Jasrasaria R, Brooker S. Global numbers of infection and disease burden of soil transmitted helminth infections in 2010. *Parasites and Vectors*. 2014;7(1):37.
8. World Health Organization. Soil-transmitted helminth infections: Fact sheets; 2019.
9. Wang H, Naghavi M, Allen C, Barber R, Bhutta Z, Carter A. Global, regional, and national life expectancy, all-cause mortality, and cause-specific mortality for 249 causes of death, 1980-2015: A systematic analysis for the global burden of disease study 2015. *Lancet*. 2016; 388(10053):1459–544.
10. Chelkeba L, Mekonnen Z, Alemu Y, Eman D. Epidemiology of intestinal parasitic infections in preschool and school-aged Ethiopian children: A systematic review and meta-analysis. *BMC Public Health*. 2020;20:117.
11. Faria C, Zanini G, Dias G, da Silva S, de Freitas M, Almendra R. Geospatial distribution of intestinal parasitic infections in Rio de Janeiro (Brazil) and its association with social determinants. *PLoS Neglected Tropical Diseases*. 2017;11(3): e0005445.
12. United Nation. Transforming our world: the 2030 agenda for sustainable development. A/RES/70/1; 2015.
13. World Health Organization. Ending the neglect to attain the Sustainable

- Development Goals: A road map for neglected tropical diseases 2021–2030; 2020.
14. Tchuem Tchuente L, Dongmo C, Ngassam P, Kenfack M, Gipwe F, Dankoni E, Tarini A, Zhang Y. Mapping of schistosomiasis and soil-transmitted helminthiasis in the regions of Littoral, North-West, South and South-West Cameroon and recommendations for treatment. *BMC Infectious Diseases*. 2013;13:602.
  15. Matsinkou M, Yamssi C, Mbong E, Noumedem A, Tateng N, Megwi L, Khan Payne V. Intestinal helminth infections and associated risk factors among school-aged children of Bamendjou Community, West Region of Cameroon. *Journal of Parasitology Research*. 2021;6665586:8.
  16. Ndifor F, Hamit M, Abakar I, Noukeu C, Issa R, Brahim O. Prevalence of intestinal parasites and polyparasitism among children less than five years in the Logbaba district hospital in Douala, Cameroon. *European Journal of Biomedical and Pharmaceutical Sciences*. 2021;8(2):38-44.
  17. Bayaga N, Guedje M, Biye H. Approche entobotanique et ethnopharmacologique des plantes utilisées dans le traitement traditionnel de l'ulcère de Buruli à Akonolinga (Cameroun). *International Journal of Biological and Chemical Sciences*. 2017;11(4):1523-1541.
  18. Allen A, Ridley D. Further observations on the formol-ether concentration technique for faecal parasites. *Technical Methods*. 1970;545-546.
  19. Bush A, Lafferty K, Lotz J, Shostak A. Parasitology meets ecology on its own terms: Margolis et al. revisited. *Journal of Parasitology*. 1997;83(4):575-583.
  20. Donohue R, Cross Z, Michael E. The extent, nature, and pathogenic consequences of helminth polyparasitism in humans: A meta-analysis. *PLoS Neglected Tropical Diseases*. 2019;13(6):e0007455.
  21. Milani C, Duranti S, Bottacini F, Casey E, Turrone F, Mahony J, Belzer C, Delgado S, Arboleya S, Mancabelli L, Lugli G, Rodriguez J, Bode L, de Vos W, Gueimonde M, Margolles A, van Sinderen D, Ventura M. The first microbial colonizers of the human gut: composition, activities, and health implications of the infant gut microbiota. *Microbiology and Molecular Biology Reviews*. 2017;81:e00036-17.
  22. Pullan R, Brooker S. The health impact of polyparasitism in humans: Are we underestimating the burden of parasitic diseases? *Parasitology*. 2008;135(7):783–94.
  23. Griffiths E, Pedersen A, Fenton A, Petchey O. The nature and consequences of coinfection in humans. *Journal of Infection*. 2011;63(3):200–6.
  24. Bilong Bilong C, Oyono M, Fosso S, Lehman L. Ecological characterization of interspecific relationships between human parasites: Conflict, cooperation or independence? *Cameroon Journal of Experimental Biology*. 2021;15(1):9-15.
  25. Oninla S, Onayade A, Owa J. Impact of intestinal helminthiasis on the nutritional status of primary-school children in Osun state, South Western Nigeria. *Annals of Tropical Medicine and Parasitology*. 2010; 104:583–594.
  26. Fürst T, Keiser J, Utzinger Jr. Global burden of human food-borne trematodiasis: A systematic review and meta-analysis. *Lancet Infectious Diseases*. 2012;12:210–221.
  27. Sayasone S, Utzinger J, Akkhavong K, Dermatt P. Multiparasitism and intensity of helminth infections in relation to symptoms and nutritional status among children: A cross-sectional study in southern Lao People's Democratic Republic. *Acta Tropica*. 2015;141:322–331.
  28. Yamssi C, Nadia N, Payne V, Sabi B, Romeo N. Gastrointestinal nematodes among residents in Melong, Moundou division, Littoral Region, Cameroon. *BioMed Research International*. 2021;9: 5368973.
  29. Ngo Ngué N, Nkengazong L, Nkoa T, Adamou M, Ngué M, Motsebo A, Moyou-Somo R, Nukenine E. Assessment of antihelminthic efficacy of Mebendazole 500 mg against soil-transmitted helminths in a high-risk area of Cameroon. *International Journal of Tropical Disease & Health*. 2017;25(3):1-13.
  30. Mbuh J, Ntonfor N, Ojong J. The epidemiology of soil-transmitted helminth and protozoan infections in South-West Cameroon. *Journal of Helminthology*. 2012;86(1):30-7.
  31. Tchuem Tchuente LA, Momo SC, Stothard JR, Rollinson D. Efficacy of praziquantel and reinfection patterns in single and mixed infection foci for intestinal and

- urogenital schistosomiasis in Cameroon. *Acta Tropica*. 2013;128:275–3.  
DOI: 10.1016/j.actatropica.2013.06.007
32. Cronk R, Guo A, Fleming L, Bartram J. Factors associated with water quality, sanitation, and hygiene in rural schools in 14 low- and middle-income countries. *Science of the Total Environment*. 2020; 144226.  
DOI: 10.1016/j.scitotenv.2020.144226
33. Bolatova Z, Tussupova K, Toleubekov B, Sukhanberdiyev K, Sharapatova K, Stafström M. Challenges of access to WASH in schools in low- and middle-income countries: Case study from Rural Central Kazakhstan. *International Journal of Environment Research and Public Health*. 2021;18:9652.  
DOI: doi.org/10.3390/ijerph18189652
34. Suntaravitun P, Dokmaikaw A. Prevalence of intestinal parasites and associated risk factors for infection among rural communities of Chachoengsao Province, Thailand. *Korean Journal of Parasitology*. 2018; 56(1):33-39.  
DOI: doi.org/10.3347/kjp.2018.56.1.33
35. Adane D, Endalkachew N, Abaineh M. Prevalence of intestinal parasitic infections and associated risk factors among pregnant women attending antenatal care center at Felege Hiwot referral hospital, Northwest Ethiopia. *BMC Infectious Diseases*. 2016;16:53.
36. Nathan C, Jedidiah S, David G, Sam H, Jason R. Deworming in pre-school age children: A global empirical analysis of health outcomes. *PLoS Neglected Tropical Diseases*. 2018;12:5.
37. Gadisa, Jote. Prevalence and factors associated with intestinal parasitic infection among underfive children in and around Haro Dumal Town, Bale Zone, Ethiopia. *BMC Pediatrics*. 2019;19:385.
38. Bordes F, Morand S. The impact of multiple infections on wild animal hosts: A review. *Infection Ecology and Epidemiology*. 2011;1:7346.  
DOI: 10.3402/iee.v1i0.7346
39. Sayasone S, Mak T, Vanmany M, Rasphone O, Vounatsou P. Helminth and intestinal protozoa infections, multiparasitism and risk factors in Champasack Province, Lao People's Democratic Republic. *PLoS Neglected Tropical Diseases*. 2011;5(4):e1037.  
DOI: 10.1371/journal.pntd.0001037
40. Vaumourin E, Vourc'h G, Gasqui P, Vayssier-Taussat M. The importance of multiparasitism: Examining the consequences of co-infection for human and animal health. *Parasites & Vectors*. 2015;8:545.

© 2022 Oyono et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/92955>