



# Influence of Different Organic Manures and Inorganic Fertilizers on Vegetative Growth of Turnip (*Brassica rapa* L.) cv Purple Top White Globe

Gyanendra Singh<sup>a++\*</sup>, Vipin Masih Prasad<sup>a#</sup>,  
Vijay Bahadur<sup>a†</sup>, Nikhil Vikram Singh<sup>a++</sup>  
and Netra Pal Yadav<sup>a++</sup>

<sup>a</sup> Department of Horticulture, NAI, SHUATS, Prayagraj, (Uttar Pradesh), India.

## Authors' contributions

This work was carried out in collaboration among all authors. All authors read and approved the final manuscript.

## Article Information

DOI: 10.9734/IJECC/2022/v12i121507

## Open Peer Review History:

This journal follows the Advanced Open Peer Review policy. Identity of the Reviewers, Editor(s) and additional Reviewers, peer review comments, different versions of the manuscript, comments of the editors, etc are available here: <https://www.sdiarticle5.com/review-history/94781>

Original Research Article

Received: 07/10/2022

Accepted: 14/12/2022

Published: 16/12/2022

## ABSTRACT

The experiment was laid out in a Randomized Block Design (RBD) with 17 treatment along with combinations of organic manure and inorganic fertilizer including control and each treatment replicated thrice. The turnip (*Brassica rapa* L.) crop in Brassicaceae family with Purple Top White Globe variety. In this study the results revealed that the application of T12 (50% NPK + 50% N through Poultry manure) significantly influenced most of the characteristics and recorded the highest values of plant height (58.94 cm, 59.85cm and 59.40 cm), number of leaves per plant (19.24, 21.03 and 20.14), leaf length (52.80 cm, 55.59 cm and 54.20 cm), leaf width (20.51 cm,

<sup>++</sup> Ph.D. Scholar;

<sup>#</sup> Professor;

<sup>†</sup> Associate Professor & Head;

\*Corresponding author: E-mail: raghuvanshisultanpur@gmail.com;

22.16cm and 21.34 cm), fresh weight of leaves (247.14, 250.84 and 248.99 g/plant) and dry weight of leaves (20.08, 22.01 and 21.05 g/plant) for the year 2018-19, 2019-20 and as pooled data respectively.

**Keywords:** Organic manure; inorganic fertilizers; vegetative growth; turnip.

## 1. INTRODUCTION

“Turnip [*Brassica rapa* L. em. Mentzg. subsp. *rapa* syn *B. compestris* L.spp. *rapifera* Sinsk, (AA, 2n=20)] is an annual or biennial plant that is cultivated worldwide as vegetable and fodder” [1,2]. “Turnip (*Brassica rapa* var. *rapifera*) is a vegetable of the cruciferous family. sometimes it is known as field mustard or turnip mustard, is a plant that is extensively grown as a leaf vegetable, root vegetable, and oilseed. Before utilising a plant medicinally, always seek the counsel of an expert. Cancer is treated using a decoction of the leaves or stems. When cooked with fat, the root is used to treat breast cancers. Skin cancer is claimed to be helped by a salve made from the flowers. Furthermore, turnip extract can help decrease uric acid and remove kidney stones. It improves visual acuity and is used to cure night blindness. Turnip syrup improves memory” [3]. “Allardice is a natural pesticide found in turnip root peelings. Turnip agriculture in India is largely limited to the northwestern states of Punjab, Haryana, Rajasthan, and Western Uttar Pradesh as an early winter season crop. Green Top, Purple Top, and 'Kenshin- Kaba' are all essential turnip cultivars for fodder production” [4].

“It is grown on 2500 acres in India, with an annual yield of 50,000 tonnes. The organics are the indigenous source of nutrients which can help in increasing production and productivity along with improvement in soil physical conditions. Use of such organic materials, which are being wasted in large amounts without proper use can help in reducing cost of cultivation, increasing productivity and improving soil as well as human and livestock health” [5-7]. “Various organic manures so far recognized in this group are green manures, farm yard manure, vermicompost, poultry manure and goat manure etc. In India, all the organic manures together have a potential to supply approximately 33 million tonnes of N,P,K per year” [8]. “Among organic manures, vermicompost is widely accepted by farmers and is produced due to the activity of earthworms. It is rich in all plant nutrients, beneficial microorganisms like N-fixers,

biologically active metabolites, particularly gibberellins, cytokinins, auxins and group B vitamins and several enzymes like lipase, cellulose, chitinase, urease, dehydrogenase and nitrogenase” (Bano et al., 1987). “The vermicompost can be applied alone or in combination with inorganic fertilizers to get better yield and quality of diverse crops. Organic farming is a method that integrates links between soil, plant, water, soil microflora and fauna. Organic farming seeks to create a healthy soil, aids in correct energy flows in the soil, crop, water, and environment, while the plant systems maintain the biological life cycle alive and aids in the maintenance of significant yield levels” [9]. In recent years, the use of organic manures for increasing agricultural yield and sustaining soil fertility and productivity has gained popularity. Organic manure improves soil structure and water retention capacity [10].

## 2. MATERIALS AND METHODS

An investigation was carried out during (2018-19 & 2019-20) at Horticulture Research Farm, Department of Horticulture, Sam Higginbottom University of Agriculture, Technology and Sciences, Prayagraj (Allahabad), U.P., India-211007. The experiment was laid out in a Randomized Block Design (RBD) with 17 treatment along with combinations of organic manure and inorganic fertilizer including control and each treatment replicated thrice. T1 - Control: RDF (100 Kg N, 80 Kg P<sub>2</sub>O<sub>5</sub> and 50 K.g. K<sub>2</sub>O /ha.), T2- (100% N through FYM), T3- (100% N through Vermicompost), T4- (100% N through Poultry manure), T5- (100 % N through Goat manure), T6- (75% NPK + 25% N through FYM), T7- (75% NPK + 25% N through Vermicompost), T8- (75% NPK + 25% N through Poultry manure), T9- (75 % NPK + 25% N through Goat manure), T10-(50% NPK + 50% N through FYM), T11- (50% NPK + 50% N through Vermicompost), T12- (50% NPK + 50% N through Poultry manure), T13- (50% NPK + 50% N through Goat manure), T14- (25% NPK + 75% N through FYM), T15- (25% NPK + 75% N through Vermicompost), T16- (25% NPK + 75% N through Poultry manure), T17- (25% NPK + 75% N through Goat manure). Source of

variables were nitrogen, phosphorus, potash, farm yard manure, Vermicompost, Poultry Manure and Goat manure. the effect of organic manure and inorganic fertilizer, vegetative growth parameters viz. plant height (cm), number of leaves per plant, leaf length (cm), leaf width (cm), fresh weight of leaves (g plant<sup>-1</sup>) and dry weight of leaves (g plant<sup>-1</sup>) were studied in the investigation. The data collected on different parameters during the course of investigation were subjected to statistical analysis as per method of analysis of variance [11]. The significance and non – significance of the treatment effect was judged with the help of F variance ratio test. Calculated F value was compared with the table value of F at 5% level of significance. If calculated value exceeds the table value, the effect was considered to be significant. The significant differences between the means were tested critical differences at 5% level of significance.

### 3. RESULTS AND DISCUSSION

The results regarding vegetative growth of turnip were statistically analyzed and are presented in Table 1a. A study of the table shows that the application of organic manures and inorganic fertilizers had significant effect on the plant height (cm), number of leaves plant<sup>-1</sup>, leaf length (cm) and leaf width (cm) of turnip in 1st year, 2nd year and when pooled.

#### 3.1 Plant Height (cm)

The result revealed that the significant maximum plant height 58.94cm, 59.85cm and 59.40cm for the year 2018-19, 2019-20 and when pooled respectively, was recorded with treatment T12(50% NPK + 50% N through Poultry manure) and followed by treatments T13 (50 % NPK + 50% N through Goat manure) 57.58cm, 58.20cm and 57.89cm at 55 days after sowing for the year 2018- 19, 2019-20 and when pooled respectively, which were statistically at par with each other and were significantly superior over T1 (control) 35.86cm, 37.14cm and 36.50cm at 55 DAS for the year 2018-19, 2019-20 and pooled data respectively. The results of the present investigation agreed with the finding of reported similarly that application of 10 to 15 t/ha of poultry manure resulted in increased height of amaranthus plants [12-15]. The increased growth parameters attributed to beneficial effect of poultry manure has been reported by [16] and [17] in radish, [18] in carrot and [19] in beetroot.

#### 3.2 Number of Leaves per Plant

The result revealed that the significant maximum number of leaves per plant 19.24, 21.03 and 20.14 for the year 2018-19, 2019-20 and when pooled respectively, was recorded with treatment T12 (50% NPK + 50% N through Poultry manure) and followed by treatments T13 (50% NPK + 50% N through Goat manure) 18.68, 20.25 and 19.47 at 55 DAS for the year 2018-19, 2019-20 and when pooled respectively, which were statistically at par with each other and were significantly superior over T1 (control) 10.83, 11.13 and 10.98 at 55 DAS for the year 2018-19, 2019-20 and pooled data respectively. These results agree with the findings. who was found significant increase in number of leaves of radish with the sole application of NPK. The results of the present investigation agreed with the finding of [20,21,22,12]. “The increased growth parameters attributed to beneficial effect of poultry manure has been reported” by [16] and [17] in radish, [18] in carrot and [19] in beetroot.

#### 3.3 Leaf Length (cm)

Among the treatments applied, treatment T12 (50% NPK + 50% N through Poultry manure)recorded significantly maximum leaf length (cm) 52.80cm, 55.59 cm and 54.20 cm for the year 2018-19, 2019-20 and as pooled data respectively, and closely followed by treatments T13 (50% NPK + 50% N through Goat manure) 51.15 cm, 53.91 cm and 52.53 cm for the year 2018-19, 2019-20 and when pooled respectively, which were statistically at par with each other and were significantly superior over T1 (control) 28.43 cm, 31.15 cm and 29.79 cm for the year 2018-19, 2019-20 and pooled data respectively. These results coincide with the previous findings of who found “maximum leaf length in radish when NPK was used, as the plants received more readily available applied nutrients, which might had increased the vegetative growth and leaf length in radish” (Islam et al. 2011). Similar results were also reported by stating an increase in cabbage leaf length, when inorganic fertilizers were used. Similar finding for the leaf length per plant were reported by [13,23,24,25] and [26] reported that the effect of application of poultry manures on growth of radish (*Raphanus sativus* L.) and pakchoi (*Brassica chinensis* L.). The increased growth parameters attributed to beneficial effect of poultry manure has been reported by [16] and [17] in radish, [18] in carrot and [19] in beetroot.

**Table 1a. Influence of different organic manures and inorganic fertilizers on vegetative growth of turnip (*Brassica rapa* L.) cv Purple Top White Globe**

Treatment	Vegetative growth											
	Plant height (cm)			Number of leaves plant-1			Leaf length (cm)			Leaf width (cm)		
	2018-19	2019-20	Pooled	2018-19	2019-20	Pooled	2018-19	2019-20	Pooled	2018-19	2019-20	Pooled
T1	35.86	37.14	36.50	10.83	11.13	10.98	28.43	31.15	29.79	8.81	9.25	9.03
T2	41.99	42.28	42.14	12.94	13.26	13.10	34.21	38.85	36.53	11.47	13.00	12.24
T3	43.05	44.34	43.70	13.57	14.14	13.86	36.26	40.47	38.37	12.64	14.23	13.44
T4	46.56	47.90	47.23	14.19	15.21	14.70	39.11	42.84	40.98	14.32	15.20	14.76
T5	44.92	45.22	45.07	13.96	14.67	14.32	38.79	41.28	40.04	13.74	15.10	14.42
T6	51.80	53.08	52.44	15.86	17.44	16.65	44.32	47.56	45.94	16.69	17.84	17.27
T7	52.72	54.12	53.42	16.20	18.15	17.18	45.68	49.90	47.79	17.26	18.81	18.04
T8	53.96	55.04	54.50	17.16	18.89	18.03	47.90	51.5	49.70	18.34	19.50	18.92
T9	53.21	54.69	53.95	16.51	18.37	17.44	46.81	50.67	48.74	17.59	19.22	18.41
T10	55.43	56.02	55.73	17.71	19.33	18.52	48.57	52.72	50.65	19.17	19.94	19.56
T11	56.81	57.39	57.10	18.10	19.69	18.90	49.95	53.13	51.54	19.48	21.12	20.30
T12	58.94	59.85	59.40	19.24	21.03	20.14	52.80	55.59	54.20	20.51	22.16	21.34
T13	57.58	58.20	57.89	18.68	20.25	19.47	51.15	53.91	52.53	19.87	21.54	20.71
T14	47.16	48.03	47.60	14.42	15.58	15.00	39.67	43.17	41.42	14.78	15.87	15.33
T15	48.00	50.29	49.15	14.68	16.05	15.37	40.84	44.63	42.74	15.29	16.75	16.02
T16	50.62	52.79	51.71	15.36	17.03	16.20	43.41	46.96	45.19	16.27	17.64	16.96
T17	49.76	51.64	50.70	14.96	16.47	15.72	41.64	45.57	43.61	15.70	17.13	16.42
<b>C.D. at 5%</b>	<b>2.25</b>	<b>2.30</b>	<b>2.27</b>	<b>0.70</b>	<b>0.76</b>	<b>0.73</b>	<b>1.94</b>	<b>2.10</b>	<b>2.02</b>	<b>0.73</b>	<b>0.79</b>	<b>0.76</b>
<b>SEd (<math>\pm</math>)</b>	<b>1.10</b>	<b>1.13</b>	<b>1.12</b>	<b>0.34</b>	<b>0.37</b>	<b>0.36</b>	<b>0.95</b>	<b>1.03</b>	<b>0.99</b>	<b>0.36</b>	<b>0.39</b>	<b>0.37</b>
<b>F-Test</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>

**Table 1b. Influence of different organic manures and inorganic fertilizers on vegetative growth of turnip (*Brassica rapa* L.) cv Purple Top White Globe**

Treatment	Vegetative growth					
	Fresh weight of leaves (g/plant)			Dry weight of leaves (g/plant)		
	2018-19	2019-20	Pooled	2018-19	2019-20	Pooled
T1	146.61	149.54	148.08	8.87	9.55	9.21
T2	152.74	156.25	154.50	11.56	12.33	11.95
T3	163.70	166.97	165.34	12.24	13.30	12.77
T4	171.01	174.94	172.98	13.70	15.08	14.39
T5	168.98	170.84	169.91	12.92	14.17	13.55
T6	191.87	197.81	194.84	16.68	17.81	17.25
T7	201.86	205.89	203.88	17.12	18.04	17.58
T8	211.97	210.29	211.13	18.51	19.62	19.07
T9	205.97	207.73	206.85	17.84	18.59	18.22
T10	216.14	222.62	219.38	18.94	19.96	19.45
T11	238.01	239.55	238.78	19.26	20.73	20.00
T12	247.14	250.84	248.99	20.08	22.01	21.05
T13	241.97	243.62	242.80	19.67	21.52	20.60
T14	172.10	178.98	175.54	13.86	15.41	14.64
T15	176.86	180.39	178.63	14.67	16.15	15.41
T16	186.93	192.35	189.64	16.34	17.74	17.04
T17	181.08	183.79	182.44	15.38	16.93	16.16
<b>C.D. at 5%</b>	<b>3.52</b>	<b>2.41</b>	<b>2.56</b>	<b>0.72</b>	<b>0.77</b>	<b>0.75</b>
<b>SEd (±)</b>	<b>1.73</b>	<b>1.19</b>	<b>1.26</b>	<b>0.35</b>	<b>0.38</b>	<b>0.37</b>
<b>F-Test</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>	<b>S</b>

### 3.4 Leaf Width (cm)

Among the treatments applied, treatment T12(50% NPK + 50% N through Poultry manure) exhibited significantly maximum leaf width (cm) 20.51cm, 22.16cm and 21.34cm for the year 2018-19, 2019-20 and as pooled data respectively, and closely followed by treatments T13 (50% NPK + 50% N through Goat manure) 19.87cm, 21.54cm and 20.71cm for the year 2018-19, 2019-20 and when pooled respectively, which were statistically at par with each other and were significantly superior over T1 (control) 8.81cm, 9.25cm and 9.03cm for the year 2018-19, 2019-20 and pooled data respectively. Similar finding for the leaf width were reported by (Chitti et al. 2018), [27,28] and [26] reported that the effect of application of poultry manures on growth of radish (*Raphanus sativus* L.) and pakchoi (*Brassica chinensis* L.). The increased growth parameters attributed to beneficial effect of poultry manure has been reported by [16] and [17] in radish, [18] in carrot and [19] in beetroot.

The results regarding vegetative growth of turnip were statistically analyzed and have been presented in Table 1b. A study of the table shows that the application of organic manures and inorganic fertilizers had significant effect on the fresh weight of leaves (g/plant) and dry

weight of leaves (g/plant) of turnip in 1st year, 2nd year and when pooled.

### 3.5 Fresh Weight of Leaves (g/plant)

Among the treatments applied, treatment T12 (50% NPK + 50% N through Poultry manure) exhibited significantly maximum fresh weight of leaves (g/plant) 247.14 g/plant, 250.84 g/plant and 248.99 g/plant for the year 2018-19, 2019-20 and as pooled data respectively, and closely followed by treatments T13 (50% NPK + 50% N through Goat manure) 241.97 g/plant, 243.62 g/plant and 242.80 g/plant for the year 2018-19, 2019-20 and when pooled respectively, which were statistically at par with each other and were significantly superior over T1 (control) 146.61 g/plant, 149.54 g/plant and 148.08 g/plant for the year 2018-19, 2019-20 and pooled data respectively. The presence of adequate amount of NPK might be the major cause of enhancing the soil fertility level which promoted plant growth thus causing an increased leaf weight plant [29-31]. The enhancements in growth parameters of radish were also reported by the application of NPK. similar finding for the fresh weight of leaves per plant were reported by [32,17,33] and [26] reported that the effect of application of poultry manures on growth of radish (*Raphanus sativus* L.) and pakchoi

(*Brassica chinensis* L.). The increased growth parameters attributed to beneficial effect of poultry manure has been reported by [16] and [17] in radish, [18] in carrot and [19] in beetroot.

### 3.6 Dry Weight of Leaves (g/plant)

Among the treatments applied, treatment T12 (50% NPK + 50% N through Poultry manure) exhibited significantly maximum dry weight of leaves (g/plant) 20.08 g/plant, 22.01 g/plant and 21.05 g/plant for the year 2018-19, 2019-20 and as pooled data respectively, and closely followed by treatments T13 (50 % NPK + 50% N through Goat manure) 19.67 g/plant, 21.52 g/plant and 20.60g/plant for the year 2018-19, 2019-20 and when pooled respectively, which were statistically at par with each other and were significantly superior over T1 (control) 8.87 g/plant, 9.55 g/plant and 9.21 g/plant for the year 2018-19, 2019-20 and pooled data respectively. The effectiveness of NPK was obvious due to greater nutrients content and their readily availability. Our results agree with the findings a significant increase in the weight of leaves plant with the application of manures and chemical fertilizers in radish. Similar results were observed by [21,34,19] and [23] reported that the effect of application of poultry manures on growth of radish (*Raphanus sativus* L.) and pakchoi (*Brassica chinensis* L.). The increased growth parameters attributed to beneficial effect of poultry manure has been reported by [13] and [14] in radish, [15] in carrot and [16] in beetroot.

## 4. CONCLUSION

Based on the current study, in both year 2018-19 and 2019-20 fiscal years. The treatment T12 (50% NPK + 50% N through Poultry manure) was determined to be the most effective in terms of turnip vegetative development. Following a few more added experiments, producers can be advised to apply this organic manure and inorganic fertilizer mix.

## COMPETING INTERESTS

Authors have declared that no competing interests exist.

## REFERENCES

1. Rakow G. Species origin and economic importance of Brassica. *Biotechnol Agric For Berlin Heidelberg*. 2004;54:3-11. DOI: 10.1007/978-3-662-06164-0\_1
2. Hammer K, Gladis TH, Laghetti G, Pignone D. The wild and the grown-remarks on Brassica. *Int J Agric Sci*. 2013; 3:453-80.
3. Khashayar P. *Alternative medicine, turnip*. Press TV, Tehran;2007.
4. Chandraprakash Y, Mishra KS, Singh KM, Sapna R, Prashant T. Effect of integrated nutrient management on growth, yield and shelf life of turnip (*Brassica rapa* L.) cv. purple top white. *The Pharm Innov J*. 2021;10(2):100-3.
5. Bano K, Kale RD. Earthworm fauna of Southern Karnataka, India. In: Veeresh GK, Rajagopal D, editors and Viraktamath CA. *Oxford and IBH, New Delhi.. Advances in management and conservation of soil fauna*. 1991; 627-34.
6. Byline M Kiran, Jilani MS, Waseem K. Impact of different organic manures and npk application on the growth and yield of turnip (*Brassica rapa* L). *Pakistan Journal of Science*. 2017;30.
7. Sandeep K, Maji S, Kumar S, Singh HD. Efficacy of organic manures on growth and yield of radish (*Raphanus sativus* L.) cv. Japanese WHITE. *Int J Plant Sci*. 2014;1(9):57-60.
8. Gaur AC, Neelkanthan S, Dargan KS. Livestock wastes characteristic and human manurial value. In: *Organic manures, pub. Indian Council of Agricultural Research*. 1990;13-95.
9. Lampkin N. In: *Organic farming, Ipswich, W.K., Farming Press Book*. 1990;801-910.
10. Kale RN, Bano K, Satyavati GP. Influence of vermicompost application on growth and yield of cereals, vegetables and ornamental plants. *Final rep. of KSCST Project N 67-04/vermin/34B*. 1991;3478;
11. Panse VG, Sukhatme PV. *Statistical methods for agricultural workers*. New Delhi: Indian Council of Agriculture Research. 1957;(p. 2ndEdn):381.
12. Kumar P, Meghwal PR, Painuli DK. Effect of organic and inorganic nutrient sources on soil health and quality of turnip. *Indian J Hortic*. 2014;71(2):222-6.
13. Mbatha AN, Ceronio GM, Coetzer GM. Response of carrot (*Daucus carota* L.) yield and quality to organic fertilizer. *S Afr J Plant Soil*. 2014;31(1):1-6. DOI: 10.1080/02571862.2013.862309

14. Verma UK, Kumar R, Kumar S, Kumar A. Integrated effect of organic manures and inorganic fertilizers on growth, yield and yield attributes of Radish cv. Kalyanpur Safed. Int J Agric Invent. 2016;1(2):158-161.
15. Okokoh SJ, Bisong BW. Effect of poultry manure and urea- N on flowering occurrence and leaf productivity of *Amranthus cruentus* in Calabar. Journ.of Apl. Sic. Environmental Management. 2011;15(1):13-5.
16. Uddain J, Chowdhury S, Rahman MJ. Efficacy of different organic manures on growth and productivity of radish (*Raphanus sativus* L.). Int J Agric Environ Biotechnol. 2010;3(2):189-93.
17. Subedi S, Srivastava A, Sharma MD, Shah SC. Effect of organic and inorganic nutrient sources on growth, yield and quality of radish (*Raphanus sativus* L.) varieties in chitwan, nepal. SAARC J Agric. 2018; 16(1):61-9.  
DOI: 10.3329/sja.v16i1.37423
18. Sylvestre H, Constance M, Esdras N, Athanase N. Effect of poultry manure and NPK on growth and yield of carrot in Rulindo district, Rwanda. Int J Novel Res Life Sci. 2015;2(1):42-8.
19. Jagadeesh C, Madhavi M, Prasad MS, Padmaja VV. Effect of organic manures on growth and yield attributes of beet root Cv. crimson globe. Int J Curr Microbiol Appl Sci. 2018; 7(11):3538-53.  
DOI: 10.20546/ijcmas.2018.711.403
20. Soheir EE, Hendawy SF, Youssef AA, Naguib NY, Hussein MS. Response of turnip (*Brassica rapa*) plants to minerals or organic fertilizers treatments. J Appl Sci Res. 2012;8(2):628-634.
21. Vijayakumari B, Sasikala V, Poornima CP. Effect of organic and inorganic manures on biometric and yield parameters of radish (*Raphanus sativus* L.). Pusaphepki. Int J Plant Sci. 2012;7(1):130-134.
22. Aisha H, Ali MR, Mahmoud S, Asmaa R, El-Desuki M. Effect of various levels of organic fertilizer and humic acid on the growth and roots quality of turnip plants (*Brassica rapa*). Current Science International. 2014;3(1):7-14.
23. Zeid HA, Wafaa HM, Abou El Seoud II, Alhadad WAA. Effect of organic materials and inorganic fertilizers on the growth, mineral composition and soil fertility of turnip plants (*Raphanus sativus* L.) grown in sandy soil. Middle East. J Agric Res. 2015;4(1):77-87.
24. Kiran M, Jilani MS, Waseem K, Sohail M. Pak J Agric Res. effect of organic manures and inorganic fertilizers on growth and yield of radish (*Raphanussativus*l). 2016; 29(4).
25. Wahocho NA, Wahocho SA, Memon N, Leghari MH, Baloch QB. Growth and yield response of turnip to various nitrogen application rates. Pac J Agri., Agril. ngg., Vet. Sci.. 2016;32(2): 143-149.
26. Zhou DM, Hao XZ, Wang YJ, Dong YH, Cang L. Copper and Zn uptake by radish and pakchoi as affected by application of livestock and poultry manures. Chemosphere. 2005;59(2):167-75.  
DOI: 10.1016/j.chemosphere.2004.11.008, PMID 15722088.
27. Dhital B, Khanal A, Sharma G, Sharma A. Effect of organic and chemical sources of nitrogen and seedling age at transplant on growth and yield of Kohlrabi. Adv Plants Agric Res. 2018;8(5):357-61.  
DOI: 10.15406/apar.2018.08.00350
28. Ingole VS, Wagh AP, Nagre PK, Bharad SG. Effect of combination of organic manure and biofertilizer for better growth and yield of beetroot (*Beta vulgaris* L.). Int J Chem Stud. 2018;6(5):1222-5.
29. Purseglove JW, Brown EG, Green CL, Robbins SRJ. Spices. 1988;7(8): 447-62.
30. Tiamiyu. Effect of sources of organic manure on growth and yields of okra (*Abelmoschus esculentus* L.). Sokoto, Nigeria: Agriculture Usmanu Danfodiyo University; 2012.
31. Thamburaj S, Singh N. Textbook of vegetable, tuber crops and spices. New Delhi: Indian Council of Agricultural Research. 2018; 161.
32. Kumar S, Dev P, Kumar J, Kumar H. Impact of organic manures and biofertilizers on the performance of radish (*Raphanus sativus* L.) HortFlora Res. Spectrum. 2018;7(2):137-40.
33. Messele B. Effects of combined application of organic – P and inorganic N fertilizers on yield of carrot (*Daucus carrota* L.). Agri

- Res & Tech: Open Access [journal]. 2016;2(2):35-9.
34. Zakir HM, Sultana MN, Saha KC. Influence of commercially available organic vs inorganic fertilizers on growth yield and quality of turnip. J Environ Sci Nat Resour. 2012;5(1): 39-45.

---

© 2022 Singh et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

*Peer-review history:*  
*The peer review history for this paper can be accessed here:*  
<https://www.sdiarticle5.com/review-history/94781>